

**NOVÁ LICHENOLOGICKÁ LITERATURA XXI.****New lichenological literature, XXII**

Zdeněk P a l i c e

*Botanický ústav AV ČR, CZ-252 43 Průhonice, e-mail: palice@ibot.cas.cz*

- Adams M. D. & Zaniewski K. (2012): Effects of recreational rock climbing and environmental variation on a sandstone cliff-face lichen community. – *Botany* 90: 253–259.
- Ahti T. & Stenroos S. (2012): New data on nomenclature, taxonomy and distribution of some species of the lichen genus *Cladonia*. – *Botanica Complutensis* 36: 31–34.
- Álvarez R. et al. (2012): Different strategies to achieve Pb-tolerance by the two *Trebouxia* algae coexisting in the lichen *Ramalina farinacea*. – *Journal of Plant Physiology* 169: 1797–1806.
- Amo de Paz G., Cubas P., Crespo A., Elix J. A. & Lumbsch H. T. (2012): Transoceanic dispersal and subsequent diversification on separate continents shaped diversity of the *Xanthoparmelia pulla* group (Ascomycota). – *PLoS ONE* 7(6): e39683.
- An K.-D. et al. (2012): Molecular phylogenetic analyses based on the nuclear rRNA genes and the intron-exon structures of the nuSSU rRNA gene in *Dictyocatenuolata alba* (anamorphic Ascomycota). – *Fungal Biology* 116: 1134–1145.
- Aptroot A. (2012): A new foliicolous *Fellhaneropsis* (*Pilocarpaceae*) from the Netherlands. – *Lichenologist* 44: 441–444.
- Aptroot A. (2012): Changes in the epiphytic flora on four *Tilia* trees in Belgium over 59 years. – *Herzogia* 25: 39–45.
- Aptroot A. & Iqbal S. H. (2012): Annotated checklist of the lichens of Pakistan, with reports of new records. – *Herzogia* 25: 211–229.
- Aptroot A. & Schumm F. (2012): A new terricolous *Trapelia* and a new *Trapeliopsis* (*Trapeliaceae*, *Baeomycetales*) from Macaronesia. – *Lichenologist* 44: 449–456.
- Aptroot A. & Schumm F. (2012): The genus *Melanophloea*, an example of convergent evolution towards polyspory. – *Lichenologist* 44: 501–509.
- Aragón G., Martínez I. & García A. (2012): Loss of epiphytic diversity along a latitudinal gradient in southern Europe. – *Science of the Total Environment* 426: 188–195.
- Arcadia L. in (2012): The publication date of fascicle 1 of volume 2 of Hoffmann's „*Descriptio et Adumbratio*“, and other notes on that work. – *Lichenologist* 44: 719–722.
- Arcadia L. in & Knudsen K. (2012): The name *Myriospora* is available for the *Acarospora smaragdula* group. – *Opuscula Philolichenum* 11: 19–25.
- Arroyo R., Serriñá E. & Araujo E. (2011): *Ramalina carminae* (Ascomycota: Ramalinaceae), a new species from Europe. – *Botanica Complutensis* 35: 5–14.
- Arup U., Fröberg L. & Pettersson L.-Å. (2011): *Placidium umbrinum* – a pyrenocarpous species new to northern Europe. – *Graphis Scripta* 23: 42–46.
- Asplund J. & Wardle D. A. (2012): Contrasting changes in palatability following senescence of the lichenized fungi *Lobaria pulmonaria* and *L. scrobiculata*. – *Fungal Ecology* 5: 710–713.
- Bates S. T. et al. (2012): A preliminary survey of lichen associated eukaryotes using pyrosequencing. – *Lichenologist* 44: 137–146.

- Beck A. & Mayr C. (2012): Nitrogen and carbon isotope variability in the green-algal lichen *Xanthoria parietina* and their implications on mycobiont–photobiont interactions. – *Ecology and Evolution* 2: 3132–3144.
- Beckett R. P., Minibayeva F. V. & Liers C. (2012): Occurrence of high tyrosinase activity in the non-Peltigeralean lichen *Dermatocarpon miniatum* (L.) W. Mann. – *Lichenologist* 44: 827–832.
- Bennett J. P., Rodriguez C. M. & Johnson C. J. (2012): Prion protein degradation by lichens of the genus *Cladonia*. – *Lichenologist* 44: 523–531.
- Berger S. A., Stamatakis A. & Lücking R. (2011): Morphology-based phylogenetic binning of the lichen genera *Graphis* and *Allographa* (Ascomycota: Graphidaceae) using molecular site weight calibration. – *Taxon* 60: 1450–1457.
- Bomble F. W. (2012): *Candelaria pacifica* und *Xanthomendoza borealis* im Aachener Raum – neu für Deutschland. – *Veröffentlichungen Bochumer Botanischer Verein* 4(1): 1–8.
- Boudreault C., Bergeron Y. & Coxson D. (2012): Epiphytic lichen colonization in regenerating black spruce forest stands of clearcut origin. – *Forest Ecology and Management* 276: 247–258.
- Braidwood D. & Ellis C. J. (2012): Bioclimatic equilibrium for lichen distributions on disjunct continental landmasses. – *Botany* 90: 1316–1325.
- Breuss O. (2012): Bemerkenswerte Flechtenfunde aus den Karnischen Alpen (Kärnten, Österreich). – *Herzogia* 25: 103–108.
- Brodo I. M. & Lendemer J. C. (2012): On the perplexing variability of reproductive modes in the genus *Ochrolechia*: Notes on *O. africana* and *O. arborea* in eastern North America. – *Opuscula Philolichenum* 11: 120–134.
- Brunialti G. et al. (2012): Can we compare lichen diversity data? A test with skilled teams. – *Ecological Indicators* 23: 509–516.
- Buck W. R. & Lendemer J. C. (2012): *Puttea* (*Pilocarpaceae*) in eastern North America. – *Opuscula Philolichenum* 11: 141–144.
- Bültmann H. (2012): The lichen syntaxa in the checklist of higher syntaxa of Europe – an overview and what we can do with them. – *Annali di Botanica* 2: 11–18.
- Bültmann H. & Daniëls F. J. A. (2012): Net photosynthesis as an alternative for relative growth rate in classifying lichens in Grime's plant strategy types. – *Bibliotheca Lichenologica* 108: 21–44.
- Burgaz A. R. & Pino-Bodas R. (2012): Notes on species of the genus *Cladonia* from Bosnia-Herzegovina and Croatia. – *Botanica Complutensis* 36: 13–18.
- Carballal R., López de Silanes M. E., Paz-Bermúdez G. & Valcárcel C. P. (2011): Flora Liquenológica Ibérica. Vol 6. Pannariaceae y *Coccocarpia* (*Coccocarpiaceae*). – Sociedad Española de Liquenología, Pontevedra.
- Cardinale M., Grube M., Vieira de Castro Jr J., Müller H. & Berg G. (2012): Bacterial taxa associated with the lung lichen *Lobaria pulmonaria* are differentially shaped by geography and habitat. – *FEMS Microbiology Letters* 194: 245–253.
- Carlsen T. et al. (2012): Species delimitation, bioclimatic range, and conservation status of the threatened lichen *Fuscopannaria confusa*. – *Lichenologist* 44: 565–575.
- Chua J. P. S. et al. (2012): Gene expression indicates a zone of heterocyst differentiation within the thallus of the cyanolichen *Pseudocyphellaria crocata*. – *New Phytologist* 196: 862–872.
- Colesie C. et al. (2012): The advantage of growing on moss: facilitative effects on photosynthetic performance and growth in the cyanobacterial lichen *Peltigera rufescens*. – *Oecologia* 169: 599–607.

- Czarnota P. (2012): Lichen protection needs natural forest disturbances – examples from some Polish Western Carpathian national parks. – In: Lipnicki L. [ed.], Lichen protection – protected lichen species, p. 53–66, Gorzów Wlkp.: Sonar Literacki.
- Czarnota P. (2012): New records of lichenized and lichenicolous fungi from Tatra National Park (W Carpathians). – In: Lipnicki L. [ed.], Lichen protection – protected lichen species, p. 287–300, Gorzów Wlkp.: Sonar Literacki.
- Czarnota P. & Guzow-Krzemińska B. (2012): ITS rDNA data confirm a delimitation of *Bacidina arnoldiana* and *B. sulphurella* and support a description of a new species within the genus *Bacidina*. – Lichenologist 44: 743–755.
- Czarnota P. & Węgrzyn M. (2012): Lichenized and lichenicolous fungi new to Babia Góra National Park (Poland, Western Carpathians). – Mycotaxon 122: 89–110.
- Dal Grande F., Widmer I., Wagner H. H. & Scheidegger C. (2012): Vertical and horizontal photobiont transmission within populations of a lichen symbiosis. – Molecular Ecology 21: 3159–3172.
- Davydov E. A. & Printzen C. (2012): Rare and noteworthy boreal lichens from the Altai Mountains (South Siberia, Russia). – Bryologist 115: 61–73.
- Davydov E. A. & Printzen C. (2012): Additions to the lichen biota of Altai Mountains (Siberia). III. – Turczaninowia 15(1): 85–91.
- Davydov E. A., Konoreva L. A., Andreev M. P., Zhdanov I. S. & Dobrysh A. A. (2012): Additions to the lichen biota of the Altai mountains (Siberia). IV. – Turczaninowia 15(3): 23–36.
- de los Ríos A. et al. (2011): Anatomical, morphological and ecophysiological strategies in *Placopsis pycnotheca* (lichenized fungi, Ascomycota) allowing rapid colonization of recently deglaciated soils. – Flora 206: 857–864.
- de Oliveira P. M. F., Timsina B. & Piercey-Normore M. D. (2012): Diversity of *Ramalina sinensis* and its photobiont in local populations. – Lichenologist 44: 649–660.
- de Vera J.-P. (2012): Lichens as survivors in space and on Mars. – Fungal Ecology 5: 472–479.
- Diederich P. et al. (2012): New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XIV. – Bulletin de la Société des naturalistes luxembourgeois 113: 95–115.
- Dillmann K. L. et al. (2012): New records, range extensions and nomenclatural innovations for lichens and lichenicolous fungi from Alaska, U.S.A. – Herzogia 25: 177–210.
- Divakar P. K. et al. (2012): Diversification of the newly recognized lichen-forming fungal lineage *Montanelia* (Parmeliaceae, Ascomycota) and its relation to key geological and climatic events. – American Journal of Botany 99: 2014–2026.
- Dobson F. S. (2011): Lichens. An illustrated guide to the British and Irish species. 6th revised and enlarged edition. – Richmond Publishing, Slough.
- Dolnik C., Schiefelbein U., Schultz M., de Bruyn U. & Richter T. (2012): New and noteworthy findings of lichenized and lichenicolous fungi from northern Germany. – Herzogia 25: 31–38.
- Dymytrova L. V., Breuss O. & Kondratyuk S. Y. (2011): *Agonimia borysthenica*, a new lichen species (Verrucariales) from Ukraine. – Österreichische Zeitschrift für Pilzkunde 20: 25–28.
- Eichler M., Cezanne R. & Teuber D. (2012): Ergänzungen zur Liste der Flechten und flechtenbewohnenden Pilze Hessens – 3. Folge. – Botanik und Naturschutz in Hessen 24: 53–69.
- Ekman S., Jonsson F. & Hermansson J. (2012): *Bacidina etayana* and *B. saxenii* new to Sweden. – Graphis Scripta 24: 14–18.

- Ellis C. J. (2012): Lichen epiphyte diversity: A species, community and trait-based review. – *Perspectives in Plant Ecology, Evolution and Systematics* 14: 131–152.
- Esslinger T. L., Morse C. A. & Leavitt S. D. (2012): A new North American species of *Hyperphyscia* (Physciaceae). – *Bryologist* 115: 31–41.
- Favero-Longo S. E. & Piervittori R. (2012): Cultivation of isidia and transplantation of adult thalli of *Xanthoparmelia tinctoria* in an abandoned asbestos mine. – *Lichenologist* 44: 840–844.
- Fazio A. T., Adler M. T., Bertoni M. D. & Maier M. S. (2012): Culture studies on the mycobiont of *Caloplaca erythrantha* (Tuck.) Zahlbr. (*Teloschistaceae*): high production of major lichen secondary metabolites. – *Lichenologist* 44: 533–542.
- Fedorenko N. M. et al. (2012): Molecular phylogeny of xanthorioid lichens (Teloschistaceae, Ascomycota), with notes on their morphology. – *Bibliotheca Lichenologica* 108: 45–64.
- Fedrowitz K., Kaasalainen U. & Rikkinen J. (2012): Geographic mosaic of symbiont selectivity in a genus of epiphytic cyanolichens. – *Ecology and Evolution* 2: 2291–2303.
- Fedrowitz K., Kuusinen M. & Snäll T. (2012): Metapopulation dynamics and future persistence of epiphytic cyanolichens in a European boreal forest ecosystem. – *Journal of Applied Ecology* 49: 493–502.
- Feurerer T. & Hertel H. (2012): The saxicolous lichens of Munich (Germany) – a preliminary evaluation. – *Bibliotheca Lichenologica* 108: 65–74.
- Flakus A. & Śliwa L. (2012): Additional information on the recently described species, *Lecanora printzenii*. – *Lichenologist* 44: 561–562.
- Fontaine K. M., Beck A., Stocker-Wörgötter E. & Piercey-Normore M. D. (2012): Photobiont relationships and phylogenetic history of *Dermatocarpon luridum* var. *luridum* and related *Dermatocarpon* species. – *Plants* 1: 39–60.
- Freitag S. et al. (2012): Metabolite profile shifts in the heathland lichen *Cladonia portentosa* in response to N deposition reveal novel biomarkers. – *Physiologia Plantarum* 146: 160–172.
- Fryday A. M. & Arcadia L. in (2012): Typification and revised basionym for *Fuscidea lygaea*, and a new name for *Amandinea lecideina*. – *Graphis Scripta* 24: 40–44.
- Fryday A. M. & Coppins B. J. (2012): New taxa, reports, and names of lichenized and lichenicolous fungi, mainly from the Scottish Highlands. – *Lichenologist* 44: 723–737.
- Gauslaa Y. & Coxson D. (2011): Interspecific and intraspecific variations in water storage in epiphytic old forest foliose lichens. – *Botany* 89: 787–798.
- Gauslaa Y., Coxson D. S. & Solhaug K. A. (2012): The paradox of higher light tolerance during desiccation in rare old forest cyanolichens than in more widespread co-occurring chloro- and cephalolichens. – *New Phytologist* 195: 812–822.
- Gauslaa Y. & Goward T. (2012): Relative growth rates of two epiphytic lichens, *Lobaria pulmonaria* and *Hypogymnia occidentalis*, transplanted within and outside of *Populus* dripzones. – *Botany* 90: 954–965.
- Gaya E. et al. (2012): Implementing a cumulative supermatrix approach for a comprehensive phylogenetic study of the Teloschistales (Pezizomycotina, Ascomycota). – *Molecular Phylogenetics and Evolution* 63: 374–387.
- Geml J. et al. (2012): Frequent circumarctic and rare transequatorial dispersals in the lichenised agaric genus *Lichenomphalia* (Hygrophoraceae, Basidiomycota). – *Fungal Biology* 116: 388–400.
- Giełwanowska I. & Olech M. (2012): New ultrastructural and physiological features of the thallus in Antarctic lichens. – *Acta Biologica Cracoviensia, ser. Bot.*, 54: 40–52.
- Giordani P. (2012): Assessing the effects of forest management on epiphytic lichens in coppiced forests using different indicators. – *Plant Biosystems* 146: 628–637.

- Giordani P., Brunialti G., Bacaro G. & Nascimbene J. (2012): Functional traits of epiphytic lichens as potential indicators of environmental conditions in forest ecosystems. – *Ecological Indicators* 18: 413–420.
- Giralt M. (2010): Flora Liquenológica Ibérica. Vol 5. Physciaceae I. – Sociedad Española de Liquenología, Barcelona.
- Goward T. (2012): Twelve readings on the lichen thallus XII. Formal propositions. – *Evansia* 29: 28–49.
- Goward T., Spribille T., Ahti T. & Hampton-Miller C. J. (2012): Four new sorediate species in the *Hypogymnia austerodes* group (lichens) from northwestern North America, with notes on thallus morphology. – *Bryologist* 115: 84–100.
- Grangeon S., Guédron S., Asta J., Sarret G. & Charlet L. (2012): Lichen and soil as indicators of an atmospheric mercury contamination in the vicinity of a chlor-alkali plant (Grenoble, France). – *Ecological Indicators* 13: 178–183.
- Groner U. (2010): Calicioid lichens and fungi in the Muota Valley, central Switzerland: high species diversity in a small area. – *Candollea* 65: 377–391.
- Grube M., Cardinale M. & Berg G. (2012): Bacteria and the Lichen Symbiosis. – In: Hock B. [ed.], *Fungal Associations, Mycota IX*, p. 363–372, Springer-Verlag, Berlin, Heidelberg.
- Grube M., Köberl M., Lackner S., Berg C. & Berg G. (2012): Host–parasite interaction and microbiome response: effects of fungal infections on the bacterial community of the Alpine lichen *Solorina crocea*. – *FEMS Microbiology Ecology* 82: 472–481.
- Grochowski P. (2012): Secondary succession with the participation of protected species of lichens in the charred areas of the Forest Inspectorate of Lubsko. – In: Lipnicki L. [ed.], *Lichen protection – protected lichen species*, p. 205–212, Gorzów Wlkp.: Sonar Literacki.
- Gruszka W. (2012): The protected and threatened lichens of the roadside trees in the Krajeńskie Lakeland. – In: Lipnicki L. [ed.], *Lichen protection – protected lichen species*, p. 277–286, Gorzów Wlkp.: Sonar Literacki.
- Hafellner J. (2012): *Caloplaca erodens* und *Protoblastenia szaferi* erstmals in Deutschland nachgewiesen. – *Herzogia* 25: 109–111.
- Hafellner J., Muggia L. & Obermayer W. (2012): *Rinodina candidogrisea*, a new sorediate species from high altitudes in the Alps. – *Bibliotheca Lichenologica* 108: 75–102.
- Hauck M. et al. (2011): Small increase in substratum pH causes the dieback of one of Europe's most common lichens, *Lecanora conizaeoides*. – *Annals of Botany* 108: 359–366.
- Hauck M. et al. (2012): Edge and land-use effects on epiphytic lichen diversity in the forest-steppe ecotone of the Mongolian Altai. – *Flora* 207: 450–458.
- Heiðmarsson S. et al. (2012): Floristic news from the NLF Iceland excursion 2009. – *Graphis Scripta* 24: 19–25.
- Henskens F. L., Green T. G. A. & Wilkins A. (2012): Cyanolichens can have both cyanobacteria and green algae in a common layer as major contributors to photosynthesis. – *Annals of Botany* 110: 555–563.
- Hertel H. (2012): Gattungseponyme bei Flechten und lichenicolen Pilzen. – *Bibliotheca Lichenologica* 107: 1–157.
- Hilmo O., Lundemo S., Holien H., Stengrundet K. & Stenøien H. K. (2012): Genetic structure in a fragmented Northern Hemisphere rainforest: large effective sizes and high connectivity among populations of the epiphytic lichen *Lobaria pulmonaria*. – *Molecular Ecology* 21: 3250–3265.
- Hodkinson B. P. (2012): An evolving phylogenetically based taxonomy of lichens and allied fungi. – *Opuscula Philolichenum* 11: 4–10.

- Hodkinson B. P., Gittel N. R., Schadt C. W. & Lutzoni F. (2012): Photoautotrophic symbiont and geography are major factors affecting highly structured and diverse bacterial communities in the lichen microbiome. – *Environmental Microbiology* 14: 147–161.
- Hodkinson B. P. & Lendemer J. C. (2012): Phylogeny and taxonomy of an enigmatic sterile lichen. – *Systematic Botany* 37: 835–844.
- Holien H. & Tønsberg T. (2012): *Biatora kodiakensis* confirmed from Europe. – *Graphis Scripta* 24: 49–52.
- Honegger R. (2012): The symbiotic phenotype of lichen-forming ascomycetes and their endo- and epibionts. – In: Hock B. [ed.], *Fungal Associations, Mycota IX*, p. 287–339, Springer-Verlag, Berlin, Heidelberg.
- Huang X., Xue J. & Guo S. (2012): Long chain n-alkanes and their carbon isotopes in lichen species from western Hubei Province: implication for geological records. – *Frontiers of Earth Science* 6: 95–100.
- Jabłońska A. (2012): Porosty z rodzaju *Porpidia* Körb. występujące w Polsce. The lichen genus *Porpidia* Körb. in Poland. – *Monographiae Botanicae* 102: 5–123.
- Jeschke M. (2012): Cryptogams in calcareous grassland restoration: perspectives for artificial vs. natural colonization. – *Tuexenia* 32: 269–279.
- Johansson O., Palmqvist K. & Olofsson J. (2012): Nitrogen deposition drives lichen community changes through differential species responses. – *Global Change Biology* 18: 2626–2635.
- John V. & Haedeke J. (2012): Neufund von *Multiclavula mucida* in Rheinland-Pfalz mit Bemerkungen zur Verbreitung und Gefährdung der Art. – *Herzogia* 25: 287–292.
- Jones T. C., Green T. G. A., Hogg I. D. & Wilkins R. J. (2012): Isolation and characterization of microsatellites in the lichen *Buellia frigida* (Physciaceae), an Antarctic endemic. – *American Journal of Botany* 99: e131–e133.
- Junttila S. & Rudd S. (2012): Characterization of a transcriptome from a non-model organism, *Cladonia rangiferina*, the grey reindeer lichen, using high-throughput next generation sequencing and EST sequence data. – *BMC Genomics* 13: 575.
- Jüriado I., Karu L. & Liira J. (2012): Habitat conditions and host tree properties affect the occurrence, abundance and fertility of the endangered lichen *Lobaria pulmonaria* in wooded meadows of Estonia. – *Lichenologist* 44: 263–276.
- Kantvilas G. & Lumbsch H. T. (2012): Reappraisal of the genera of Megalosporaceae (Teloschistales, Ascomycota). – *Australian Systematic Botany* 25: 210–216.
- Kärnefelt I., Scholz P., Seaward M. R. D. & Thell A. (2012): Lichenology in Germany: past, present and future. – *Schlechtendalia* 23: 1–90.
- Ketner-Oostra R., Aptroot A., Jungerius P. D. & Sýkora K. V. (2012): Vegetation succession and habitat restoration in Dutch lichen-rich inland drift sands. – *Tuexenia* 32: 245–268.
- Kim J. A., Hong S. G., Cheong Y. H., Koh Y. J. & Hur J.-S. (2012): A new reducing polyketide synthase gene from the lichen-forming fungus *Cladonia metacorallifera*. – *Mycologia* 104: 362–370.
- Kirschbaum U., Cezanne R., Eichler M., Hanewald K. & Windisch U. (2012): Long-term monitoring of environmental change in German towns through the use of lichens as biological indicators: comparison between the surveys of 1970, 1980, 1985, 1995, 2005 and 2010 in Wetzlar and Giessen. – *Environmental Sciences Europe* 24: 19.
- Kononenko C. P., Burkin A. A. & Tolpysheva T. Yu. (2012): Enzyme immunoassay of the secondary metabolites of micromycetes as components of lichen substances. – *Applied Biochemistry and Microbiology* 48: 71–76.

- Kosanić M., Ranković B. & Stanojković T. (2012): Antioxidant, antimicrobial, and anticancer activity of 3 Umbilicaria species. – *Journal of Food Science* 71: T20–T25.
- Kosanić M. M., Ranković B. R. & Stanojković T. P. (2012): Antioxidant, antimicrobial and anticancer activities of three *Parmelia* species. – *Journal of the Science of Food and Agriculture* 92: 1909–1916.
- Kovaleva N. M. & Ivanova G. A. (2012): Features of the biomass distribution of epiphytic lichens on scotch pine (Lower Angara Region). – *Contemporary Problems of Ecology* 5: 319–322.
- Kubiak D. (2011): Nowe stanowiska *Calicium parvum* Tibell w północnej Polsce. – *Acta Botanica Cassubica* 10: 75–81.
- Kukwa M. (2012): Typification of *Lecanora expallens* and *L. expallens* var. *conizaea* (Lecanorales, Ascomycota). – *Mycotaxon* 119: 197–200.
- Kukwa M., Łubek A., Szymczyk R. & Zalewska A. (2012): Seven lichen species new to Poland. – *Mycotaxon* 120: 105–118.
- Kukwa M. & Motiejūnaitė J. (2012): Revision of the lichen genera *Cetrelia* and *Punctelia* (Lecanorales, Ascomycota) in Lithuania with implications for their conservation. – *Herzogia* 25: 5–14.
- Kukwa M., Pietnoczko M. & Czyżewska K. (2012): The lichen family Parmeliaceae in Poland. II. The genus *Cetrelia*. – *Acta Societatis Botanicorum Poloniae* 81: 43–52.
- Kutorga E. et al. (2012): A checklist of mycobiota recorded in burnt and unburnt *Pinus mugo* plantations in the Curonian Spit (Lithuania). – *Botanica Lithuanica* 18: 66–79.
- Kuznetsova E. S., Motiejūnaitė J., Stepanchikova I. S., Himelbrant D. E. & Czarnota P. (2012): New records of lichens and allied fungi from the Leningrad Region, Russia. III. – *Folia Cryptogamica Estonica* 49: 31–37.
- Lang S. I. et al. (2012): Arctic warming on two continents has consistent negative effects on lichen diversity and mixed effects on bryophyte diversity. – *Global Change Biology* 18: 1096–1107.
- Lakatos M., Obregón A., Büdel B. & Bendix J. (2012): Midday dew – an overlooked factor enhancing photo-synthetic activity of corticolous epiphytes in a wet tropical rain forest. – *New Phytologist* 194: 245–253.
- Larsson P., Solhaug K. A. & Gauslaa Y. (2012): Seasonal partitioning of growth into biomass and area expansion in a cephalolichen and a cyanolichen of the old forest genus *Lobaria*. – *New Phytologist* 194: 991–1000.
- Leavitt S. D., Esslinger T. L., Divakar P. K. & Lumbsch H. T. (2012): Miocene and Pliocene dominated diversification of the lichen-forming fungal genus *Melanohalea* (Parmeliaceae, Ascomycota) and Pleistocene population expansions. – *BMC Evolutionary Biology* 12: 176
- Leavitt S. D., Esslinger T. L., Divakar P. K. & Lumbsch H. T. (2012): Miocene divergence, phenotypically cryptic lineages, and contrasting distribution patterns in common lichen-forming fungi (Ascomycota: Parmeliaceae). – *Biological Journal of the Linnean Society* 107: 920–937.
- Leavitt S. D., Esslinger T. L. & Lumbsch H. T. (2012): Neogene-dominated diversification in neotropical montane lichens: Dating divergence events in the lichen-forming fungal genus *Oropogon* (Parmeliaceae). – *American Journal of Botany* 99: 1764–1777.
- Lendemer J. C. (2011): A taxonomic revision of the North American species of *Lepraria* s. l. that produce divaricatic acid, with notes on the type species of the genus *L. incana*. – *Mycologia* 103: 1216–1229.
- Lindblom L. & Ekman S. (2012): RAPDs distinguish the lichens *Xanthoria aureola* and *X. parietina* in a mixed seashore rock population. – *Nova Hedwigia* 94: 279–285.

- Lipnicki L., Grochowski P. & Gruszka W. (2012): The protected and threatened lichens of the roadside trees in the Krajeńskie Lakeland. – In: Lipnicki L. [ed.], Lichen protection – protected lichen species, p. 187–196, Gorzów Wlkp.: Sonar Literacki.
- Lisov A., Zavarzina A., Zavarzin A., Demin V. & Leontievsky A. (2012): Dimeric and monomeric laccases of soil-stabilizing lichen *Solorina crocea*: Purification, properties and reactions with humic acids. – *Soil Biology and Biochemistry* 45: 161–167.
- Llop E., Pinho P., Matos P., Pereira M. J. & Branquinho C. (2012): The use of lichen functional groups as indicators of air quality in a Mediterranean urban environment. – *Ecological Indicators* 13: 215–221.
- Lobakova E. S. & Smirnov I. A. (2012): Experimental Lichenology. – In: Petre M. [ed.], *Advances in Applied Microbiology*, p. 257–276, InTech.
- Lõhmus P., Leppik E., Motiejunaite J., Suija A. & Lõhmus A. (2012): Old selectively cut forests can host rich lichen communities – lessons from an exhaustive field survey. – *Nova Hedwigia* 95: 493–515.
- Łubek A. (2012): Pionowe zróżnicowanie bioty porostów na pniu jesionu wyniosłego *Fraxinus excelsior* oraz znaczenie tego drzewa w zachowaniu różnorodności gatunkowej porostów w rezerwacie Oleszno (Przedborski Park Krajobrazowy). – *Leśne Prace Badawcze (Forest Research Papers)* 73: 23–32.
- Lücking R. (2012): Predicting species richness in tropical lichenized fungi with ‘modular’ combinations of character states. – *Biodiversity and Conservation* 21: 2341–2360.
- Lücking R. & McCune B. (2012): *Graphis pergracilis* new to North America, and a new name for *Graphis britannica* sensu Staiger auct. – *Evansia* 28: 77–84.
- Manojlovic N. T., Vasiljevic P. J., Maskovic P. Z., Juskovic M. & Bogdanovic-Dusanovic G. (2012): Chemical composition, antioxidant, and antimicrobial activities of lichen *Umbilicaria cylindrica* (L.) Delise (Umbilicariaceae). – *Evidence-Based Complementary and Alternative Medicine* 2012: 452431.
- Mansournia M. R., Wu B., Matsushita N. & Hogetsu T. (2012): Genotypic analysis of the foliose lichen *Parmotrema tinctorum* using microsatellite markers: association of mycobiont and photobiont, and their reproductive modes. – *Lichenologist* 44: 419–440.
- Mark M., Saag L., Saag A., Thell A. & Randle T. (2012): Testing morphology-based delimitation of *Vulpicida juniperinus* and *V. tubulosus* (Parmeliaceae) using three molecular markers. – *Lichenologist* 44: 757–772.
- Marmor L., Tõrra T., Saag L. & Randle T. (2012): Species richness of epiphytic lichens in coniferous forests: the effect of canopy openness. – *Annales Botanici Fennici* 49: 352–358.
- Martellos S. (2012): From a textual checklist to an information system: The case study of ITALIC, the Information System on Italian Lichens. – *Plant Biosystems* 146: 764–770.
- Martin L., Suija A., Schmeimann M., Leppik E. & Pykälä J. (2012): New Estonian records and amendments: Lichenized and lichenicolous fungi. – *Folia Cryptogamica Estonica* 49: 93–96.
- Martínez I. et al. (2012): Multiple-scale environmental modulation of lichen reproduction. – *Fungal Biology* 116: 1192–1201.
- McCarthy D. P. & Henry N. (2012): Measurement of growth in the lichen *Rhizocarpon geographicum* using a new photographic technique. – *Lichenologist* 44: 679–693.
- McIlroy de la Rosa J. P., Warke P. A. & Smith B. J. (2012): Microscale biopitting by the endolithic lichen *Verrucaria baldensis* and its proposed role in mesoscale solution basin development on limestone. – *Earth Surface Processes and Landforms* 37: 374–384.
- McMullin R. T., Bell F. W. & Newmaster S. G. (2012): The effects of triclopyr and glyphosate on lichens. – *Forest Ecology and Management* 264: 90–97.



- Meichik N. R. & Vorob'ev D. V. (2012): Chitin–glucan complex in cell walls of the *Peltigera aphthosa* lichen. – Applied Biochemistry and Microbiology 48: 307–311.
- Meinunger L. (2011): Kommentierte Checkliste der Flechten Thüringens. – Haussknechtia Beiheft 16: 1–160.
- Mežaka A., Brūmelis G. & Piterāns A. (2012): Tree and stand-scale factors affecting richness and composition of epiphytic bryophytes and lichens in deciduous woodland key habitats. – Biodiversity and Conservation 21: 3221–3241.
- Mežaka A., Brūmelis G., Piterāns A. & Printzen C. (2012): Distribution of *Lepraria* in Latvia in relation to tree substratum and deciduous forest type. – Annales Botanici Fennici 49: 162–170.
- Miao V. P. W., Manoharan S. S., Snæbjarnarson V. & Andrésón Ó. S. (2012): Expression of lec-1, a mycobiont gene encoding a galectin-like protein in the lichen *Peltigera membranacea*. – Symbiosis 57: 23–31.
- Millot M., Di Meo F., Tomasi S., Boustie J. & Trouillas P. (2012): Photoprotective capacities of lichen metabolites: A joint theoretical and experimental study. – Journal of Photochemistry and Photobiology B: Biology 111: 17–26.
- Miralles I., Jorge-Villar S. E., Cantón Y. & Domingo F. (2012): Using a mini-raman spectrometer to monitor the adaptive strategies of extremophile colonizers in arid deserts: relationships between signal strength, adaptive strategies, solar radiation, and humidity. – Astrobiology 12: 743–753.
- Miyake H. et al. (2011): Multiple dissipation components of excess light energy in dry lichen revealed by ultrafast fluorescence study at 5 K. – Photosynthesis Research 110: 39–48.
- Moberg R. (2012): Lichenes selecti exsiccati Upsalienses. Fasc. 17 & 18 (Nos 401–450). – Thunbergia 39: 1–18.
- Muñiz D. & Hladun N. (2010): Flora Liquenológica Ibérica. Vol 7. Calicioides. – Sociedad Española de Liquenología, Barcelona.
- Munzi S., Paoli L., Fiorini E. & Loppi S. (2012): Physiological response of the epiphytic lichen *Evernia prunastri* (L.) Ach. to ecologically relevant nitrogen concentrations. – Environmental Pollution 171: 25–29.
- Nakajima H. et al. (2012): Effect of copper stress on cup lichens *Cladonia humilis* and *C. subconistea* growing on copper-hyperaccumulating moss *Scopelophila cataractae* at copper-polluted sites in Japan. – Ecotoxicology and Environmental Safety 84: 341–346.
- Nascimbene J., Nimis P. L. & Benesperi R. (2012): Mature non-native black-locust (*Robinia pseudoacacia* L.) forest does not regain the lichen diversity of the natural forest. – Science of the Total Environment 421–422: 197–202.
- Nascimbene J., Thor G. & Nimis P. L. (2012): Habitat types and lichen conservation in the Alps: Perspectives from a case study in the Stelvio National Park (Italy). – Plant Biosystems 146: 428–442.
- Neumann P. & Dolnik C. (2012): *Peltigera extenuata* und andere seltene Flechten aus Schleswig-Holstein. – Kieler Notizen zur Pflanzenkunde 38: 39–47.
- Nicolardi V. et al. (2012): The adaptive response of lichens to mercury exposure involves changes in the photosynthetic machinery. – Environmental Pollution 160: 1–10.
- Nordén B., Paltto H., Claesson C. & Götmark F. (2012): Partial cutting can enhance epiphyte conservation in temperate oak-rich forests. – Forest Ecology and Management 270: 35–44.
- Oberwinkler F. (2012): Basidiolichens. – In: Hock B. [ed.], Fungal Associations, Mycota IX, p. 341–362, Springer-Verlag, Berlin, Heidelberg.
- Orange A. (2012): Semi-cryptic marine species of *Hydropunctaria* (Verrucariaceae, lichenized Ascomycota) from north-west Europe. – Lichenologist 44: 299–320.

- Orock E. A., Leavitt S. D., Fonge B. A., St. Clair L. L. & Lumbsch H. T. (2012): DNA-based identification of lichen-forming fungi: can publicly available sequence databases aid in lichen diversity inventories of Mount Cameroon (West Africa)? – *Lichenologist* 44: 833–839.
- Osyczka P. (2012): The lichens of *Cladonia*, ‘supergroup’ *Perviae*, in Poland. – *Herzogia* 25: 15–30.
- Otte V. (2012): The value of larch (*Larix* Mill.) plantations for the protection of threatened lichens in southern East Germany and adjacent areas. – In: Lipnicki L. [ed.], *Lichen protection – protected lichen species*, p. 333, Gorzów Wlkp.: Sonar Literacki.
- Otte V. & Jeremies M. (2011): Die Flechten des Baruther Schafberges und der Dubrauker Horken. – *Berichte der Naturforschenden Gesellschaft der Oberlausitz, Supplement zu Band 18*: 95–96.
- Oukarroum A., Strasser R. J. & Schansker G. (2012): Heat stress and the photosynthetic electron transport chain of the lichen *Parmelina tiliacea* (Hoffm.) Ach. in the dry and the wet state: differences and similarities with the heat stress response of higher plants. – *Photosynthesis Research* 111: 303–314.
- Ozimec S. (2011): Diversity and ecology of epiphytic and terricolous lichen mycota in Gorski kotar and Kvarner littoral (Croatia). – *Acta Biologica Slovenica* 54: 15–41.
- Pankratov T. A. (2012): Acidobacteria in microbial communities of the bog and tundra lichens. – *Microbiology* 81: 51–58.
- Paoli L. et al. (2012): Long-term biological monitoring of environmental quality around a solid waste landfill assessed with lichens. – *Environmental Pollution* 161: 70–75.
- Parnmen S., Lücking R. & Lumbsch H. T. (2012): Phylogenetic classification at generic level in the absence of distinct phylogenetic patterns of phenotypical variation: a case study in *Graphidaceae* (Ascomycota). – *PLoS ONE* 7(12): e51392.
- Parnmen S. et al. (2012): Using phylogenetic and coalescent methods to understand the species diversity in the *Cladia aggregata* complex (Ascomycota, Lecanorales). – *PLoS ONE* 7(12): e52245.
- Partl A. (2011): Lichen flora of Žumberak-Samoborsko gorje Nature Park, NW Croatia. – *Acta Botanica Croatica* 70: 99–107.
- Paukov A. G. & Teptina A. Yu. (2012): New records of lichens from Middle Urals, Russia. – *Folia Cryptogamica Estonica* 49: 39–43.
- Pérez-Ortega S. et al. (2012): Extreme phenotypic variation in *Cetraria aculeata* (lichenized Ascomycota): adaptation or incidental modification?. – *Annals of Botany* 109: 1133–1148.
- Pérez-Ortega S., Ortiz-Álvarez R., Green T. G. A. & de los Ríos A. (2012): Lichen myco- and photobiont diversity and their relationships at the edge of life (McMurdo Dry Valleys, Antarctica). – *FEMS Microbiology Ecology* 82: 429–448.
- Peršoh D. & Rambold G. (2012): Lichen-associated fungi of the Letharietum vulpinae. – *Mycological Progress* 11: 753–760.
- Pinho P. et al. (2012): Lichen functional groups as ecological indicators of the effects of land-use in Mediterranean ecosystems. – *Ecological Indicators* 15: 36–42.
- Pino-Bodas R., Ahti T., Stenroos S., Martín M. P. & Burgaz A. R. (2012): *Cladonia conista* and *C. humilis* (Cladoniaceae) are different species. – *Bibliotheca Lichenologica* 108: 161–176.
- Pino-Bodas R., Burgaz A. R., Martín M. P. & Lumbsch H. T. (2012): Species delimitations in the *Cladonia cariosa* group (Cladoniaceae, Ascomycota). – *Lichenologist* 44: 121–135.

- Pino-Bodas R., Martín M. P. & Burgaz A. R. (2012): *Cladonia subturgida* and *C. iberica* (Cladoniaceae) form a single, morphologically and chemically polymorphic species. – *Mycological Progress* 11: 269–278.
- Polovinka M. P. et al. (2012): Secondary metabolites of the lichen *Cladonia stellaris*. – *Chemistry of Natural Compounds* 48: 392–395.
- Prieto M., Martínez I., Aragón G., Gueidan C. & Lutzoni F. (2012): Molecular phylogeny of *Heteropladidium*, *Placidium*, and related catapyrenioid genera (Verrucariaceae, lichen-forming Ascomycota). – *American Journal of Botany* 99: 23–35.
- Printzen C., Cezanne R., Eichler M. & Lumbsch H. T. (2012): The genera *Aphanopsis* and *Steinia* represent basal lineages within Leotiomyceta. – *Bibliotheca Lichenologica* 108: 177–186.
- Printzen C., Fernández-Mendoza F., Muggia L., Berg G. & Grube M. (2012): Alphaproteobacterial communities in geographically distant populations of the lichen *Cetraria aculeata*. – *FEMS Microbiology Ecology* 82: 316–325.
- Pykälä J., Stepanchikova I. S., Himelbrant D. E., Kuznetsova E. S. & Alexeeva N. M. (2012): The lichen genera *Thelidium* and *Verrucaria* in the Leningrad Region (Russia). – *Folia Cryptogamica Estonica* 49: 45–57.
- Rai H., Upreti D. K. & Gupta R. K. (2012): Diversity and distribution of terricolous lichens as indicator of habitat heterogeneity and grazing induced trampling in a temperate-alpine shrub and meadow. – *Biodiversity and Conservation* 21: 97–113.
- Rajakaruna N. et al. (2012): Investigation of the importance of rock chemistry for saxicolous lichen communities of the New Idria serpentinite mass, San Benito County, California, USA. – *Lichenologist* 44: 695–714.
- Randlane T., Saag A., Martin L., Timdal E. & Nimis P. L. (2011): Epiphytic macrolichens of Estonia. – *Ülikooli Kirjastus, Tartu*.
- Rapai S. B., McMullin R. T. & Newmaster S. G. (2012): The importance of macrolichen traits and phylogeny in forest community assemblage along a high elevation gradient in southwestern British Columbia. – *Forest Ecology and Management* 274: 231–240.
- Riddell J., Padgett P. E. & Nash III T. H. (2012): Physiological responses of lichens to factorial fumigations with nitric acid and ozone. – *Environmental Pollution* 170: 202–210.
- Rivas Plata E., Lücking R. & Lumbsch H. T. (2012): A new classification for the family Graphidaceae (Ascomycota: Lecanoromycetes: Ostropales). – *Fungal Diversity* 52: 107–121.
- Rohrer A., Bilowitz P. O. & Mayrhofer H. (2012): Lichenized fungi from the Jakupica mountain range (Macedonia, FYROM). – *Herzogia* 25: 167–175.
- Roof S. & Werner A. (2011): Indirect Growth Curves remain the best choice for lichenometry: Evidence from directly measured growth rates from Svalbard. – *Arctic, Antarctic, and Alpine Research* 43: 621–631.
- Roux C. (2012): Liste des lichens et champignons lichénicoles de France. Listo de la likenoj kaj nelikeniĝintaj fungoj de Francio. – *Bulletin de la Société linnéenne de Provence* 16: 3–220.
- Ruprecht U., Brunauer G. & Printzen C. (2012): Genetic diversity of photobionts in Antarctic lecideoid lichens from an ecological viewpoint. – *Lichenologist* 44: 661–678.
- Ruprecht U., Lumbsch H. T., Brunauer G., Green T. G. A. & Türk R. (2012): Insights into the Diversity of Lecanoraceae (Lecanorales, Ascomycota) in continental Antarctica (Ross Sea region). – *Nova Hedwigia* 94: 287–306.

- Sadowsky A., Hussner A. & Ott S. (2012): Submersion tolerance in a habitat of *Stereocaulon paschale* (Stereocaulaceae) and *Cladonia stellaris* (Cladoniaceae) from the high mountain region Rondane, Norway. – *Nova Hedwigia* 94: 323–334.
- Sánchez F. J. et al. (2012): The resistance of the lichen *Circinaria gyrosa* (nom. provis.) towards simulated Mars conditions—a model test for the survival capacity of an eukaryotic extremophile. – *Planetary and Space Science* 72: 102–110.
- Sanders W. B. & de Los Ríos A. (2012): Development of thallus axes in *Usnea longissima* (Parmeliaceae, Ascomycota), a fruticose lichen showing diffuse growth. – *American Journal of Botany* 99: 998–1009.
- Savić S. & Tibell L. (2012): *Polyblastia* in Northern Europe and the adjacent Arctic. – *Symbolae Botanicae Upsalienses* 36/1: 1–69.
- Scarciglia F. et al. (2012): Role of lichens in weathering of granodiorite in the Sila uplands (Calabria, southern Italy). – *Sedimentary Geology* 280: 119–134.
- Schaper T. & Ott S. (2012): Initial developmental processes and interactions in the xerophilic lichen community of Gotland, Sweden: in situ culture experiments using the crustaceous cyanolichen *Placynthium nigrum*. – *Bibliotheca Lichenologica* 108: 187–200.
- Schei F. H. et al. (2012): Fine-scale distribution and abundance of epiphytic lichens: environmental filtering or local dispersal dynamics?. – *Journal of Vegetation Science* 23: 459–470.
- Scheidegger C., Bilovitz P. O., Werth S., Widmer I. & Mayrhofer H. (2012): Hitchhiking with forests: population genetics of the epiphytic lichen *Lobaria pulmonaria* in primeval and managed forests in southeastern Europe. – *Ecology and Evolution* 2: 2223–2240.
- Schiefelbein U., Czarnota P., Thüs H. & Kukwa M. (2012): The lichen biota of the Drawieński National Park (NW Poland, Western Pomerania). – *Folia Cryptogamica Estonica* 49: 59–71.
- Schmitt I., Otte J., Parnmen S., Sadowska-Deś A. D., Lücking R. & Lumbsch H. T. (2012): A new circumscription of the genus *Varicellaria* (Pertusariales, Ascomycota). – *MycoKeys* 4: 23–36.
- Schoch C. L. et al. (2012): Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. – *PNAS* 109: 6241–6246.
- Sérusiaux E., van den Boom P. P. G., Brand M. A., Coppins B. J. & Magain N. (2012): *Lecania falcata*, a new species from Spain, the Canary Islands and the Azores, close to *Lecania chlorotiza*. – *Lichenologist* 44: 577–590.
- Sheard J. W., Lendemer J. C., Spribille T., Thor G. & Tønsberg T. (2012): Further contributions to the genus *Rinodina* (Physciaceae, Lecanoromycetidae): two species new to science and a new record for the Canadian High Arctic. – *Herzogia* 25: 125–143.
- Shivarov V. V. & Stoykov D. Y. (2012): New records of pyrenocarpous lichenized fungi from Bulgaria. – *Mycotaxon* 121: 133–138.
- Shrestha G., Petersen S. L. & St. Clair L. L. (2012): Predicting the distribution of the air pollution sensitive lichen species *Usnea hirta*. – *Lichenologist* 44: 511–521.
- Singh K. P. & Sinha G. P. (2010): Indian lichens : an annotated checklist. – *Botanical Survey of India, Dehra Dun*.
- Śliwa L. & Krzewicka B. (2012): New records of pyrenocarpous crustose lichens in the Polish Tatra Mts and surroundings. – *Polish Botanical Journal* 57: 463–471.
- Śliwa L. & Kukwa M. (2012): New distribution data for sterile crustose lichens in the Polish Tatra Mts and its surroundings. – *Polish Botanical Journal* 57: 259–278.
- Śliwa L., Miadlikowska J., Redelings B. D., Molnar K. & Lutzoni F. (2012): Are widespread morphospecies from the *Lecanora dispersa* group (lichen-forming Ascomycota) monophyletic?. – *Bryologist* 115: 265–277.

- Smith R. J. et al. (2012): Rare inland reindeer lichens at Mima Mounds in southwest Washington State. – *North American Fungi* 7(3): 1–25.
- Solhaug K. A. & Gauslaa Y. (2012): Secondary lichen compounds as protection against excess solar radiation and herbivores. – *Progress in Botany* 73: 283–304.
- Song L., Liu W.-Y. & Nadkarni N. M. (2012): Response of non-vascular epiphytes to simulated climate change in a montane moist evergreen broad-leaved forest in southwest China. – *Biological Conservation* 152: 127–135.
- Spitale D. & Nascimbene J. (2012): Spatial structure, rock type, and local environmental conditions drive moss and lichen distribution on calcareous boulders. – *Ecological Research* 27: 633–638.
- Staniaszek-Kik M. & Szczepańska K. (2011): Lichen biota of tree fall disturbances in the Polish part of the Karkonosze Mts (West Sudety Mts). – *Časopis Slezského Muzea, Opava (A)* 60: 139–146.
- Stapper N. J. (2012): Baumflechten in Düsseldorf unter dem Einfluss von Luftverunreinigungen, Stadtklima und Klimawandel. – *Bibliotheca Lichenologica* 108: 221–240.
- Stapper N. J. (2012): Illustrierte Bestimmungshilfe zur Unterscheidung von *Candelaria concolor* und *Candelaria pacifica*. – *Archive for Lichenology* 7: 1–12.
- Stenroos S., Ahti T., Lohtander K. & Myllys L. [eds] (2011): Suomen Jäkäläopas [Lichen flora of Finland]. – *Norrinia* 21, Kasvimuseo, Luonnontieteellinen keskus-museo, Helsinki.
- Stevens C. J. et al. (2012): Terricolous lichens as indicators of nitrogen deposition: Evidence from national records. – *Ecological Indicators* 20: 196–203.
- Stoy P. C., Street L. E., Johnson A. V., Prieto-Blanco A. & Ewing S. A. (2012): Temperature, heat flux, and reflectance of common subarctic mosses and lichens under field conditions: might changes to community composition impact climate-relevant surface fluxes? – *Arctic, Antarctic, and Alpine Research* 44: 500–508.
- Sweetwood G., Lücking R., Nelsen M. P. & Aptroot A. (2012): Ascospore ontogeny and discharge in megalosporous Trypetheliaceae and Graphidaceae (Ascomycota: Dothideomycetes and Lecanoromycetes) suggest phylogenetic relationships and ecological constraints. – *Lichenologist* 44: 277–296.
- Szczepańska K. & Staniaszek-Kik M. (2012): New lichenized fungi of the Polish Karkonosze Mountains. – *Polish Botanical Journal* 57: 279–283.
- Thell A. et al. (2012): A review of the lichen family Parmeliaceae – history, phylogeny and current taxonomy. – *Nordic Journal of Botany* 30: 641–664.
- Timdal E. & Tønsberg T. (2012): *Cladonia straminea*, the correct name for *C. metacorallifera*. – *Graphis Scripta* 24: 33–35.
- Timsina B. A., Stocker-Wörgötter E. & Piercey-Normore M. D. (2012): Monophyly of some North American species of *Ramalina* and inferred polyketide synthase gene function. – *Botany* 90: 1295–1307.
- Tretiach M., Baruffo L. & Piccotto M. (2012): Effects of Mediterranean summer conditions on chlorophyll a fluorescence emission in the epiphytic lichen *Flavoparmelia soredians*: a field study. – *Plant Biosystems* 146: 171–180.
- Tretiach M., Pavanetto S., Pittao E., Sanità di Toppi L. & Piccotto M. (2012): Water availability modifies tolerance to photo-oxidative pollutants in transplants of the lichen *Flavoparmelia caperata*. – *Oecologia* 168: 589–599.
- Ungethüm K., Kison H.-U. & Stordeur R. (2011): Die epiphytischen Flechten in drei verschiedenen Laubwaldgesellschaften im Nationalpark Harz. – *Hercynia, N. F.* 44: 191–210.

- van den Boom P. & Ertz D. (2012): Lichens and lichenicolous fungi from El Hierro (Canary Islands), a survey, including five new species. – *Cryptogamie, Mycologie* 33: 59–97.
- Wang Y., Kim J. A., Cheong Y. H., Koh Y. J. & Hur J.-S. (2012): Isolation and characterization of a non-reducing polyketide synthase gene from the lichen-forming fungus *Usnea longissima*. – *Mycological Progress* 11: 75–83.
- Werth S. (2011): Biogeography and phylogeography of lichen fungi and their photobionts. – In: Fontaneto D. [ed.], *Biogeography of microorganisms. Is everything small everywhere. The Systematic Association special series, vol. 79*, p. 191–208, Cambridge, Cambridge University Press.
- Werth S. (2012): Fungal-algal interactions in *Ramalina menziesii* and its associated epiphytic lichen community. – *Lichenologist* 44: 543–560.
- Westberg M. & Clerc P. (2012): Five species of *Candelaria* and *Candelariella* (Ascomycota, Candelariales) new to Switzerland. – *MycoKeys* 3: 1–12.
- Whelan P. (2011): Lichens of Ireland. An illustrated introduction to over 250 species. – The Collins Press, Doughcloyne, Cork.
- Widmer I., Dal Grande F., Excoffier L., Holderegger R., Keller C., Mikryukov V. S. & Scheidegger C. (2012): European phylogeography of the epiphytic lichen fungus *Lobaria pulmonaria* and its green algal symbiont. – *Molecular Ecology* 21: 5827–5844.
- Wilk K. (2012): Calcicolous species of the genus *Caloplaca* in the Polish Western Carpathians. – *Polish Botanical Studies* 29: 1–91.
- Xavier B. B., Miao V. P. W., Jónsson Z. O. & Andrésson Ó. S. (2012): Mitochondrial genomes from the lichenized fungi *Peltigera membranacea* and *Peltigera malacea*: Features and phylogeny. – *Fungal Biology* 116: 802–814.
- Zalewska A. & Bohdan A. (2012): New records of *Lobaria amplissima* (Lobariaceae, Ascomycota) in Poland. – *Acta Mycologica* 47: 109–119.
- Zambare V. P. & Christopher L. P. (2012): Biopharmaceutical potential of lichens. – *Pharmaceutical Biology* 50: 778–798.
- Zambell C. B., Adams J. M., Gorring M. L. & Schwartzman D. W. (2012): Effect of lichen colonization on chemical weathering of hornblende granite as estimated by aqueous elemental flux. – *Chemical Geology* 291: 166–174.
- Zduńczyk A., Dziejczak M. & Kukwa M. (2012): The lichen genus *Pertusaria* in Poland II. Secondary chemistry of *P. flavida*. – *Folia Cryptogamica Estonica* 49: 77–81.
- Zduńczyk A. & Kukwa M. (2012): Revised distribution of the lichen species *Haematomma ochroleucum* in Poland. – *Polish Botanical Journal* 57: 473–477.
- Zimmermann D. G. & Guderley E. (2012): Flechten und flechtenbewohnende Pilze auf dem Gelände des ehemaligen Rangierbahnhofes Wuppertal-Vohwinkel (VohRang) unter besonderer Berücksichtigung ephemerer Arten. – *Jahresbericht des Naturwissenschaftlichen Vereins Wuppertal* 62: 223–240.