

NOVÁ LICHENOLOGICKÁ LITERATURA XXIII.**New lichenological literature, XXIII**

Zdeněk P a l i c e

Botanický ústav AV ČR, CZ–252 43 Průhonice, e-mail: zdenek.palice@ibot.cas.cz

- Ahti T., Stenroos S. & Moberg R. (2013): Nordic Lichen Flora Vol. 5. Cladoniaceae. – Museum of Evolution, Uppsala University.
- Allen J. L. & Lendemer J. C. (2013): Uncovering chemical variability: molecular data reveal the identity of a sterile crustose lichen from the Yukon and affirm an expanded circumscription for *Buellia griseovirens*. – North American Fungi 8(12): 1–14.
- Alphandary E. & McCune B. (2013): A new chemical spot test for miriquidic acid. – Lichenologist 45: 697–699.
- Aragón G., Belinchón R., Martínez I. & Prieto M. (2013): Estimating epiphytic lichen richness by single families in Mediterranean forests. – Forest Ecology and Management 310: 187–193.
- Arcadia L. in (2013): Lichen biogeography at the largest scales. – Lichenologist 45: 565–578.
- Arcadia L. in (2013): *Usnea dasopoga*, a name to be reinstated for *U. filipendula*, and its orthography. – Taxon 62: 604–605.
- Ardelean I. V., Keller C. & Scheidegger C. (2013): Lichen flora of Rodnei Mountains National Park (Eastern Carpathians, Romania) including new records for the Romanian mycoflora. – Folia Cryptogamica Estonica 50: 101–115
- Armstrong R. A. (2013): Development of areolae and growth of the peripheral prothallus in the crustose lichen *Rhizocarpon geographicum*: an image analysis study. – Symbiosis 60: 7–15.
- Arup U., Söchting U. & Frödén P. (2013): A new taxonomy of the family Teloschistaceae. – Nordic Journal of Botany 31: 16–83.
- Asplund J., Sandling A. & Wardle D. A. (2012): Lichen specific thallus mass and secondary compounds change across a retrogressive fire-driven chronosequence. – PLoS ONE 7(11): e49081.
- Asplund J., Bokhorst S. & Wardle D. A. (2013): Secondary compounds can reduce the soil micro-arthropod effect on lichen decomposition. – Soil Biology and Biochemistry 66: 10–16.
- Asplund J. & Wardle D. A. (2013): The impact of secondary compounds and functional characteristics on lichen palatability and decomposition. – Journal of Ecology 101: 689–700.
- Baloch E., Lumbsch H. T., Lücking R. & Wedin M. (2013): New combinations and names in *Gyalecta* for former *Belonia* and *Pachyphiale* (Ascomycota, Ostropales) species. – Lichenologist 45: 723–727.
- Bartholmeß H., Stark U. & Nobel W. (2012): Wirkungsbezogene Langzeitbeobachtung von Umweltwirkungen – 25 Jahre Flechtenkartierungen im Umweltmonitoring-Programm Esslingen/Altbach von 1983 bis 2008. – Gefahrstoffe – Reinhaltung der Luft 72: 132–138.

- Beckett R. P., Minibayeva F. V. & Liers C. (2013): On the occurrence of peroxidase and laccase activity in lichens. – *Lichenologist* 45: 277–283.
- Beckett R. P., Zavarzina A. G. & Liers C. (2013): Oxidoreductases and cellulases in lichens: Possible roles in lichen biology and soil organic matter turnover. – *Fungal Biology* 117: 431–438.
- Benatti M. N. (2012): A worldwide key for the genus *Parmelinopsis* Elix & Hale (Parmeliaceae; lichenized Ascomycetes). – *Opuscula Philolichenum* 11: 304–312.
- Bendiksby M. & Timdal E. (2013): Molecular phylogenetics and taxonomy of *Hypocenomyce* sensu lato (Ascomycota: Lecanoromycetes): Extreme polyphyly and morphological/ecological convergence. – *Taxon* 62: 940–956.
- Berger F. (2013): Zusammenfassung der in den Jahren 1990 bis 2012 in Oberösterreich publizierten Neufunde von lichenisierten Ascomyceten und nicht lichenisierten lichenicolen Pilzen. – *Beiträge zur Naturkunde Oberösterreichs* 23: 159–194.
- Bertuzzi S. & Tretiach M. (2013): Hydrogen sulphide inhibits PSII of lichen photobionts. – *Lichenologist* 45: 101–113.
- Bertuzzi S., Davies L., Power S. A. & Tretiach M. (2013): Why lichens are bad biomonitors of ozone pollution?. – *Ecological Indicators* 34: 391–397.
- Bidussi M., Gauslaa Y. & Solhaug K. A. (2013): Prolonging the hydration and active metabolism from light periods into nights substantially enhances lichen growth. – *Planta* 237: 1359–1366.
- Bidussi M., Goward T. & Gauslaa Y. (2013): Growth and secondary compound investments in the epiphytic lichens *Lobaria pulmonaria* and *Hypogymnia occidentalis* transplanted along an altitudinal gradient in British Columbia. – *Botany* 91: 621–630.
- Bilovitz P. O. & Grube M. (2012): Flechten im Ostalpenraum – sensible Zeiger von Umweltbedingungen. – *Stapfia* 96: 141–161.
- Boch S., Müller J., Prati D., Blaser S. & Fischer M. (2013): Up in the tree – The overlooked richness of bryophytes and lichens in tree crowns. – *PLoS ONE* 8(12): e84913.
- Boch S., Prati D., Hessenmöller D., Schulze E.-D. & Fischer M. (2013): Richness of lichen species, especially of threatened ones, is promoted by management methods furthering stand continuity. – *PLoS ONE* 8(1): e55461.
- Boltersdorf S. & Werner W. (2013): Source attribution of agriculture-related deposition by using total nitrogen and $\delta^{15}\text{N}$ in epiphytic lichen tissue, bark and deposition water samples in Germany. – *Isotopes in Environmental and Health Studies* 49: 197–218.
- Bosch-Roig P., Barca D., Crisci G. M. & Lalli C. (2013): Lichens as bioindicators of atmospheric heavy metal deposition in Valencia, Spain. – *Journal of Atmospheric Chemistry* 70: 373–388.
- Boudreault C., Coxson D., Bergeron Y., Stevenson S. & Bouchard M. (2013): Do forests treated by partial cutting provide growth conditions similar to old-growth forests for epiphytic lichens?. – *Biological Conservation*, 159: 458–467.
- Boudreault C., Zouaoui S., Drapeau P., Bergeron Y. & Stevenson S. (2013): Canopy openings created by partial cutting increase growth rates and maintain the cover of three *Cladonia* species in the Canadian boreal forest. – *Forest Ecology and Management* 304: 473–481.
- Breuss O. (2013): Neue Flechtenfunde aus Frankreich. – *Herzogia* 26: 417–422.
- Breuss O. & Berger F. (2012): Die Validierung von *Verrucaria finitima* und Bemerkungen über den Formenkreis von *Verrucaria tristis* (lichenisierte Ascomyceten, Verrucariaceae). – *Österreichische Zeitschrift für Pilzkunde* 21: 117–126.

- Brisdelli F. et al. (2013): Cytotoxic activity and antioxidant capacity of purified lichen metabolites: an in vitro study. – *Phytotherapy Research* 27: 431–437.
- Brodo I. M., Freebury C. & Alfonso N. (2013): Notes on the lichens *Physcia aipolia* and *P. alnophila* in North America. – *Evansia* 30: 110–119.
- Brunialti G., Ravera S. & Frati L. (2013): Mediterranean old-growth forests: the role of forest type in the conservation of epiphytic lichens. – *Nova Hedwigia* 96: 367–381.
- Burgaz A. R. (2011): Flora Liquenológica Ibérica. Vol 8. Peltigerales: Massalongiaceae y Placynthiaceae. – Sociedad Española de Liquenología, Madrid.
- Burkin A. A. & Kononenko G. P. (2013): Peculiarities of mycotoxin accumulation in lichens. – *Applied Biochemistry and Microbiology* 49: 521–528.
- Calviño-Cancela M., López de Silanes M. E., Rubido-Bará M. & Uribarri J. (2013): The potential role of tree plantations in providing habitat for lichen epiphytes. – *Forest Ecology and Management* 291: 386–395.
- Carvalho P. (2012): Flora Liquenológica Ibérica. Vol 10. *Collema* (Coccocarpiaceae). – Sociedad Española de Liquenología, Pontevedra.
- Casanovas P., Lynch H. J. & Fagan W. F. (2013): Multi-scale patterns of moss and lichen richness on the Antarctic Peninsula. – *Ecography* 36: 209–219.
- Cetin H. et al. (2012): Larvicidal activity of some secondary lichen metabolites against the mosquito *Culiseta longiareolata* Macquart (Diptera: Culicidae). – *Natural Product Research* 26: 350–355.
- Cezanne R., Eichler M. & Windisch U. (2012): Flechten und flechtenbewohnende Pilze von 25 Untersuchungsstationen in Bayern. – *Hoppea* 73: 153–190.
- Christensen S. N. & Alstrup V. (2013): Notes on epilithic, epigeic and muscicolous lichens and lichenicolous fungi from rock outcrops in the mountains of northern Greece. – *Mycobiota* 1: 25–50.
- Cornejo C. & Scheidegger C. (2013): New morphological aspects of cephalodium formation in the lichen *Lobaria pulmonaria* (Lecanorales, Ascomycota). – *Lichenologist* 45: 77–87.
- Cornejo C. & Scheidegger C. (2013): Morphological aspects associated with repair and regeneration in *Lobaria pulmonaria* and *L. amplissima* (Peltigerales, Ascomycota). – *Lichenologist* 45: 285–289.
- Cuzman O. A. et al. (2013): Evaluation of treatments efficiency against lichens growing on monumental stones by electrical conductivity. – *International Biodeterioration & Biodegradation* 84: 314–321.
- Dal-Forno M. et al. (2013): Starting from scratch: Evolution of the lichen thallus in the basidiolichen *Dictyonema* (Agaricales: Hygrophoraceae). – *Fungal Biology* 117: 584–598.
- del Campo E. M. et al. (2013): The genetic structure of the cosmopolitan three-partner lichen *Ramalina farinacea* evidences the concerted diversification of symbionts. – *FEMS Microbiology Ecology* 83: 310–323.
- Del-Prado R. et al. (2013): Molecular phylogeny and historical biogeography of the lichen-forming fungal genus *Flavoparmelia* (Ascomycota: Parmeliaceae). – *Taxon* 62: 928–939.
- Dittrich S. et al. (2013): Response of ground vegetation and epiphyte diversity to natural age dynamics in a Central European mountain spruce forest. – *Journal of Vegetation Science* 24: 675–687.

- Dittrich S. et al. (2013): Separating forest continuity from tree age effects on plant diversity in the ground and epiphyte vegetation of a Central European mountain spruce forest. – *Flora* 208: 238–246.
- Divakar P. K. et al. (2013): A molecular perspective on generic concepts in the *Hypotrachyna* clade (Parmeliaceae, Ascomycota). – *Phytotaxa* 132: 21–38.
- Divakar P. K., Kauff F., Crespo A., Leavitt S. D. & Lumbsch H. T. (2013): Understanding phenotypical character evolution in parmelioid lichenized fungi (Parmeliaceae, Ascomycota). – *PLoS ONE* 8(11): e83115.
- Dobson F. S. (2013): A field key to lichens on trees. – Published by the author.
- Dymytrova L., Nadyeina O., Naumovych A., Keller C. & Scheidegger C. (2013): Primeval beech forests of Ukrainian Carpathians are sanctuaries for rare and endangered epiphytic lichens. – *Herzogia* 26: 73–89.
- Edwards D., Axe L. & Honegger R. (2013): Contributions to the diversity in cryptogamic covers in the mid-Palaeozoic: *Nematothallus* revisited. – *Botanical Journal of the Linnean Society* 173: 505–534.
- Ekman S. & Blaalid R. (2011): The devil in the details: Interactions between the branch-length prior and likelihood model affect node support and branch lengths in the phylogeny of the Psoraceae. – *Systematic Biology* 60: 541–561.
- Ellis C. J. (2013): A risk-based model of climate change threat: hazard, exposure, and vulnerability in the ecology of lichen epiphytes. – *Botany* 91: 1–11.
- Ellis C. J. & Ellis S. C. (2013): Signatures of autogenic epiphyte succession for an aspen chronosequence. – *Journal of Vegetation Science* 24: 688–701.
- Ellis C. J., Coppins B. J., Eaton S. & Simkin J. (2013): Implications of ash dieback for associated epiphytes. – *Conservation Biology* 27: 899–901.
- Ertz D. & Tehler A. (2011): The phylogeny of Arthoniales (Pezizomycotina) inferred from nucLSU and RPB2 sequences. – *Fungal Diversity* 49: 47–71.
- Ertz D., Fischer E., Killmann D., Razafindrahaja T. & Sérusiaux E. (2013): *Savoronala*, a new genus of Malmideaceae (Lecanorales) from Madagascar with stipes producing sporodochia. – *Mycological Progress* 12: 645–656.
- Evju M. & Bruteig I. E. (2013): Lichen community change over a 15-year time period: effects of climate and pollution. – *Lichenologist* 45: 35–50.
- Favero-Longo S. E. et al. (2013): Lichen deterioration of asbestos and asbestiform minerals of serpentinite rocks in Western Alps. – *International Biodeterioration & Biodegradation* 84: 342–350.
- Fernández-Brime S., Llimona X., Lutzoni F. & Gaya E. (2013): Phylogenetic study of *Diploschistes* (lichen-forming Ascomycota: Ostropales: Graphidaceae), based on morphological, chemical, and molecular data. – *Taxon* 62: 267–280.
- Fernández-Mendoza F. & Printzen C. (2013): Pleistocene expansion of the bipolar lichen *Cetraria aculeata* into the Southern hemisphere. – *Molecular Ecology* 22: 1961–1983.
- Flakus A. & Farkas E. (2013): A contribution to the taxonomy of *Lyromma* (Lyrommataceae, lichenized Ascomycota) with a species key. – *Mycotaxon* 124: 127–141.
- Frahm J.-P. (2013): Contents of amino acids and osmotic values of epiphytic lichens as indicators for regional atmospheric nitrogen loads. – *Archive for Lichenology* 9: 1–11.
- Gauslaa Y., Bidussi M., Solhaug K. A., Asplund J. & Larsson P. (2013): Seasonal and spatial variation in carbon based secondary compounds in green algal and cyanobacterial members of the epiphytic lichen genus *Lobaria*. – *Phytochemistry* 94: 91–98.

- Gazzano C., Favero-Longo S. E., Iacomussi P. & Piervittori R. (2013): Biocidal effect of lichen secondary metabolites against rock-dwelling microcolonial fungi, cyanobacteria and green algae. – *International Biodeterioration & Biodegradation* 84: 300–306.
- Ghate N. B. et al. (2013): An antioxidant extract of tropical lichen, *Parmotrema reticulatum*, induces cell cycle arrest and apoptosis in breast carcinoma cell line MCF-7. – *PLoS ONE* 8(12): e82293.
- Gheza G., Assini S. & Valucia-Passadore M. (2013): Contribution to the knowledge of lichen flora of inland sand dunes in the western Po Plain (N Italy). – *Plant Biosystems* 147: 115–124.
- Gustafsson L., Fedrowitz K. & Hazell P. (2013): Survival and vitality of a macrolichen 14 years after transplantation on aspen trees retained at clearcutting. – *Forest Ecology and Management* 291: 436–441.
- Guzow-Krzemińska B. & Stocker-Wörgötter E. (2013): In vitro culturing and resynthesis of the mycobiont *Protoparmeliopsis muralis* with algal bionts. – *Lichenologist* 45: 65–76.
- Guzow-Krzemińska B. & Stocker-Wörgötter E. (2013): Development of microsatellite markers in *Protoparmeliopsis muralis* (lichenized Ascomycete) – a common lichen species. – *Lichenologist* 45: 791–798.
- Hauck M., Böning J., Jacob M., Dittrich S., Feussner I. & Leuschner C. (2013): Lichen substance concentrations in the lichen *Hypogymnia physodes* are correlated with heavy metal concentrations in the substratum. – *Environmental and Experimental Botany* 85: 58–63.
- Hauck M., de Bruyn U. & Leuschner C. (2013): Dramatic diversity losses in epiphytic lichens in temperate broad-leaved forests during the last 150 years. – *Biological Conservation* 157: 136–145.
- Hauck M. & Lkhagvadorj D. (2013): Epiphytic lichens as indicators of grazing pressure in the Mongolian forest-steppe. – *Ecological Indicators* 32: 82–88.
- Hauck M., Tønsberg T., Mayrhofer H. & Breuss O. (2013): Lichen-forming and lichenicolous fungi new to Kazakhstan. – *Herzogia* 26: 103–116.
- Hertel H., Scholz P., Krempelhuber D. & Krempelhuber A. (2013): „Mit der Kugelbüchse in der Hand und Hammer, Meißel nebst Lupe im Bergsacke“. Der Flechtenforscher August von Krempelhuber im Spiegel eigener Briefe. – *Berichte der Bayerischen Botanischen Gesellschaft* 83: 5–56.
- Hilmo O., Gauslaa Y., Rocha L., Lindmo S. & Holien H. (2013): Vertical gradients in population characteristics of canopy lichens in boreal rainforests of Norway. – *Botany* 91: 814–821.
- Himelbrant D. E., Motiejūnaitė J., Pykälä J., Schiefelbein U. & Stepanchikova I. S. (2013): New records of lichens and allied fungi from the Leningrad Region, Russia. IV. – *Folia Cryptogamica Estonica* 50: 23–31.
- Hodkinson B. P. & Lendemer J. C. (2013): Next-generation sequencing reveals sterile crustose lichen phylogeny. – *Mycosphere* 4: 1028–1039.
- Honegger R., Axe L. & Edwards D. (2013): Bacterial epibionts and endolichenic actinobacteria and fungi in the Lower Devonian lichen *Chlorolichenomycites salopenensis*. – *Fungal Biology* 117: 512–518.
- Honegger R., Edwards D. & Axe L. (2013): The earliest records of internally stratified cyanobacterial and algal lichens from the Lower Devonian of the Welsh Borderland. – *New Phytologist* 197: 264–275.

- Humphries R. N. (2013): Case study: the contribution of active surface mines in the conservation of lichen communities in the South Wales Coalfield, United Kingdom. – *Journal American Society of Mining and Reclamation* 2: 80–98.
- Hutten M. et al. (2013): Lichens and lichenicolous fungi of Yosemite National Park, California. – *North American Fungi* 8(11): 1–47.
- Johansson V., Ranius T. & Snäll T. (2013): Epiphyte metapopulation persistence after drastic habitat decline and low tree regeneration: time-lags and effects of conservation actions. – *Journal of Applied Ecology* 50: 414–422.
- Johansson V., Snäll T. & Ranius T. (2013): Estimates of connectivity reveal non-equilibrium epiphyte occurrence patterns almost 180 years after habitat decline. – *Oecologia* 172: 607–615.
- Jones T. C., Hogg I. D., Wilkins R. J. & Green T. G. A. (2013): Photobiont selectivity for lichens and evidence for a possible glacial refugium in the Ross Sea Region, Antarctica. – *Polar Biology* 36: 767–774.
- Jönsson M. T. & Thor G. (2012): Estimating coextinction risks from epidemic tree death: affiliate lichen communities among diseased host tree populations of *Fraxinus excelsior*. – *PLoS ONE* 7(9): e45701.
- Junttila S., Laiho A., Gyenesei A. & Rudd S. (2013): Whole transcriptome characterization of the effects of dehydration and rehydration on *Cladonia rangiferina*, the grey reindeer lichen. – *BMC Genomics* 14: 870
- Kaasalainen U., Fewer D. P., Jokela J., Wahlsten M., Sivonen K. & Rikkinen J. (2013): Lichen species identity and diversity of cyanobacterial toxins in symbiosis. – *New Phytologist* 198: 647–651.
- Kantvilas G. & Elix J. A. (2013): The lichen genus *Lecidella* (Lecanoraceae), with special reference to the Tasmanian species. – *Muelleria* 31: 31–47.
- Klepsland J. T. (2013): *Nephroma helveticum* and *N. tangeriense* new to Norway. – *Graphis Scripta* 25: 33–38.
- La Greca S. & Lumbsch H. T. (2013): Taxonomic investigations of *Lecanora strobilina* and *L. symmicta* (Lecanoraceae, Lecanorales) in northeastern North America. – *Bryologist* 116: 287–295.
- Leavitt S. D. et al. (2013): Multilocus phylogeny of the lichen-forming fungal genus *Melanohalea* (Parmeliaceae, Ascomycota): Insights on diversity, distributions, and a comparison of species tree and concatenated topologies. – *Molecular Phylogenetics and Evolution* 66: 138–152.
- Leavitt S. D. et al. (2013): DNA barcode identification of lichen-forming fungal species in the *Rhizoplaca melanophthalma* species-complex (Lecanorales, Lecanoraceae), including five new species. – *MycKeys* 7: 1–22.
- Leavitt S. D., Lumbsch H. T. & St. Clair L. L. (2013): Contrasting demographic histories of two species in the lichen-forming fungal genus *Xanthomendoza* (Teloschistaceae, Ascomycota). – *Bryologist* 116: 337–349.
- Leavitt S. D., Nelsen M. P., Lumbsch H. T., Johnson L. A. & St. Clair L. L. (2013): Symbiont flexibility in subalpine rock shield lichen communities in the Southwestern USA. – *Bryologist* 116: 149–161.
- Lendemer J. C. (2013): A monograph of the crustose members of the genus *Lepraria* Ach. s. str. (Stereocaulaceae, Lichenized Ascomycetes) in North America north of Mexico. – *Opuscula Philolichenum* 12: 27–141.

- Lendemer J. C., Harris R. C. & Tripp E. A. (2013): The lichens and allied fungi of Great Smoky Mountains National Park, an annotated checklist with comprehensive keys. – *Memoirs of the New York Botanical Garden* 104: i–vii, 1–156.
- Lendemer J. C. & Hodkinson B. P. (2013): A radical shift in the taxonomy of *Lepraria* s. l.: Molecular and morphological studies shed new light on the evolution of asexuality and lichen growth form diversification. – *Mycologia* 105: 994–1018.
- Leppik E., Jüriado I., Suija A. & Liira J. (2013): The conservation of ground layer lichen communities in alvar grasslands and the relevance of substitution habitats. – *Biodiversity and Conservation*, 22: 591–614.
- Lindblom L. & Söchting U. (2013): Genetic diversity of the photobiont of the bipolar lichen-forming ascomycete *Xanthomendoza borealis*. – *Herzogia* 26: 307–322.
- Lohézic-Le Dévéhat F., Legouin B., Couteau C., Boustie J. & Coiffard L. (2013): Lichenic extracts and metabolites as UV filters. – *Journal of Photochemistry and Photobiology B: Biology* 120: 17–28.
- Lõhmus A., Suija A. & Lõhmus P. (2013): Intensive local surveys can complement rapid survey techniques to provide insights into the population size and ecology of lichenised fungi. – *Fungal Ecology* 6: 449–452.
- Lord J. M. et al. (2013): Rediscovery of pycnidia in *Thamnolia vermicularis*: implications for chemotype occurrence and distribution. – *Lichenologist* 45: 397–411.
- Łubek A. & Jaroszewicz B. (2012): New, rare and noteworthy species of lichens and lichenicolous fungi from Białowieża Forest. – *Polish Journal of Natural Sciences* 27: 275–287.
- Lücking R., Tehler A., Bungartz F., Plata E. R. & Lumbsch H. T. (2013): Journey from the West: Did tropical Graphidaceae (lichenized Ascomycota: Ostropales) evolve from a saxicolous ancestor along the American Pacific coast?. – *American Journal of Botany* 100: 844–856.
- Lücking R. et al. (2013): Ten new species of lichenized Basidiomycota in the genera *Dictyonema* and *Cora* (Agaricales: Hygrophoraceae), with a key to all accepted genera and species in the *Dictyonema* clade. – *Phytotaxa* 139: 1–38.
- Lundström J., Jonsson F., Perhans K. & Gustafsson L. (2013): Lichen species richness on retained aspens increases with time since clear-cutting. – *Forest Ecology and Management* 293: 49–56.
- MacDonald A. M., Coxson D. & Björk C. (2013): A framework for climate biomonitoring with lichens in British Columbia's inland temperate rainforest. – *Journal of Ecosystems and Management* 114: 1–13.
- Máguas C., Pinho P., Branquinho C., Hartard B. & Lakatos M. (2013): Carbon-Water-Nitrogen relationships between lichens and the atmosphere: Tools to understand metabolism and ecosystem change. – *MycoKeys* 6: 95–106.
- Manoharan S. S., Miao V. P. W. & Andrésón Ó. S. (2012): LEC-2, a highly variable lectin in the lichen *Peltigera membranacea*. – *Symbiosis* 58: 91–98.
- Marmor L., Törre T., Saag L., Leppik E. & Randle T. (2013): Lichens on *Picea abies* and *Pinus sylvestris* – from tree bottom to the top. – *Lichenologist* 45: 51–63.
- Marques J., Schultz M. & Paz-Bermúdez G. (2013): A *Peltula* Nyl. diversity hotspot in north-east Portugal, with one species new to science and three species new to mainland Europe. – *Lichenologist* 45: 483–496.
- Matsunaga K. K. S., Stockey R. A. & Tomescu A. M. F. (2013): *Honeggeriella complexa* gen. et sp. nov., a heteromerous lichen from the Lower Cretaceous of Vancouver Island (British Columbia, Canada). – *American Journal of Botany* 100: 450–459

- Mayer W. et al. (2013): Significant decrease in epiphytic lichen diversity in a remote area in the European Alps, Austria. – *Basic and Applied Ecology* 14: 396–403.
- Mayrhofer H. & Coppins B. J. (2013): Remarkable records of lichenized fungi from Slovenia. – *Herzogia* 26: 201–206.
- McDonald T. R., Gaya E. & Lutzoni F. (2013): Twenty-five cultures of lichenizing fungi available for experimental studies on symbiotic systems. – *Symbiosis* 59: 165–171.
- McDonald T. R., Mueller O., Dietrich F. S. & Lutzoni F. (2013): High-throughput genome sequencing of lichenizing fungi to assess gene loss in the ammonium transporter/ammonia permease gene family. – *BMC Genomics* 14: 225.
- McIlroy de la Rosa J. P., Casares Porcel M. & Warke P. A. (2013): Mapping stone surface temperature fluctuations: Implications for lichen distribution and biomodification on historic stone surfaces. – *Journal of Cultural Heritage* 14: 346–353.
- McMullin R. T., Thompson I. D. & Newmaster S. G. (2013): Lichen conservation in heavily managed boreal forests. – *Conservation Biology* 27: 1020–1030.
- Meeßen J. & Ott S. (2013): Recognition mechanisms during the pre-contact state of lichens: I. Mycobiont-photobiont interactions of the mycobiont of *Fulgensia bracteata*. – *Symbiosis* 59: 121–130.
- Meeßen J., Eppenstein S. & Ott S. (2013): Recognition mechanisms during the pre-contact state of lichens: II. Influence of algal exudates and ribitol on the response of the mycobiont of *Fulgensia bracteata*. – *Symbiosis* 59: 131–143.
- Meeßen J. et al. (2013): Extremotolerance and resistance of lichens: Comparative studies on five species used in astrobiological research I. Morphological and anatomical characteristics. – *Origins of Life and Evolution of Biospheres* 43: 283–303.
- Miadlikowska J., Kauff F., Högnabba F., Oliver J. C., Molnár K., Fraker E., Gaya E., Hafelner J., Hofstetter V., Gueidan C., Otálora M. A. G., Hodkinson B., Kukwa M., Lücking R., Björk C., Sipman H. J. M., Burgaz A. R., Thell A., Passo A., Myllys L., Goward T., Fernández-Brime S., Hestmark G., Lendemer J., Lumbsch H. T., Schmall M, Schoch C. L., Sérusiaux E., Maddison D. R., Arnold A. E., Lutzoni F. & Stenroos S. (2014): A multigene phylogenetic synthesis for the class Lecanoromycetes (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. *Molecular Phylogenetics and Evolution* 79: 132–168.
- Molnár K., Lőkös L., Schrett-Major A. & Farkas E. (2012): Molecular genetic analysis of *Xanthoparmelia pulvinaris* (Ascomycota, Lecanorales, Parmeliaceae). – *Acta Botanica Hungarica* 54: 125–130.
- Moncada B., Lücking R. & Betancourt-Macuase L. (2013): Phylogeny of the Lobariaceae (lichenized Ascomycota: Peltigerales), with a reappraisal of the genus *Lobariella*. – *Lichenologist* 45: 203–263.
- Muggia L., Gueidan C., Knudsen K., Perlmutter G. & Grube M. (2013): The lichen connections of black fungi. – *Mycopathologia* 175: 523–535.
- Muggia L., Klug B., Berg G. & Grube M. (2013): Localization of bacteria in lichens from Alpine soil crusts by fluorescence in situ hybridization. – *Applied Soil Ecology* 68: 20–25.
- Nadyeina O., Lutsak T., Blum O., Grakhov V. & Scheidegger C. (2013): *Cetraria steppae* Savicz is conspecific with *Cetraria aculeata* (Schreb.) Fr. according to morphology, secondary chemistry and ecology. – *Lichenologist* 45: 841–856.
- Nakajima H., Yamamoto Y., Yoshitani A. & Itoh K. (2013): Effect of metal stress on photosynthetic pigments in the Cu-hyperaccumulating lichens *Cladonia humilis* and

- Stereocaulon japonicum* growing in Cu-polluted sites in Japan. – *Ecotoxicology and Environmental Safety* 97: 154–159.
- Nardini A., Marchetto A. & Tretiach M. (2013): Water relation parameters of six *Peltigera* species correlate with their habitat preferences. – *Fungal Ecology* 6: 397–407.
- Nascimbene J., Dainese M. & Sitzia T. (2013): Contrasting responses of epiphytic and dead wood-dwelling lichen diversity to forest management abandonment in silver fir mature woodlands. – *Forest Ecology and Management* 289: 325–332.
- Nascimbene J., Marini L. & Ódor P. (2012): Drivers of lichen species richness at multiple spatial scales in temperate forests. – *Plant Ecology & Diversity* 5: 355–363.
- Nascimbene J., Nimis P. L. & Ravera S. (2013): Evaluating the conservation status of epiphytic lichens of Italy: A red list. – *Plant Biosystems* 147: 898–904.
- Nascimbene J., Nimis P. L. & Thüs H. (2013): Lichens as bioindicators in freshwater ecosystems – challenges and perspectives. – *Annali di Botanica* 3: 45–50.
- Nascimbene J., Thor G. & Nimis P. L. (2013): Effects of forest management on epiphytic lichens in temperate deciduous forests of Europe – A review. – *Forest Ecology and Management* 298: 27–38.
- Nascimbene J. et al. (2013): Patterns and drivers of β -diversity and similarity of *Lobaria pulmonaria* communities in Italian forests. – *Journal of Ecology* 101: 493–505.
- Nelson P. R., Roland C., Macander M. J. & McCune B. (2013): Detecting continuous lichen abundance for mapping winter caribou forage at landscape spatial scales. – *Remote Sensing of Environment* 137: 43–54.
- Neuwirth G. (2013): Der *Graphis scripta*-Komplex in Oberösterreich. Ein erster Überblick zur Verbreitung neuer Morphotypen der „Schriftflechte“. – *Stapfia* 99: 61–74.
- Nguyen K.-H., Chollet-Krugler M., Gouault N. & Tomasi S. (2013): UV-protectant metabolites from lichens and their symbiotic partners. – *Natural Product Reports* 30: 1490–1508.
- Nyati S., Bhattacharya D., Werth S. & Honegger R. (2013): Phylogenetic analysis of LSU and SSU rDNA group I introns of lichen photobionts associated with the genera *Xanthoria* and *Xanthomendoza* (Teloschistaceae, lichenized Ascomycetes). – *Journal of Phycology* 49: 1154–1166.
- Nyati S., Werth S. & Honegger R. (2013): Genetic diversity of sterile cultured *Trebouxia* photobionts associated with the lichen-forming fungus *Xanthoria parietina* visualized with RAPD-PCR fingerprinting techniques. – *Lichenologist* 45: 825–840.
- O'Brien H. E., Miadlikowska J. & Lutzoni F. (2013): Assessing population structure and host specialization in lichenized cyanobacteria. – *New Phytologist* 198: 557–566.
- Ódor P., Király I., Tinya F., Bortignon F. & Nascimbene J. (2013): Patterns and drivers of species composition of epiphytic bryophytes and lichens in managed temperate forests. – *Forest Ecology and Management* 306: 256–265.
- Oetl S. K. et al. (2013): Imbricarinic acid and perlatolic acid: multi-targeting anti-inflammatory depsides from *Cetrelia monachorum*. – *PLoS ONE* 8(10): e76929.
- Ohmura Y., Hosaka K., Kasuya T., Abe J. P. & Kakishima M. (2013): Activity concentrations of radionuclides in lichens following the Fukushima nuclear accident. – *Lichenologist* 45: 685–689.
- Orange A. (2013): Four new species of *Verrucaria* (Verrucariaceae, lichenized Ascomycota) from freshwater habitats in Europe. – *Lichenologist* 45: 305–322.
- Osyczka P. (2013): A morphometric evaluation of the *Cladonia chlorophaea* group and allied taxa (Cladoniaceae, Ascomycota). – *Herzogia* 26: 49–64.

- Osyczka P. & Rola K. (2013): Phenotypic plasticity of primary thallus in selected *Cladonia* species (lichenized Ascomycota: Cladoniaceae). – *Biologia* 68: 365–372.
- Osyczka P. & Rola K. (2013): *Cladonia* lichens as the most effective and essential pioneers in strongly contaminated slag dumps. – *Central European Journal of Biology* 8: 876–887.
- Osyczka P. & Rola K. (2013): Response of the lichen *Cladonia rei* Schaer. to strong heavy metal contamination of the substrate. – *Environmental Science and Pollution Research* 20: 5076–5084.
- Otálora M. A. G., Aragón G., Martínez I. & Wedin M. (2013): Cardinal characters on a slippery slope – A re-evaluation of phylogeny, character evolution, and evolutionary rates in the jelly lichens (Collemaataceae s. str.). – *Molecular Phylogenetics and Evolution* 68: 185–198.
- Otálora M. A. G. & Wedin M. (2013): *Collema fasciculare* belongs in Arctomiaceae. – *Lichenologist* 45: 295–304.
- Otte V., Yakovchenko L., Clerc P. & Westberg M. (2013): *Candelariella commutata* sp. nov. for *C. unilocularis* auct. medioeur. – an arctic-alpine lichen on calcareous substrata from the Caucasus and Europe. – *Herzogia* 26: 217–222.
- Paoli L., Fiorini E., Munzi S., Sorbo S., Basile A. & Loppi S. (2013): Antimony toxicity in the lichen *Xanthoria parietina* (L.) Th. Fr.. – *Chemosphere* 93: 2269–2275.
- Paoli L., Munzi S., Fiorini E., Gaggi C. & Loppi S. (2013): Influence of angular exposure and proximity to vehicular traffic on the diversity of epiphytic lichens and the bioaccumulation of traffic-related elements. – *Environmental Science and Pollution Research* 20: 250–259.
- Park S.-Y. et al. (2013): *Agrobacterium tumefaciens*-mediated transformation of the lichen fungus, *Umbilicaria muehlenbergii*. – *PLoS ONE* 8(12): e83896.
- Parrot D., Jan S., Baert N., Guyot S. & Tomasi S. (2013): Comparative metabolite profiling and chemical study of *Ramalina siliquosa* complex using LC–ESI-MS/MS approach. – *Phytochemistry* 89: 114–124.
- Pino-Bodas R., Ahti T., Stenroos S., Martín M. P. & Burgaz A. R. (2013): Multilocus approach to species recognition in the *Cladonia humilis* complex (Cladoniaceae, Ascomycota). – *American Journal of Botany* 100: 664–678.
- Pino-Bodas R., Martín M. P., Burgaz A. R. & Lumbsch H. T. (2013): Species delimitation in *Cladonia* (Ascomycota): a challenge to the DNA barcoding philosophy. – *Molecular Ecology Resources* 13: 1058–1068.
- Prieto M. & Wedin M. (2013): Dating the diversification of the major lineages of Ascomycota (Fungi). – *PLoS ONE* 8(6): e65576.
- Prieto M., Baloch E., Tehler A. & Wedin M. (2013): Mazaedium evolution in the Ascomycota (Fungi) and the classification of mazaediate groups of formerly unclear relationship. – *Cladistics* 29: 296–308.
- Printzen C., Domaschke S., Fernández-Mendoza F. & Pérez-Ortega S. (2013): Biogeography and ecology of *Cetraria aculeata*, a widely distributed lichen with a bipolar distribution. – *MycKeys* 6: 33–53.
- Pykälä J. (2013): Additions to the lichen flora of Finland. VII. – *Graphis Scripta* 25: 21–29.
- Randlane T., Saag A., Thell A. & Ahti T. (2013): Third world list of cetrarioid lichens – in a new databased form, with amended phylogenetic and type information. – *Cryptogamie, Mycologie* 34: 79–84.

- Ravera S. & Brunialti G. (2013): Epiphytic lichens of a poorly explored National Park: Is the probabilistic sampling effective to assess the occurrence of species of conservation concern?. – *Plant Biosystems* 147: 115–124.
- Rico V. J. & Barrasa J. M. (2011): Flora Liquenológica Ibérica. Vol 9. Basidiomycota liquenizados y liquenícolas. Agaricales: Hygrophoraceae. Cantharellales: Clavulinaceae e Hydnaceae. – Sociedad Española de Liquenología, Madrid.
- Rikkinen J. (2013): Molecular studies on cyanobacterial diversity in lichen symbioses. – *MycKeys* 6: 3–32.
- Rivas Plata E. et al. (2013): A molecular phylogeny of Graphidaceae (Ascomycota, Lecanoromycetes, Ostropales) including 428 species. – *MycKeys* 6: 55–94.
- Rizzi G. & Giordani P. (2013): The ecology of the lichen genus *Xanthoparmelia* in Italy: An investigation throughout spatial scales. – *Plant Biosystems* 147: 33–39.
- Rolstad J., Ekman S., Andersen H. L. & Rolstad E. (2013): Genetic variation and reproductive mode in two epiphytic lichens of conservation concern: a transatlantic study of *Evernia divaricata* and *Usnea longissima*. – *Botany* 91: 69–81.
- Rubio-Salcedo M., Martínez I., Carreño F. & Escudero A. (2013): Poor effectiveness of the Natura 2000 network protecting Mediterranean lichen species. – *Journal for Nature Conservation* 21: 1–9.
- Runnel K., Rosenvald R. & Lõhmus A. (2013): The dying legacy of green-tree retention: Different habitat values for polypores and wood-inhabiting lichens. – *Biological Conservation* 159: 187–196.
- Saag A., Randlane T., Saag L., Thell A. & Ahti T. (2013). Third world list of cetrarioid lichens, a databased tool for documentation of nomenclatural data – lessons learned. – *Taxon* 62: 591–603.
- Sadowska-Deś A. D., Bálint M., Otte J. & Schmitt I. (2013): Assessing intraspecific diversity in a lichen-forming fungus and its green algal symbiont: Evaluation of eight molecular markers. – *Fungal Ecology* 6: 141–151.
- Sadowsky A. & Ott S. (2012): Photosynthetic symbionts in Antarctic terrestrial ecosystems: the physiological response of lichen photobionts to drought and cold. – *Symbiosis* 58: 81–90.
- Schei F. H., Blom H. H., Gjerde I., Grytnes J.-A., Heegaard E. & Sætersdal M. (2013): Conservation of epiphytes: Single large or several small host trees?. – *Biological Conservation* 168: 144–151.
- Schiefelbein U. (2013): Additions to the lichenized and lichenicolous fungi of Mecklenburg-Western Pomerania (Germany). – *Herzogia* 26: 65–72.
- Schiefelbein U. & Jansen F. (2013): Human impact on the lichen floras of two forest landscapes in Pomerania (Poland, Germany). – *Plant Diversity and Evolution* 130: 273–293.
- Seed L. et al. (2013): Modelling relationships between lichen bioindicators, air quality and climate on a national scale: Results from the UK OPAL air survey. – *Environmental Pollution* 182: 437–447.
- Selva S. B. (2013): The calicioid lichens and fungi of the Acadian Forest Ecoregion of northeastern North America, I. New species and range extensions. – *Bryologist* 116: 248–256.
- Shivarov V. V. (2013): New records of Verrucariaceae (Ascomycota) from Bulgaria. – *Mycobiota* 3: 11–17.
- Shrestha G. & St. Clair L. L. (2013): Lichens: a promising source of antibiotic and anticancer drugs. – *Phytochemistry Reviews* 12: 229–244.

- Shriver R. K., Cutler K. & Doak D. F. (2012): Comparative demography of an epiphytic lichen: support for general life history patterns and solutions to common problems in demographic parameter estimation. – *Oecologia* 170: 137–146.
- Singh G. et al. (2012): Genetic basis of self-incompatibility in the lichen-forming fungus *Lobaria pulmonaria* and skewed frequency distribution of mating-type idiomorphs: implications for conservation. – *PLoS ONE* 7(12): e51402.
- Singh G. et al. (2013): The sister-group relationships of the largest family of lichenized fungi, Parmeliaceae (Lecanorales, Ascomycota). – *Fungal Biology* 117: 715–721.
- Sipman H. et al. (2013): *Hyperphyscia adglutinata* ist zurück in Berlin – über die Erholung der Flechtenflora auf der Pfaueninsel. – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 145 [2012]: 139–150.
- Sohrabi M. et al. (2013): *Teuvoa*, a new lichen genus in Megasporaceae (Ascomycota: Pertusariales), including *Teuvoa junipericola* sp. nov. – *Lichenologist* 45: 347–360.
- Sohrabi M., Stenroos S., Myllys L., Søchting U., Ahti T. & Hyvönen J. (2013): Phylogeny and taxonomy of the ‘manna lichens’. – *Mycological Progress* 12: 231–269.
- Søndergaard J., Bach L. & Asmund G. (2013): Modelling atmospheric bulk deposition of Pb, Zn and Cd near a former Pb–Zn mine in West Greenland using transplanted *Flavocetraria nivalis* lichens. – *Chemosphere* 60: 2549–2556.
- Speranza M. et al. (2013): Nd-YAG laser irradiation damages to *Verrucaria nigrescens*. – *International Biodeterioration & Biodegradation* 84: 281–290.
- Spribile T. & Muggia L. (2013): Expanded taxon sampling disentangles evolutionary relationships and reveals a new family in Peltigerales (Lecanoromycetidae, Ascomycota). – *Fungal Diversity* 58: 171–184.
- Svensson M., Dahlberg A., Ranius T. & Thor G. (2013): Occurrence patterns of lichens on stumps in young managed forests. – *PLoS ONE* 8(4): e62825.
- Szczeptańska K., Kossowska M. & Wilk K. (2013): *Caloplaca subpallida* (Teloschistaceae), a lichen species new to Poland: distribution, ecology and taxonomic affinities. – *Acta Societatis Botanicorum Poloniae* 82: 85–89.
- Tehler A., Ertz D. & Irestedt M. (2013): The genus *Dirina* (Roccellaceae, Arthoniales) revisited. – *Lichenologist* 45: 427–476.
- Timsina B. A., Sorensen J. L., Weihrauch D. & Piercey-Normore M. D. (2013): Effect of aposymbiotic conditions on colony growth and secondary metabolite production in the lichen-forming fungus *Ramalina dilacerata*. – *Fungal Biology* 117: 731–743.
- Toni S. A. & Piercey-Normore M. D. (2013): Chemical ecology of lichens and species composition of cryptogams among three boreal habitats in eastern Manitoba. – *Botany* 91: 53–61.
- Tregidgo D. J., West S. E. & Ashmore M. R. (2013): Can citizen science produce good science? Testing the OPAL Air Survey methodology, using lichens as indicators of nitrogenous pollution. – *Environmental Pollution* 182: 448–451.
- Truong C., Divakar P. K., Yahr R., Crespo A. & Clerc P. (2013): Testing the use of ITS rDNA and protein-coding genes in the generic and species delimitation of the lichen genus *Usnea* (Parmeliaceae, Ascomycota). – *Molecular Phylogenetics and Evolution* 68: 357–372.
- Urbanavichus G. & Ismailov A. (2013): The lichen flora of Gunib plateau, inner-mountain Dagestan (North-East Caucasus, Russia). – *Turkish Journal of Botany* 37: 753–768.
- Urbanavichus G. & Urbanavichene I. (2013): New records of pyrenocarpous lichens from the NW Caucasus (Russia). – *Herzogia* 26: 123–129.