

## NOVÁ BRYOLOGICKÁ LITERATURA XX.

## New bryological literature, XX

Jan Kučera<sup>1</sup>, Svatava Kuběšová<sup>2</sup>, Michal Hájek<sup>3</sup> & Vítězslav Plásek<sup>4</sup>

<sup>1</sup> Jihočeská Univerzita, Přírodovědecká fakulta, Branišovská 31, CZ-370 05 České Budějovice, e-mail: kucera@prf.jcu.cz, <sup>2</sup> Botanické oddělení, Moravské zemské muzeum, Hvězdoslavova 29a, CZ-62700 Brno, skubesova@mzm.cz; <sup>3</sup> Masarykova univerzita, Ústav botaniky a zoologie, Kotlářská 2, CZ-61137 Brno, hajek@sci.muni.cz; <sup>4</sup> Ostravská univerzita, KBE PŘF, Chittussiho 10, CZ-71000 Ostrava, vitezslav.plasek@osu.cz

## Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Aboal J. R., Couto J. A., Fernández J. A. & Carballeira A. (2008): Physiological responses to atmospheric fluorine pollution in transplants of *Pseudoscleropodium purum*. – Environmental Pollution 153: 602–609.
- Afonina O. M. & Ignatova E. A. (2008 [2007]): A new species of *Didymodon* (Pottiaceae, Musci) from Asian Russia. – Arctoa 16: 133–138.
- Afonina O. M. & Ignatova E. A. (2008 [2007]): East Asian species of genus *Stereodon* (Brid.) Mitt. (Pylaisiaceae, Musci) in Russia. – Arctoa 16: 7–20.
- Ah-Peng C., Bardat J., Strasberg D. & Stieperaere H. (2008): Additions to the bryoflora of Réunion Island 2: *Anthocerotopsida*, *Marchantiopsida* and *Jungermanniopsida* from the herbarium of the National Botanic Garden of Belgium. – Journal of Bryology 30: 185–191.
- Aničić M., Tasić M., Frontasyeva M. V., Tomašević M., Rajšić S., Strelková L. P. & Steinnes E. (2007): Active moss biomonitoring of atmospheric trace element deposition in Belgrade urban area using ENAA and AAS. – In: AIP Conference Proceedings, Vol. 958, Nuclear Physics Methods And Accelerators In Biology And Medicine, p. 222–223, Prague.
- Arróniz-Crespo M., Phoenix G., Núñez-Olivera E. & Martínez-Abaigar J. (2008): Age-specific physiological responses to UV radiation in the aquatic liverwort *Jungermannia exsertifolia* subsp. *cordifolia*. – Cryptogamie Bryologie 29: 115–126.
- Asakawa Y., Toyota M., Hashimoto T., Tori M., Nagashima F. & Harinantenaina L. (2008): *Marchantiophyta* (liverworts): rich sources of macrocyclic bis(bibenzyls). – Heterocycles 76: 99–127.
- Austrheim G., Mysterud A., Pedersen B., Halvorsen R., Hassel K. & Evju M. (2008): Large scale experimental effects of three levels of sheep densities on an alpine ecosystem. – Oikos 117: 837–846.
- Baars C. & Edwards D. (2008): Effects of elevated atmospheric CO<sub>2</sub> on spore capsules of the moss *Leptobryum pyriforme*. – Journal of Bryology 30: 36–40.
- Bakalin V. A. & Vetrova V. P. (2008): Vegetation-permafrost relationships in the zone of sporadic permafrost distribution in the Kamchatka Peninsula. – Russian Journal of Ecology 39: 318–326.
- Bako S. P. & Afolabi S. (2008): Spatial distribution and heavy metal content of some bryophytes and lichens in relation to air pollution in Nigeria's Guinea Savanna. – International Journal of Environment and Pollution 33: 195–206.
- Barry M. J., Andreas B. K. & De Szalay F. A. (2008): Long-term plant community changes in managed fens in Ohio, USA. – Aquatic Conservation-Marine and Freshwater Ecosystems 18: 392–407.
- Bisang I., Ehrlén J. & Hedenäs L. (2008): Are annual growth intervals independent units in the moss *Pseudocalliergon trifarium* (Amblystegiaceae)? – Bryologist 111: 435–443.
- Bisang I. & Hedenäs L. (2008): Mate limitation does not explain the lack of capsules in a *Pseudocalliergon trifarium* population. – Journal of Bryology 30: 97–99.
- Breeuwer A., Heijmans M., Robroek B. J. M., Limpens J. & Berendse F. (2008): The effect of increased temperature and nitrogen deposition on decomposition in bogs. – Oikos 117: 1258–1268.
- Burch J. (2008): The relationship of bryophyte regeneration to heather canopy height following moorland burning on the North York Moors. – Journal of Bryology 30: 208–216.
- Buscardo E., Smith G. F., Kelly D. L., Freitas H., Iremonger S., Mitchell F. J. G., O'Donoghue S. & McKee A.-M. (2008): The early effects of afforestation on biodiversity of grasslands in Ireland. – Biodiversity and Conservation 17: 1057–1072.
- Callaghan D. A. & Ashton P. A. (2008): Attributes of rarity in a regional bryophyte assemblage. – Journal of Bryology 30: 101–107.

- Callaghan D. A. & Ashton P. A. (2008): Bryophyte distribution and environment across an oceanic temperate landscape. – *Journal of Bryology* 30: 23–35.
- Callaghan D. A. & Ashton P. A. (2008): Knowledge gaps in bryophyte distribution and prediction of species-richness. – *Journal of Bryology* 30: 147–158.
- Camilo V. J., Tobón C. & Breshears D. D. (2008): Fog interception by non-vascular epiphytes in tropical montane cloud forests: dependencies on gauge type and meteorological conditions. – *Hydrological Processes* 22: 2484–2492.
- Cao T., An L., Wang M., Lou Y., Yu Y., Wu J., Zhu Z., Qing Y. & Glime J. (2008): Spatial and temporal changes of heavy metal concentrations in mosses and its indication to the environments in the past 40 years in the city of Shanghai, China. – *Atmospheric Environment* 42: 5390–5402.
- Carballeira C. B., Aboal J. R., Fernández J. A. & Carballeira A. (2008): Comparison of the accumulation of elements in two terrestrial moss species. – *Atmospheric Environment* 42: 4904–4917.
- Cesa M., Campisi B., Bizzotto A., Ferraro C., Fumagalli F. & Nimis P. L. (2008): A factor influence study of trace element bioaccumulation in moss bags. – *Archives of Environmental Contamination and Toxicology* 55: 386–396.
- Chiarucci A., Calderisi M., Casini F. & Bonini I. (2008): Vegetation at the limits for vegetation: Vascular plants, bryophytes and lichens in a geothermal field. – *Folia Geobotanica* 43: 19–33.
- Chiyoda S., Ishizaki K., Kataoka H., Yamato K. T. & Kohchi T. (2008): Direct transformation of the liverwort *Marchantia polymorpha* L. by particle bombardment using immature thalli developing from spores. – *Plant Cell Reports* 27: 1467–1473.
- Clarke L. J. & Robinson S. A. (2008): Cell wall-bound ultraviolet-screening compounds explain the high ultraviolet tolerance of the Antarctic moss, *Ceratodon purpureus*. – *New Phytologist* 179: 776–783.
- Cleavitt N. L., Eschtruth A. K., Battles J. J. & Fahey T. J. (2008): Bryophyte response to eastern hemlock decline caused by hemlock woolly adelgid infestation. – *The Journal of the Torrey Botanical Society* 135: 12–25.
- Cole H. A., Newmaster S. G., Bell F. W., Pitt D. & Stinson A. (2008): Influence of microhabitat on bryophyte diversity in Ontario mixedwood boreal forest. – *Canadian Journal of Forest Research – Reue Canadienne de Recherche Forestière* 38: 1867–1876.
- Cooté L., Smith G. F., Kelly D. L., O'Donoghue S., Dowding P., Iremonger S. & Mitchell F. J. G. (2008): Epiphytes of Sitka spruce (*Picea sitchensis*) plantations in Ireland and the effects of open spaces. – *Biodiversity and Conservation* 17: 953–968.
- Cros R. M., Sáez L. & Brugués M. (2008): The bryophytes of the Balearic Islands: an annotated checklist. – *Journal of Bryology* 30: 74–95.
- Czernadyeva I. V. (2008 [2007]): The genus *Anacamptodon* (*Amblystegiaceae*, *Bryophyta*) in Russia and Transcaucasia. – *Arctoa* 16: 1–6.
- Da Costa D. P. & De Faria C. P. (2008): Conservation priorities for the bryophytes of Rio de Janeiro State, Brazil. – *Journal of Bryology* 30: 133–142.
- Dauphin G., Morales T. & Moreno E. J. (2008): Preliminar Catalogue of *Lejeuneaceae* (*Hepaticae*) of Venezuela (in Spanish). – *Cryptogamie Bryologie* 29: 215–265.
- Davis E. Christine & Shaw A. Jonathan (2008): Biogeographic and phylogenetic patterns in diversity of liverwort-associated endophytes. – *American Journal of Botany* 95: 914–924.
- Dengler J. & Boch S. (2007): Taxonomy, ecology, and distribution of six remarkable plant taxa (*Spermatophyta*, *Bryophyta*, and *Lichenes*) of the Estonian dry grassland flora. – *Phyton (Horn)* 47: 47–71.
- Dolnik C. & Napreenko M. G. (2008 [2007]): The bryophytes of the Southern Curonian spit (Baltic Sea coast). – *Arctoa* 16: 35–46.
- Dolný A., Plášek V., Drozd P. & Kočárek P. (2008): Affinity of dragonflies for *Sphagnum* ecological-groups. – In: 4<sup>th</sup> International Meeting on the Biology of *Sphagnum*, 2008, p. 7, University of Alaska, Juneau.
- Dragičević S., Veljić M. & Marin P. D. (2008): Liverworts from the Morača River Basin – taxonomic and ecological analysis. – *Archives of Biological Sciences* 60: 469–474.
- Drozd P., Dolný A., Jašík M., Kočárek P., Krupař M., Plášek V. & Ševčík J. (2008): Structure of invertebrate community associated with moss cushions. – In: 4<sup>th</sup> International Meeting on the Biology of *Sphagnum*, 2008, p. 7, University of Alaska, Juneau.
- Duckett J. G. & Ligrone R. (2008): Basidiomycetous endophytes in New Zealand *Aneuraceae* (simple thalloid liverworts, *Metzgeriidae*) and the derived status of the monotypic genus *Verdoornia*. – *Botany – Botanique* 86: 346–358.

- Epstein H. E., Walker, D. A., Raynolds M. K., Jia G. J. & Kelley A. M. (2008): Phytomass patterns across a temperature gradient of the North American arctic tundra. – *Journal of Geophysical Research-Biogeosciences* 113: G03S02.
- Erzberger P., Papp B. & Dragičević S. (2008): Notes on some newly recorded bryophytes from Montenegro. – *Journal of Bryology* 30: 167–170.
- Faubert J. (2007): Catalogue des bryophytes du Québec et du Labrador. – *Provancheria* 30: 1–138.
- Fenton N. J. & Bergeron Y. (2008): Does time or habitat make old-growth forests species rich? Bryophyte richness in boreal *Picea mariana* forests. – *Biological Conservation* 141: 1389–1399.
- Fernando E. S., Co L. C., Lagunzad D. A., Gruezo W. S., Barcelona J. F., Madulid D. A., Lapis A. B., Texon G. I., Manila A. C. & Zamora P. M. (2008): Threatened plants of the Philippines: A preliminary assessment. – *Asia Life Sciences, Suppl.* 3: 1–52.
- Forsum Å., Laudon H. & Nordin A. (2008): Nitrogen uptake by *Hylocomium splendens* during snowmelt in a boreal forest. – *Ecoscience* 15: 315–319.
- Frey W. & Stech M. (2008): New suprageneric taxa of liverworts (*Marchantiophyta*) and mosses (*Bryophyta*). – *Nova Hedwigia* 87: 261–267.
- Goncharova I. A. (2008): Features of moss mat structure and productivity in forest-bog complexes of West Siberia. – *Contemporary Problems of Ecology* 1: 467–475.
- Goncharova I. A. (2008): The hygroscopic moisture of mosses in bogs of Tomsk oblast. – *Russian Journal of Ecology* 39: 299–301.
- González-Mancebo J., Romaguera F., Ros R. M., Patino J. & Werner O. (2008): Bryophyte flora of the Canary Islands : an updated compilation of the species list with an analysis of distribution patterns in the context of the Macaronesian Region. – *Cryptogamie Bryologie* 29: 315–357.
- Gorr G. (2008): Recombinant production of therapeutic proteins using bryotechnology. – *Pharmazeutische Industrie* 70: 532–534.
- Grdović S. & Sabovljević M. (2008): An ecological approach to bryophytes of beehive yards: Is there a bee-moss relationship?. – *Acta Veterinaria, Beograd* 58: 287–294.
- Hallingbäck T., Lönnell N. & Weibull H. (2008): Nationalnyckeln till Sveriges flora och fauna. Bladmossor: Kompaktmossor – kapmossor. *Bryophyta: Anoetangium – Orthodontium*. – ArtDatabanken, SLU, Uppsala [504 pp.]
- Harris A. (2008): Spectral reflectance and photosynthetic properties of *Sphagnum* mosses exposed to progressive drought. – *Ecohydrology* 1: 35–42.
- Harris E. S. J. (2008): Paraphyly and multiple causes of phylogenetic incongruence in the moss genus *Plagiomnium* (*Mniaceae*). – *Taxon* 57: 417–433.
- Hawksworth D. L. (2008): Czech national park fungi, lichens and bryophytes [Review of: Antonín V., Gruna B., Hradílek Z., Vágner A. & Vězda A., 2000, Houby, lišejníky a mechorosty Národního parku Podyjí, Masarykova Univerzita, Brno]. – *Bryologist* 111: 521–522.
- Hedderson T. A. & Longton R. E. (2008): Local adaptation in moss life histories: population-level variation and a reciprocal transplant experiment. – *Journal of Bryology* 30: 1–11.
- Hedderson T. A. & Zander R. H. (2008): *Algaria nataliei* (*Pottiaceae*), a new moss genus and species from the Western Cape Province of South Africa. – *Journal of Bryology* 30: 192–195.
- Hedderson T. A. & Zander R. H. (2008): *Vrolijkheidia circumscissa* (*Pottiaceae*), a new moss genus and species from the Succulent Karoo of South Africa. – *Journal of Bryology* 30: 143–146.
- Hedenäs L. (2008): Molecular variation in *Drepanocladus aduncus* s.l. does not support recognition of more than one species in Europe. – *Journal of Bryology* 30: 108–120.
- Hernández-Maqueda R., Quandt D. & Muñoz J. (2008): Testing reticulation and adaptive convergence in the *Grimmiaceae* (*Bryophyta*). – *Taxon* 57: 500–510.
- Hock Zs., Szövényi P., Schneller J. J., Urmi E. & Tóth Z. (2008): Are sexual or asexual events determining the genetic structure of populations in the liverwort *Mannia fragrans*?. – *Journal of Bryology* 30: 66–73.
- Hooijmaijers C. (2008): Membrane integrity, oxidative damage and chlorophyll fluorescence during dehydration of the thalloid liverwort *Monoclea forsteri* Hook. – *Journal of Bryology* 30: 217–222.
- Hrivňák R., Hájek M., Blanár D., Kochjarová J. & Hájková P. (2008): Mire vegetation of the Muranska Planina Mts - formalised classification, ecology, main environmental gradient and influence of geographical position. – *Biologia* 63: 368–377.
- Hugonnot V. (2008): Chorologie et écologie d'*Orthotrichum rogeri* Brid. en France. – *Cryptogamie Bryologie* 29: 275–297.
- Hugonnot V. (2008): *Antitrichia californica* Sull. (*Leucodontaceae*) in France. Identification, distribution, habitat and communities. – *Cryptogamie Bryologie* 29: 359–385.

- Hutsemekers V., Risterucci A. M., Ricca M., Boles S., Hardy O. J., Shaw A. J. & Vanderpoorten A. (2008): Identification and characterization of nuclear microsatellite loci in the aquatic moss *Platyhypnidium riparioides* (*Brachytheciaceae*). – *Molecular Ecology Resources* 8: 1130–1132.
- Huttunen S., Hedenäs L., Ignatov M. S., Devos N. & Vanderpoorten A. (2008): Origin and evolution of the northern hemisphere disjunction in the moss genus *Homalothecium* (*Brachytheciaceae*). – *American Journal of Botany* 95: 720–730.
- Ignatov M. S. & Milyutina I. A. (2008 [2007]): A revision of the genus *Sciuro-hypnum* (*Brachytheciaceae*, *Bryophyta*) in Russia. – *Arctoa* 16: 63–86.
- Ignatov M. S. & Milyutina I. A. (2008 [2007]): On *Sciuro-hypnum oedipodium* and *S. curtum* (*Brachytheciaceae*, *Bryophyta*). – *Arctoa* 16: 47–61.
- Ignatova E. A. & Ignatov M. S. (2008 [2007]): *Didymodon glaucus* Ryan (*Pottiaceae*, *Musci*) – the first record from Siberia. – *Arctoa* 16: 133–138.
- Jamilena M., Mariotti B. & Manzano S. (2008): Plant sex chromosomes: molecular structure and function. – *Cytogenetic and Genome Research* 120: 255–264.
- Jasmin J.-N., Rochefort L. & Gauthier G. (2008): Goose grazing influences the fine-scale structure of a bryophyte community in arctic wetlands. – *Polar Biology* 31: 1043–1049.
- Jeschke M. & Kiehl K. (2008): Effects of a dense moss layer on germination and establishment of vascular plants in newly created calcareous grasslands. – *Flora* 203: 557–566.
- Jonášova M. & Prach K. (2008): The influence of bark beetles outbreak vs. salvage logging on ground layer vegetation in Central European mountain spruce forests. – *Biological Conservation* 141: 1525–1535.
- Kalniņa L. (2007): Diversity of mire origin and history in Latvia. – *Peatlands International 2007*: 54–56.
- Karlin E. F., Boles S. B. & Shaw J. A. (2008): Resolving boundaries between species in *Sphagnum* section *Subsecunda* using microsatellite markers. – *Taxon* 57: 1189–1200, E1–E2.
- Karlin E. F., Boles S. B. & Shaw J. A. (2008): Systematics of *Sphagnum* section *Sphagnum* in New Zealand: a microsatellite-based analysis. – *New Zealand Journal of Botany* 46: 105–118.
- Kleppin L., Pesch R. & Schröder W. (2008): CHAID Models on boundary conditions of metal accumulation in mosses collected in Germany in 1990, 1995 and 2000. – *Atmospheric Environment* 42: 5220–5231.
- Klos A., Rajfur M., Waclawek M. & Waclawek W. (2008): Accumulation of micro- and macroelements in mosses and lichens. – *Ecological Chemistry and Engineering S/Chemia i Inżynieria Ekologiczna S* 15: 397–423.
- Klos A., Rajfur M., Waclawek M., Waclawek W., Frontasyeva M. V. & Pankratova J. S. (2008): The influence of unidentified pollution sources on the irregularity of biomonitoring tests results. – *Water, Air and Soil Pollution* 191: 345–352.
- Knapp S., Kühn I., Mosbrugger V. & Klotz S. (2008): Do protected areas in urban and rural landscapes differ in species diversity?. – *Biodiversity and Conservation* 17: 1595–1612.
- Kočárek P., Grucmanová Š., Filipcová Z., Bradová L., Plášek V. & Holuša, J. (2008): Mosses as a food component of groundhopper *Tetrix ceperoi* (*Orthoptera: Tetrigidae*). – In: 4th International Meeting on the Biology of *Sphagnum*, 2008, p. 9, University of Alaska, Juneau.
- Krumnikl M., Sojka E., Gaura J. & Motyka O. (2008): A new method for bryophyte canopy analysis based on 3D surface reconstruction. – In: Seventh International Conference On Computer Information Systems and Industrial Management Applications, Proceedings, p. 210–211, Ostrava.
- Kučerová A., Rektoris L., Štechová T. & Bastl M. (2008): Disturbances on a wooded raised bog – How windthrow, bark beetle and fire affect vegetation and soil water quality? – *Folia Geobotanica* 43: 49–67.
- Lappalainen N. M., Huttunen S. & Suokanerva H. (2007): Red-stemmed feather moss *Pleurozium schreberi* (Britt.) Mitt. – A bioindicator of UV radiation? – *Progress In Environmental Science and Technology*, Vol 1: 14–18.
- Lara F., Garillete R. & Mazimpaka V. (2007): Nyckel til släktet *Orthotrichum* i Norden. – *Myrinia* 17: 105–112.
- Leith I. D., Mitchell R. J., Truscott A. -M., Cape J. N., van Dijk N., Smith R. I., Fowler D. & Sutton M. A. (2008): The influence of nitrogen in stemflow and precipitation on epiphytic bryophytes, *Isoetecium myosuroides* Brid., *Dicranum scoparium* Hewd. and *Thuidium tamariscinum* (Hedw.) Schimp of Atlantic oakwoods. – *Environmental Pollution* 155: 237–246.
- Limpens J. & Heijmans M. M. P. D. (2008): Swift recovery of *Sphagnum* nutrient concentrations after excess supply. – *Oecologia* 157: 153–161.
- Liu X.-Y., Xiao H.-Y., Liu C.-Q., Li Y.-Y. & Xiao H.-W. (2008): Stable carbon and nitrogen isotopes of the moss *Haplcladium microphyllum* in an urban and a background area (SW China): The role of

- environmental conditions and atmospheric nitrogen deposition. – *Atmospheric Environment* 42: 5413–5423.
- Liu X.-Y., Xiao H.-Y., Liu C.-Q., Li Y.-Y. & Xiao H.-W. (2008): Tissue N content and N-15 natural abundance in epilithic mosses for indicating atmospheric N deposition in the Guiyang area, SW China. – *Applied Geochemistry* 23: 2708–2715.
- Liu Y. & Jia Y. (2008): *Lejeuneaceae* (*Marchantiophyta*) share an indel in chloroplast *rps4* gene. – *Journal of Bryology* 30: 174–177.
- Liu Y., Lia Y., Wang W., Chen Z.-D., Davis E. C. & Qiu Y.-L. (2008): Phylogenetic relationships of two endemic genera from East Asia: *Trichocoleopsis* and *Neotrichocolea* (*Hepaticae*). – *Annals of the Missouri Botanical Garden* 95: 459–470.
- Lõhmus A. & Lõhmus P. (2008): First-generation forests are not necessarily worse than long-term managed forests for lichens and bryophytes. – *Restoration Ecology* 16: 231–239.
- Maksimov A. I. (2008 [2007]): *Sphagnum imbricatum* complex (*Sphagnaceae*, *Bryophyta*) in Russia. – *Arctoa* 16: 21–24.
- Martínez-Abaigar J., Otero S., Tomás R. & Núñez-Olivera E. (2008): High-level phosphate addition does not modify UV effects in two aquatic bryophytes. – *Bryologist* 111: 444–454.
- McClymont E. L., Mauquoy D., Yeloff D., Broekens P., van Geel B., Charman D. J., Pancost R. D., Chambers F. M. & Evershed R. P. (2008): The disappearance of *Sphagnum imbricatum* from Butterburn Flow, UK. – *Holocene* 18: 991–1002.
- Meyer M., Seibt U. & Griffiths H. (2008): To concentrate or ventilate? Carbon acquisition, isotope discrimination and physiological ecology of early land plant life. – *Philosophical Transactions of the Royal Society B, Biological Sciences*, 363: 2767–2778.
- Mietelski J. W., Olech M. A., Sobiech-Matura K., Howard B. J., Gaca P., Zwolak M., Błażej S. & Tomankiewicz E. (2008): <sup>137</sup>Cs, <sup>40</sup>K, <sup>238</sup>Pu, <sup>239+240</sup>Pu and <sup>90</sup>Sr in biological samples from King George Island (Southern Shetlands) in Antarctica. – *Polar Biology* 31: 1081–1089.
- Mosseichik Y. V., Ignatov M. S. & Ignatiev I. A. (2008 [2007]): A bryophyte-like plant from the Lower Carboniferous of the Moscow Coal Basin. – *Arctoa* 16: 99–122.
- Mykrä H., Aroviita J., Hämäläinen H., Kotanen J., Vuori K.-M. & Muotka T. (2008): Assessing stream condition using macro invertebrates and macrophytes: concordance of community responses to human impact. – *Fundamental and Applied Limnology* 172: 191–203.
- Németh C. (2007): Data on the distribution of some rare alpine-boreal bryophytes in the Vértes Mts (Hungary). – *Studia Botanica Hungarica* 38: 59–70.
- Osipov S. V., Cherdantseva V. Ya., Galanina I. A. & Yakubov V. V. (2008): Species composition and ecologo-phytocenotic spectra of vascular plants, mosses, and lichens on gold-mining sites in the taiga zone of the lower Amur River basin, the Russian Far East. – *Contemporary Problems of Ecology* 1: 425–439.
- Otnyukova T. N. (2008 [2007]): Notes on *Dicranum* (*Dicranaceae*, *Musci*) in Russia. 2. *Dicranum pseudacutifolium*, sp. nov. from North Siberia. – *Arctoa* 16: 163–168.
- Pajunen A., Virtanen R. & Roininen H. (2008): The effects of reindeer grazing on the composition and species richness of vegetation in forest-tundra ecotone. – *Polar Biology* 31: 1233–1244.
- Paltto H., Nordén B. & Götmark F. (2008): Partial cutting as a conservation alternative for oak (*Quercus* spp.) forest – Response of bryophytes and lichens on dead wood. – *Forest Ecology and Management* 256: 536–547.
- Paton A. J., Brummitt N., Govaerts R., Harman K., Hinchcliffe S., Allkin B. & Lughadha E. N. (2008): Towards Target 1 of the Global Strategy for Plant Conservation: a working list of all known plant species - progress and prospects. – *Taxon* 57: 602–611.
- Pesch R., Schröder W., Dieffenbach-Fries H., Genssler L. & Kleppin L. (2008): Improving the design of environmental monitoring networks. Case study on the heavy metals in mosses survey in Germany. – *Ecological Informatics* 3: 111–121.
- Pesch R., Schröder W., Schmidt G. & Genssler L. (2008): Monitoring nitrogen accumulation in mosses in central European forests. – *Environmental Pollution* 155: 528–536.
- Picardi E. & Quagliariello C. (2008): Is plant mitochondrial RNA editing a source of phylogenetic incongruence? An answer from *in silico* and *in vivo* data sets. – *BMC Bioinformatics* 9, Suppl. 2: S14.
- Piessens K., Stieperaere H., Honnay O. & Hermy M. (2008): Effects of management and adjacent forest on the heathland bryophyte layer. – *Basic and Applied Ecology* 9: 253–262.
- Plášek V. & Marková I. (2008): *Orthotrichum pulchellum*. – In: Blockeel T. L. & al., New national and regional bryophyte records, 18, p. 163–164, *Journal of Bryology* 30: 161–166.

- Pohjamo M., Korpelainen H. & Kalinauskaitė N. (2008): Restricted gene flow in the clonal hepatic *Trichocolea tomentella* in fragmented landscapes. – *Biological Conservation* 141: 1204–1217.
- Popper Z. A. (2008): Evolution and diversity of green plant cell walls. – *Current Opinion in Plant Biology* 11: 286–292.
- Porter P. R., Evans A. J., Hodson A. J., Lowe A. T. & Crabtree M. D. (2008): Sediment-moss interactions on a temperate glacier: Falljökull, Iceland. – *Annals of Glaciology* 48: 25–31.
- Pressel S., Ligrone R. & Duckett J. G. (2008): Cellular differentiation in moss protonemata: A morphological and experimental study. – *Annals Of Botany* 102: 227–245.
- Rawat K. K. & Srivastava S. C. (2007): Genus *Plagiochila* in Eastern Himalaya (India). – Bishen Singh Mahendra Pal Singh, Dehra Dun, India. [xiv + 259 pp.]
- Real C., Fernández J. A., Aboal J. R. & Carballeira A. (2008): Detection of pulses of atmospheric mercury deposition with extensive surveys and frequently sampled stations: A comparison. – *Ecotoxicology and Environmental Safety* 70: 392–399.
- Rüdinger M., Polsakiewicz M. & Knoop V. (2008): Organellar RNA editing and plant-specific extensions of pentatricopeptide repeat proteins in jungermanniid but not in marchantiid liverworts. – *Molecular Biology and Evolution* 25: 1405–1414.
- Sabovljević M., Sabovljević Aneta, Radulović J. & Dragičević Ivana (2008): Genetic variability within Serbian populations of the rare and endangered pottioid moss *Hilpertia velenovskyi* (Schiffn.) Zander inferred by isozyme analyses. – *Archives of Biological Sciences* 60: 207–213.
- Sahoo S. K., Enomoto H., Tokonami S., Ishikawa Te., Ujić P., Čeliković I. & Zunić Z. S. (2008): Determination of depleted uranium in environmental bio-monitor samples and soil from target sites in western Balkan region. – *Natural Radiation Environment* 1034: 287–290.
- Sardans J. & Peñuelas J. (2008): Drought changes nutrient sources, content and stoichiometry in the bryophyte *Hypnum cupressiforme* Hedw. growing in a Mediterranean forest. – *Journal of Bryology* 30: 59–65.
- Sedeľnikov V. P., Naurnenko Yu. V., Sedeľnikova N. V., Gorbunova I. A., Pisarenko O. Yu. & Shauro D. N. (2008): Biodiversity and spatial arrangement of Siberian flora. – *Contemporary Problems of Ecology* 1: 14–21.
- Sérgio C., Sim-Sim M., Fontinha S. & Figueira R. (2008): List of bryophytes (*Bryophyta*). – In: Borges P. A. V., Abreu C., Aguiar A. M. F., Carvalho P., Jardim R., Melo I., Oliveira P., Sérgio C., Serrano A. R. M. & Vieira. P. (eds.), A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos, p. 143–156, Direcção Regional do Ambiente da Madeira and Universidade dos Açores, Funchal and Angra do Heroísmo.
- Sérgio C., Sim-Sim M., Fontinha S. & Figueira R. (2008): The bryophytes (*Bryophyta*) of the Madeira and Selvagens archipelagos. – In: Borges P. A. V., Abreu C., Aguiar A. M. F., Carvalho P., Jardim R., Melo I., Oliveira P., Sérgio C., Serrano A. R. M. & Vieira. P. (eds.), A list of the terrestrial fungi, flora and fauna of Madeira and Selvagens archipelagos, p. 123–141, Direcção Regional do Ambiente da Madeira and Universidade dos Açores, Funchal and Angra do Heroísmo.
- Shaw A. J., Pokorny L., Shaw B., Ricca M., Boles S. & Szövényi P. (2008): Genetic structure and genealogy in the *Sphagnum subsecundum* complex (*Sphagnaceae: Bryophyta*). – *Molecular Phylogenetics and Evolution* 49: 304–317.
- Shaw A. J., Boles S. & Shaw B. (2008): A phylogenetic delimitation of the „*Sphagnum subsecundum* complex“ (*Sphagnaceae, Bryophyta*). – *American Journal of Botany* 95: 731–744."
- Shetler G., Turetsky M. R., Kane E. & Kasischke E. (2008): *Sphagnum* mosses limit total carbon consumption during fire in Alaskan black spruce forests. – *Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière* 38: 2328–2336.
- Sim-Sim M., Luís L., Garcia C., Fontinha S., Lobo C., Martins S. & Stech M. (2008): New data on the status of threatened bryophytes of Madeira Island. – *Journal of Bryology* 30: 226–228.
- Skrzypczak R. (2007): *Barbula amplexifolia* (Mitt.) A. Jaeger présent en France (Savoie). – *Bulletin de la Société Botanique du Centre-Ouest* n.s. 38: 321–326.
- Smith G. F., Gittings T., Wilson M., French L., Oxbrough A., O'Donoghue S., O'Halloran J., Kelly D. L., Mitchell F. J. G., Kelly T., Iremonger S., McKee A.-M. & Giller P. (2008): Identifying practical indicators of biodiversity for stand-level management of plantation forests. – *Biodiversity and Conservation* 17: 991–1015.
- Sotiaux A., Sotiaux O. & Vanderpoorten A. (2008): Additions to the bryophyte flora of Corsica. – *Cryptogamie Bryologie* 29: 267–274.
- Steinnes E. (2008): Use of mosses to study atmospheric deposition of trace elements: contributions from investigations in Norway. – *International Journal of Environment and Pollution* 32: 499–508.

- Stevenson C. R. & Hill M. O. (2008): Urban myths exploded: results of a bryological survey of King's Lynn (Norfolk, UK). – *Journal of Bryology* 30: 12–22.
- Suyama Y., Gunnarsson U. & Parducci L. (2008): Analysis of short DNA fragments from Holocene peatmoss samples. – *Holocene* 18: 1003–1006.
- Teleganova V. V. & Ignatov M. S. (2008 [2007]): On the reproductive biology of *Ptilium crista-castrensis* (*Pylaisiaceae*, *Bryophyta*). – *Arctoa* 16: 87–98.
- Thiebaut G., Giamberini L. & Ghanbaja J. (2008): Cellular distribution of metals in a liverwort and a moss transplanted to two streams of differing acidity. – *Water, Air and Soil Pollution* 192: 263–272.
- Tsegmed Ts. & Ignatova E. A. (2008 [2007]): Distribution and ecology of *Grimmia* (*Grimmiaceae*, *Bryophyta*) in Mongolia. – *Arctoa* 16: 157–162.
- Vergel K. N., Frontasyeva M. V., Pavlov S. S. & Povtoreyko E. A. (2007): Air pollution studies in Tver region of Russia using moss-biomonitoring with nuclear analytical methods. – *Nuclear Physics Methods and Accelerators In Biology and Medicine* 958: 240–241.
- Verma P. K. & Srivastava S. C. (2008): Diversity and distribution of epiphytic liverworts (*Hepaticae*) in Nilgiri hills (Tamil Nadu). – *Proceedings of the National Academy of Sciences India, Section B, Biol. Sciences* 78: 253–264.
- Vilnet A. A., Konstantinova N. A. & Troitsky A. V. (2008): Phylogeny and systematics of the genus *Lophozia* s. str. (Dumort.) Dumort. (*Hepaticae*) and related taxa from nuclear ITS1–2 and chloroplast *trnL*–F sequences. – *Molecular Phylogenetics and Evolution* 47: 403–418.
- Wahrmund U., Groth-Malonek M. & Knoop V. (2008): Tracing plant mitochondrial DNA evolution: Rearrangements of the ancient mitochondrial gene cluster *trnA-trnT-nad7* in liverwort phylogeny. – *Journal of Molecular Evolution* 66: 621–629.
- Wallmann K. (2008): Mesozoic climate – Liverworts and all. – *Nature Geoscience* 1: 14–15.
- Wickett N. J., Fan Y., Lewis P. O. & Goffinet B. (2008): Distribution and evolution of pseudogenes, gene losses, and a gene rearrangement in the plastid genome of the nonphotosynthetic liverwort, *Aneura mirabilis* (*Metzgeriales*, *Jungermannniopsida*). – *Journal of Molecular Evolution* 67: 111–122.
- Wilbraham J. (2008): Bryophyte Flora of Uganda. 8. *Orthotrichaceae* Part 1 – *Macromitrioideae*. – *Journal of Bryology* 30: 201–207.
- Wójciak H. (2007): Porosty, Mszaki, Paprotniki. Wydanie II. – MULTICO, Warsaw, Poland. [368 pp.]
- Wyatt H. D. M., Ashton N. W. & Dahms T. E. S. (2008): Cell wall architecture of *Physcomitrella patens* is revealed by atomic force microscopy. – *Botany – Botanique* 86: 385–397.
- Xiao H.-Y., Tang C.-G., Liu X.-Y., Xiao H.-W. & Liu Co.-Q. (2008): Sulphur isotopic ratios in mosses indicating atmospheric sulphur sources in Southern Chinese mountainous areas. – *Geophysical Research Letters* 35: –L19807.
- Xie C. & Lou H. (2008): Chemical constituents from the Chinese bryophytes and their reversal of fungal resistance. – *Current Organic Chemistry* 12: 619–628.
- Xu S., Yin C., He M. & Wang Y. (2008): A Technology for Rapid Reconstruction of Moss-Dominated Soil Crusts. – *Environmental Engineering Science* 25: 1129–1137.
- Yin-Long Q. (2008): Phylogeny and evolution of charophytic algae and land plants. – *Journal of Systematics and Evolution* 46: 287–306.
- Yu J., Ma P.-J., Shi D.-J., Li S.-M. & Wang C.-L. (2008): Homologous comparisons of photosynthetic system I genes among cyanobacteria and chloroplasts. – *Journal of Integrative Plant Biology* 50: 929–940.
- Zander R.-H. (2008): Evolutionary inferences from non-monophyly on molecular trees. – *Taxon* 57: 1182–1188.

## **Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]**

### 1. Články a knižní publikace [Papers and books]

- Bączkiewicz A., Klama H. & Buczkowska K. (2008): Genetic structure of leafy liverwort *Lepidozia reptans* detected by isozyme electrophoresis. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 112–118, Ostravská Univerzita, Ostrava.
- Fojcik B. (2008): Impact of anthropogenic habitats occurrence on the local pattern of moss species distribution. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 119–125, Ostravská Univerzita, Ostrava.

- Klama H., Salachna A., Smieja A., Buczkowska K. & Bączkiewicz A. (2008): The species of genus *Chiloscyphus* in Tatra National Park (S Poland). – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 126–133, Ostravská Univerzita, Ostrava.
- Kliment J. (ed.) (2008): Příroda Velké Fatry. Lišajníky, machorasty, cievnaté rastliny. – Univerzita Komenského, Bratislava [408 pp.]
- Kočárek P., Grucmanová Š., Filipcová Z., Bradová L., Plášek V. & Holuša J. (2008): Bryophagy in the groundhopper *Tetrix ceperoi* (Orthoptera: Tetrigidae): analysis of alimentary tract contents. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 348–352, Ostravská Univerzita, Ostrava.
- Kubešová S. & Novotný I. (2008): The mosses *Orthodontium lineare* and *Dicranum tauricum* in nature reserves in the Žďárské vrchy Protected Landscape Area (Czech Republic). – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 134–137, Ostravská Univerzita, Ostrava.
- Kučera J. (ed.) (2008): Zajímavé bryofloristické nálezy XI. [Interesting bryofloristic records, XI]. – Bryonora 41: 25–27.
- Kučera J., Kubešová S. & Plášek V. (2008): Nová bryologická literatura XIX. [New bryological literature, XIX]. – Bryonora 41: 34–43.
- Lackovičová A., Pišút P. & Guttová A. (2008): Epiphytic lichens – bioindicators of air pollution in Bratislava (SW Slovakia). – In: Kočárek P., Plášek V., Malachová K. & Cimalová O. (eds) Environmental changes and biological assessment IV (Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186), p. 138–142, Univ. Ostrava, Ostrava.
- Machowska A. (2008): Epixylous organisms on the decaying spruce wood (*Picea abies*) in the upper forest belt on the Polish part of Karkonosze Mts. (Poland). The first results. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 143–149, Ostravská Univerzita, Ostrava.
- Motyka O., Jašík M. & Vondřejc T. E. (2008): Analysis of possible factors affecting moss canopy density. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 150–153, Ostravská Univerzita, Ostrava.
- Novotný I. (2008): The *Ceratodon* moss genus in the Czech Republic and Slovakia. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 154–162, Ostravská Univerzita, Ostrava.
- Petrášová A. & Šuvada R. (2008): Bryofloristicko-ekologická štúdia Národného parku Slovenský kras. – Bulletin Slovenskej Botanickéj Spoločnosti 30: 169–177.
- Plášek V. & Cimalová Š. (eds.) (2008): Zajímavé botanické nálezy z regionu severní Moravy a Slezska I. – Časopis Slezského Zemského Muzea, Ser. A, 57: 45–48.
- Plášek V., Zmrhalová M. & Číhal L. (2008): Epiphytic mosses within *Orthotrichaceae* family in the Hrubý Jeseník Mts. A preliminary comparison of historical & recent data. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 163–169, Ostravská Univerzita, Ostrava.
- Sawicki J. & Zieliński R. (2008): Phylogenetic relationships between five *Sphagnum* species of the section *Acutifolia* based on DNA markers. – Časopis Slezského Zemského Muzea, Ser. A, 57: 63–80.
- Sawicki J., Plášek V. & Szczecińska M. (2008): Testing candidate *trnH-psbA* noncoding plant barcode region in the *Orthotrichum* genus. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 170–177, Ostravská Univerzita, Ostrava.
- Saxena D.K., Srivastava K. & Singh S. (2008): Retrospective metal data of past 10 years by analysis of moss *Barbula* sp. from Mussoorie city of India. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 178–182, Ostravská Univerzita, Ostrava.
- Stebel A., Smieja A., Stachurska-Swakoń A. & Żarnowiec J. (2008): *Bryum weigelii* (Bryophyta, Bryaceae) in the Polish part of the Carpathians. – In: Kočárek P., Plášek V., Malachová K. & Cimalová Š. (eds.), Environmental changes and biological assessment IV, Scripta Facultatis Rerum Naturalium Universitatis Ostraviensis 186, p. 183–188, Ostravská Univerzita, Ostrava.