

## NOVÁ BRYOLOGICKÁ LITERATURA XVII.

### New bryological literature, XVII

Jan Kučera

Jihočeská Univerzita, biologická fakulta, Branišovská 31, CZ–370 05 České Budějovice, e-mail:  
kucera@bf.jcu.cz

### Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Adamo P., Crisafulli P., Giordano S., Minganti V., Modenesi P., Monaci F., Pittao E., Tretiach M. & Bargagli R. (2007): Lichen and moss bags as monitoring devices in urban areas. Part II: Trace element content in living and dead biomonitoring and comparison with synthetic materials. – Environmental Pollution 146: 392–399.
- Adams B. J., Bardgett R. D., Ayres E., Wall D. H., Aislabie J., Bamforth S., Bargagli R., Cary C., Cavacini P., Connell L., Convey P., Fell J. W., Frati F., Hogg I. D., Newsham K. K., O'Donnell A., Russell N., Seppelt R. D. & Stevens M. I. (2006): Diversity and distribution of Victoria Land biota. – Soil Biology & Biochemistry 38: 3003–3018.
- Anderson H. M., Gale M. R., Jurgensen M. F. & Trettin C. C. (2007): Vascular and non-vascular plant community response to silvicultural practices and resultant microtopography creation in a forested wetland. – Wetlands 27: 68–79.
- Andrus R. E. (2006): Six new species of *Sphagnum* (*Bryophyta: Sphagnaceae*) from North America. – Sida 22: 959–972.
- Andrus R. E. (2007): *Sphagnum atlanticum* a new species from eastern North America. – Bryologist 110: 274–278.
- Andrus R. E. (2007): *Sphagnum oregonense* sp. nov. from western North America. – Bryologist 110: 123–125.
- Aoki S. (2006): The moss *Physcomitrella patens* as a potential model plant for studying circadian rhythms. – Bryological Research. 9: 50–59.
- Aryanti N. S. & Gradstein S. R. (2007): Wallace's line and the distribution of the liverworts of Sulawesi. – Cryptogamie Bryologie 28: 3–14.
- Bakalin V. A. 2006. Ekologicheskie osobennosti i zakonomernosti formirovaniya flory pechenochnikov pod vliyaniem aktivnogo vulkanizma i sovremennoy lednikovoy deyatel'nosti (na primere vulkana Ushkovskiy, Kamchatka) [Ecological features and trends in formation of the flora of liverworts under the impact of current volcanic and glacial activities: an example of Ushkovskii Volcano, Kamchatka]. – Ekologia – Russian Journal of Ecology 37: 16–22.
- Bakalin V. A. & Cherdantseva V. Ya. (2007 [2006]): Bryophytes of northern Kuril Islands (North-West Pacific). – Arctoa 15: 131–153.
- Banu-Fattah K. & Hadiuzzaman S. (2006): Acrocarpous mosses of Bangladesh. XI family: Bartramiaceae-2. – Bangladesh Journal of Botany 35: 91–97.
- Belinchón R., Martínez I., Escudero A., Aragón G. & Valladares F. (2007): Edge effects on epiphytic communities in a Mediterranean *Quercus pyrenaica* forest. – Journal of Vegetation Science 18: 81–90.
- Belyea L. R. & Baird A. J. (2006): Beyond "The limits to peat bog growth": cross-scale feedback in peatland development. – Ecological Monographs 76: 299–322.
- Benscoter B. W. (2006): Post-fire bryophyte establishment in a continental bog. – Journal of Vegetation Science 17: 647–652.
- Benscoter B. W. & Vitt D. H. (2007): Evaluating feathermoss growth: a challenge to traditional methods and implications for the boreal carbon budget. – Journal of Ecology 95: 151–158.
- Biggs L. & Wittkuhn R. S. (2006): Bryophyte diaspore banks in soil and litter of an urban bushland in Perth, Western Australia. – Hikobia 14: 375–382.
- Björk R. G. & Molau U. (2007): Ecology of alpine snowbeds and the impact of global change. – Arctic, Antarctic, and Alpine Research 39: 34–43.
- Björn L. O. (2007): Stratospheric ozone, ultraviolet radiation, and cryptogams. – Biological Conservation 135: 326–333.

- Blom H. H., Ignatova E. A. & Afonina O. M. (2007 [2006]): New records of *Schistidium* (*Grimmiaceae, Musci*) in Russia. – *Arctoa* 15: 187–194.
- Braun U. (2006): Type material in the herbarium of the Martin-Luther-University Halle-Wittenberg (HAL). *Bryophyta*: Species described by G. E. L. Hampe, C. F. Hornschuch, J. K. (C.) A. F. W. Müller (Müll. Hal.), etc. – *Schlechtendalia* 14: 25–31.
- Budke J. M. & Goffinet B. (2006): Phylogenetic analyses of *Timmiaceae* (*Bryophyta: Musci*) based on nuclear and chloroplast sequence data. – *Systematic Botany* 31: 633–641.
- Budke J. M., Jones C. S. & Goffinet B. (2007): Development of the enigmatic peristome of *Timmia megapolitana* (*Timmiaceae; Bryophyta*). – *American Journal of Botany* 94: 460–467.
- Calabrese G. M. (2006): A taxonomic revision of *Zygodon* (*Orthotrichaceae*) in southern South America. – *Bryologist* 109: 453–509.
- Chiarucci A., D'Auria F. & Bonini I. (2007): Is vascular plant species diversity a predictor of bryophyte species diversity in Mediterranean forests?. – *Biodiversity and Conservation* 16: 525–545.
- Chirino C., Campeau S. & Rochefort L. (2006): *Sphagnum* establishment on bare peat: The importance of climatic variability and *Sphagnum* species richness. – *Applied Vegetation Science* 9: 285–294.
- Cleavitt N. L., Klima K. L. & Fahey T. J. (2007): The role of water availability in seasonal growth of *Bazzania trilobata* on boulders in an eastern hemlock stand. – *Bryologist* 110: 295–308.
- Craw D., Rufaut C., Haffert L. & Paterson L. (2007): Plant colonization and arsenic uptake on high arsenic mine wastes, New Zealand. – *Water, Air, & Soil Pollution* 179: 351–364.
- Crosby M. R. & Engel J. J. (2006): Index of Hepatics, 1974–2000. – The Hattori Botanical Laboratory, Nichinan. [368 pp.]
- Davey M. L. & Currah R. S. (2006): Interactions between mosses (*Bryophyta*) and fungi. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 84: 1509–1519.
- Davies L., Bates J. W., Bell J. N. B., James P. W. & Purvis O. W. (2007): Diversity and sensitivity of epiphytes to oxides of nitrogen in London. – *Environmental Pollution* 146: 299–310.
- Davies T. D. (2007): Sulphate toxicity to the aquatic moss, *Fontinalis antipyretica*. – *Chemosphere* 66: 444–451.
- Diviš P., Dočekalová H., Brulík L., Pavliš M. & Hekera P. (2007): Use of the diffusive gradients in thin films technique to evaluate (bio)available trace metal concentrations in river water. – *Analytical and Bioanalytical Chemistry* 387: 2239–2244.
- Dmochowski W. (2005): Use of plant bioindicators in assessment of environmental contamination with heavy metals. – *Reports of the Botanical Garden of the Polish Academy of Sciences, Series: Monographs and Treatises* 7: 1–138.
- Dolník C. (2006): Different gemma formation in *Bryum barnesii*. – *Lindbergia* 31: 63–77.
- Dovčiak M., Halpern C. B., Saracco J. F., Evans S. A. & Liguori D. A. (2006): Persistence of ground-layer bryophytes in a structural retention experiment: initial effects of level and pattern of overstory retention. – *Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière* 36: 3039–3052.
- Doyle W. T. & Stotler R. E. (2006): Contributions toward a bryoflora of California III. Keys and annotated species catalogue for liverworts and hornworts. – *Madroño* 53: 89–197.
- Draper I., Lara F., Albertos B., Garilleti R. & Mazimpaka V. (2006): Epiphytic bryoflora of the Atlas and AntiAtlas Mountains, including a synthesis of the distribution of epiphytic bryophytes in Morocco. – *Journal of Bryology* 28: 312–330.
- Duff R. J., Villareal J. C., Cargill D. C. & Renzaglia K. S. (2007): Progress and challenges toward developing a phylogeny and classification of the hornworts. – *Bryologist* 110: 214–243.
- Dunn J. L. & Robinson S. A. (2006): Ultraviolet B screening potential is higher in two cosmopolitan moss species than in a co-occurring Antarctic endemic moss: implications of continuing ozone depletion. – *Global Change Biology* 12: 2282–2296.
- During H. J. (2006): Trends in bryophyte population dynamics. – *Lindbergia* 31: 6–15.
- During H. J., Verduyn B. & van Tooren B. F. (2006): On the increase of *Trematodon ambiguus* in lowland Belgium and the Netherlands. – *Lindbergia* 31: 101–108.
- Dynesius M. & Hylander K. (2007): Resilience of bryophyte communities to clear-cutting of boreal stream-side forests. – *Biological Conservation* 135: 423–434.
- Ellwood N. T. W. & Whitton B. A. (2007): Phosphatase activities of the aquatic moss *Warnstorffia fluitans* (Hedw.) Loeske from an acidic stream in North-East England. – *Hydrobiologia* 575: 95–107.
- Eppley S. M., Taylor P. T. & Jesson L. K. (2007): Self-fertilization in mosses: a comparison of heterozygote deficiency between species with combined versus separate sexes. – *Heredity* 98: 38–44.

- Fältmarsch R., Peltola P., Åström M. & Raitio H. (2007): Abundance, correlations and spatial patterns of nutrients and metals in till, humus, moss and pine needles in a boreal forest, western Finland. – *Geochemistry: Exploration, Environment, Analysis* 7: 57–69.
- Fattash I., Voss B., Reski R., Hess W. R. & Frank W. (2007): Evidence for the rapid expansion of microRNA-mediated regulation in early land plant evolution. – *BMC Plant Biology* 7: 13.
- Fedosov V. E. & Ignatova E. A. (2007 [2006]): The genus *Pseudocrossidium* R.S.Williams (*Pottiaceae, Musci*) in Russia. – *Arctoa* 15: 203–210.
- Feldberg K. & Heinrichs J. (2007):
- Feldberg K., Hentschel J., Wilson R., Rycroft D. S., Glenny D. & Heinrichs J. (2007): Phylogenetic biogeography of the leafy liverwort *Herbertus* (*Jungermanniales, Herbertaceae*) based on nuclear and chloroplast DNA sequence data: correlation between genetic variation and geographical distribution. – *Journal of Biogeography* 34: 688–698.
- Fernández J. A., Aboal J. R., Real C. & Carballeira A. (2007): A new moss biomonitoring method for detecting sources of small scale pollution. – *Atmospheric Environment* 41: 2098–2110.
- Finka A., Schaefer D. G., Saidi Y., Goloubinoff P. & Zryd J.-P. (2007): In vivo visualization of F-actin structures during the development of the moss *Physcomitrella patens*. – *New Phytologist* 174: 63–76.
- Fisher K. M., Wall D. P., Yip K. L. & Mishler B. D. (2007): Phylogeny of the *Calymperaceae* with a rank-free systematic treatment. – *Bryologist* 110: 46–73.
- Floyd S. K. & Bowman J. L. (2007): The ancestral developmental tool kit of land plants. – *International Journal of Plant Sciences* 168: 1–35.
- Frahm J.-P. (2006): Deutsche bryologische Literatur 1990–2005. – *Limprichtia* 29: 131–153.
- Freestone A. L. (2006): Facilitation drives local abundance and regional distribution of a rare plant in a harsh environment. – *Ecology* 87: 2728–2735.
- Freitas M. C., Pacheco A. M. G., Vieira B. J. & Rodrigues A. F. (2006): Neutron activation analysis of atmospheric biomonitoring from the Azores: A comparative study of lower and higher plants. – *Journal of Radioanalytical and Nuclear Chemistry* 270: 21–27.
- Friedel A., Oheimb G. v., Dengler J. & Härdtle W. (2006): Species diversity and species composition of epiphytic bryophytes and lichens – a comparison of managed and unmanaged beech forests in NE Germany. – *Feddes Repertorium* 117: 172–185.
- Galuszka A. (2007): Distribution patterns of PAHs and trace elements in mosses *Hylocomium splendens* (Hedw.) B.S.G. and *Pleurozium schreberi* (Brid.) Mitt. from different forest communities: A case study, south-central Poland. – *Chemosphere* 67: 1415–1422.
- Garilletti R., Lara F. & Mazimpaka V. (2007): Typification of twenty-six varieties of *Orthotrichum* (*Orthotrichaceae, Bryopsida*) published by G. Venturi. – *Nova Hedwigia* 84: 503–512.
- Glenn A. J., Flanagan L. B., Syed K. H. & Carlson P. J. (2006): Comparison of net ecosystem CO<sub>2</sub> exchange in two peatlands in western Canada with contrasting dominant vegetation, *Sphagnum* and *Carex*. – *Agricultural and Forest Meteorology* 140: 115–135.
- Górski P. (2007): Expansion of the liverwort *Tetralophozia setiformis* in the Polish Tatra Mts. (Western Carpathians). – *Acta Societatis Botanicorum Poloniae* 76: 75–79.
- Gough L. P., Eppinger R. G., Briggs P. H. & Giles S. (2006): Biogeochemical characterization of an undisturbed highly acidic, metal-rich bryophyte habitat, East-Central Alaska, USA. – *Arctic, Antarctic, and Alpine Research* 38: 522–529.
- Gradstein S. R. (2006): Stephani's Species Hepaticarum revisited. – *Willdenowia* 36 (Spec. Iss.) 557–563.
- Groth-Malonek M., Wahrmund U., Polsakiewicz M. & Knoop V. (2007): Evolution of a pseudogene: Exclusive survival of a functional mitochondrial *nad7* gene supports *Haplomitrium* as the earliest liverwort lineage and proposes a secondary loss of RNA editing in *Marchantiidae*. – *Molecular Biology and Evolution* 24: 1068–1074.
- Grundmann M., Ansell S. W., Russell S. J., Koch M. A. & Vogel J. C. (2007): Genetic structure of the widespread and common Mediterranean bryophyte *Pleurochaete squarrosa* (Brid.) Lindb. (*Pottiaceae*) – evidence from nuclear and plastidic DNA sequence variation and allozymes. – *Molecular Ecology* 16: 709–722.
- Gunnarsson U., Shaw A. J. & Lönn M. (2007): Local-scale genetic structure in the peatmoss *Sphagnum fuscum*. – *Molecular Ecology* 16: 305–312.
- Gunnarsson U. & Söderström L. (2007): Can artificial introductions of diaspore fragments work as a conservation tool for maintaining populations of the rare peatmoss *Sphagnum angermanicum*? – *Biological Conservation* 135: 450–458.

- Guo S. L., Enroth J. & Koponen T. (2007): Bryophyte flora of Hunan Province, China. 11. *Orthotrichaceae (Musci)*. – Annales Botanici Fennici 44 (Suppl. A): 1–34.
- Hájek M., Horská M., Hájková P. & Dítě D. (2006): Habitat diversity of central European fens in relation to environmental gradients and an effort to standardise fen terminology in ecological studies. – Perspectives in Plant Ecology, Evolution and Systematics 8: 97–114.
- Hájková P., Hájek M. & Apostolova I. (2006): Diversity of wetland vegetation in the Bulgarian high mountains, main gradients and context-dependence of the pH role. – Vegetatio 184: 111–130.
- Hájková P. & Hájek M. (2007): *Sphagnum* distribution patterns along environmental gradients in Bulgaria. – Journal of Bryology 29: 18–26.
- Hallingbäck T. (2007): Working with Swedish cryptogam conservation. – Biological Conservation 135: 334–340.
- Hallingbäck T., Lönnell N., Weibull H., Hedenäs L. & von Knorring P. (2006): Nationalnyckeln till Sveriges flora och fauna. Bladmossor: Sköldmossor – blåmossor. Bryoflora: *Buxbaumia* – *Leucobryum*. – ArtDatabanken, SLU, Uppsala. [416 pp.]
- Hartmann F. A., Wilson R., Gradstein S. R., Schneider H. & Heinrichs J. (2006): Testing hypotheses on species delimitations and disjunctions in the liverwort *Bryopteris* (*Jungermanniopsida*: *Lejeuneaceae*). – International Journal of Plant Sciences 167: 1205–1214.
- Hedderson T. A. & Blockeel T. L. (2006): *Oncophorus dendrophilus*, a new moss species from Cyprus and Crete. – Journal of Bryology 28: 357–359.
- Hedderson T. A. & Nowell T. L. (2006): Phylogeography of *Homalothecium sericeum* (Hedw.) Br. Eur.; toward a reconstruction of glacial survival and postglacial migration. – Journal of Bryology 28: 283–292.
- Hedenäs L. (2007): Global diversity patterns among pleurocarpous mosses. – Bryologist 110: 319–331.
- Heinrichs J., Hentschel J., Wilson R., Feldberg K. & Schneider H. (2007): Evolution of leafy liverworts (*Jungermanniidae*, *Marchantiophyta*): estimating divergence times from chloroplast DNA sequences using penalized likelihood with integrated fossil evidence. – Taxon 56: 31–44.
- Heras P., Infante M. & Buck W. R. (2006): On the presence of *Clasmatodon parvulus* (*Bryopsida*) in Europe. – Herzogia 19: 317–321.
- Hernández-Maqueda R., Quandt D., Werner O. & Muñoz J. (2007): Chloroplast data reveal two conflicting hypotheses for the positions of *Campylostelium* and *Grimmia pitardii* (*Bryophyta*). – Taxon 56: 89–94.
- Heslewood M. M. & Brown E. A. (2007): A molecular phylogeny of the liverwort family *Lepidoziaceae* Limpr. in Australasia. – Plant Systematics and Evolution 265: 193–219.
- Holz I. (2006): Epiphytic communities of bryophytes and macrolichens in a Costa Rican montane oak forest. – In: Kappelle M., ed., Ecology and conservation of Neotropical montane oak forests, Ecological Studies 185: 83–98.
- Humbert L., Gagnon D., Kneeshaw D. & Messier C. (2007): A shade tolerance index for common understory species of northeastern North America. – Ecological Indicators 7: 195–207.
- Hylander K. & Jonsson B. G. (2007): The conservation ecology of cryptogams. – Biological Conservation 135: 311–314.
- Ignatov M. S., Afonina O. M. & Ignatova E. A. (eds.) (2007 [2006]): Check-list of mosses of East Europe and North Asia. – Arctoa 15: 1–130.
- Ignatova E. A., Maksimov A., Maksimova T. & Belkina O. (2007 [2006]): Notes on distribution of *Schistidium* species (*Grimmiaceae*, *Bryophyta*) in Murmansk Province and Karelia. – Arctoa 15: 237–247.
- Ingerpuu N. & Kupper T. (2007): Response of calcareous grassland vegetation to mowing and fluctuating weather conditions. – Journal of Vegetation Science 18: 141–146.
- Ingerpuu N., Vellak K. & Möls T. (2007): Growth of *Neckera pennata*, an epiphytic moss of old-growth forests. – Bryologist 110: 309–318.
- Itouga M., Komatsu-Kato Y., Yamaguchi I., Ono Y. & Sakakibara H. (2006): Phytoremediation using bryophytes, 2. Bryo-filtration of copper in water using two species of *Scopelophila*. – Hikobia 14: 413–418.
- Jägerbrand A. K., Lindblad K. E. M., Björk R. G., Alatalo J. M. & Molau U. (2006): Bryophyte and lichen diversity under simulated environmental change compared with observed variation in unmanipulated alpine tundra. – Biodiversity and Conservation 15: 4453–4475.

- Juslén A. (2006): Phylogeny of *Vetaformaceae*, *Lepicoleaceae* and *Herbertaceae* (including *Mastigophoraceae*) inferred from chloroplast *trnL-F*, nuclear ITS2, and morphology. – *Annales Botanici Fennici* 43: 349–362.
- Juslén A. (2006): Revision of Asian *Herbertus* (*Herbertaceae*, *Marchantiophyta*). – *Annales Botanici Fennici* 43: 409–436.
- Köckinger H., Bednarek-Ochyra H. & Ochyra R. (2007): *Bucklandiella nivalis* (*Grimmiaceae*), a new moss from the Alps of Austria. – *Bryologist* 110: 92–99.
- Köckinger H. & Kučera J. (2007): *Barbula amplexifolia* (Mitt.) A.Jaeger in Europe. – *Journal of Bryology* 29: 33–40.
- Kooijman A. M. & Paulissen M. P. C. P. (2006): Higher acidification rates in fens with phosphorus enrichment. – *Applied Vegetation Science* 9: 205–212.
- Koranda M., Kerschbaum S., Wanek W., Zechmeister H. & Richter A. (2007): Physiological responses of bryophytes *Thuidium tamariscinum* and *Hylocomium splendens* to increased nitrogen deposition. – *Annals of Botany* 99: 161–169.
- Kučera J. (2007): New hepatic keys for Europe [Review of: Schumacker R. & Váňa J. 2005. Identification Keys to the Liverworts and Hornworts of Europe and Macaronesia (Distribution and Status), 2nd ed.]. – *Bryologist* 110: 335–336.
- Kürschner H. (2007): A key to the *Pottiaceae* (*Bryopsida* – *Bryophytina*) of the Near and Middle East. Towards a bryophyte flora of the Near and Middle East, 6. – *Nova Hedwigia* 84: 21–50.
- Kürschner H., Parolly G., Erdağ A. & Eren Ö. (2007): Synanthropic bryophyte communities new to western Turkey – syntaxonomy, synecology and life syndromes. – *Nova Hedwigia* 84: 459–478.
- Kürschner H., Stech M., Fontinha S., Sim-Sim M. & Frey W. (2007): Epiphytic bryophyte communities of the Madeiran laurel and ericaceous forests – a phytosociological analysis. – *Nova Hedwigia* 84: 333–362.
- Lara F., San Miguel E. & Mazimpaka V. (2006): Mosses and other plants used in nativity sets: a sampling study in northern Spain. – *Journal of Bryology* 28: 374–381.
- Larsen R. S., Bell J. N. B., James P. W., Chimonides P. J., Rumsey F. J., Tremper A. & Purvis O. W. (2007): Lichen and bryophyte distribution on oak in London in relation to air pollution and bark acidity. – *Environmental Pollution* 146: 332–340.
- Lavoie M., Pare D. & Bergeron Y. (2007): Quality of growth substrates of post-disturbed lowland black spruce sites for black spruce (*Picea mariana*) seedling growth. – *New Forests* 33: 207–216.
- Löhmus A., Löhmus P. & K. Vellak (2007): Substratum diversity explains landscape-scale co-variation in the species-richness of bryophytes and lichens. – *Biological Conservation* 135: 405–414.
- Long D. G. (2006): New higher taxa of complex thalloid liverworts (*Marchantiophyta* – *Marchantiopsida*). – *Edinburgh Journal of Botany* 63: 257–262.
- Longton R. E. (2006): Reproductive ecology of bryophytes: what does it tell us about the significance of sexual reproduction? – *Lindbergia* 31: 16–23.
- Malcolm B. & Malcolm N. (2006): Mosses and Other Bryophytes, An Illustrated Glossary. Second Edition. – Micro-Optics Press, Nelson, New Zealand. [iv + 336 pp.]
- Mälson K. & Rydin H. (2007): The regeneration capabilities of bryophytes for rich fen restoration. – *Biological Conservation* 135: 435–442.
- Mallón R., Barros P., Luzardo A. & González M. L. (2007): Encapsulation of moss buds: an efficient method for the in vitro conservation and regeneration of the endangered moss *Splachnum ampullaceum*. – *Plant Cell, Tissue and Organ Culture* 88: 41–49.
- Manzke W. (2006): Das Kugel-Hornmoos *Notothylas orbicularis* im Vogelsberg (Hessen): Verbreitung, Ökologie, Gefährdung, Forschungsbedarf. – *Limprichtia* 29: 57–88.
- Mathews S. (2006): Phytochrome-mediated development in land plants: red light sensing evolves to meet the challenges of changing light environments. – *Molecular Ecology* 15: 3483–3503.
- Mineyuki Y. (2007): Plant microtubule studies: past and present. – *Journal of Plant Research* 120: 45–51.
- Müller F. (2006): Bryologische Beobachtungen im Erzgebirge II. – *Limprichtia* 29: 155–160.
- Natcheva R. & Cronberg N. (2007): Recombination and introgression of nuclear and chloroplast genomes between the peat mosses, *Sphagnum capillifolium* and *Sphagnum quinquefarium*. – *Molecular Ecology* 16: 811–818.
- Natcheva R. & Ganeva A. (2006): Bryophytes on loess cliffs in Bulgaria: a preliminary study. – *Phytologia Balcanica* 12: 47–50.

- Natcheva R., Ganeva A. & Spiridonov G. (2006): Red List of the bryophytes in Bulgaria. – *Phytologia Balcanica* 12: 55–62.
- Niemelä M., Piispanen J., Poikolainen J. & Perämäki P. (2007): Preliminary study of the use of terrestrial moss (*Pleurozium schreberi*) for biomonitoring traffic-related Pt and Rh deposition. – *Archives of Environmental Contamination and Toxicology* 52: 347–354.
- Nishiyama T. (2007): Evolutionary developmental biology of nonflowering land plants. – *International Journal of Plant Sciences* 168: 37–47.
- Nordén B., Paltto H., Götmark F. & Wallin K. (2007): Indicators of biodiversity, what do they indicate? – Lessons for conservation of cryptogams in oak-rich forest. – *Biological Conservation* 135: 369–379.
- Oesau A. (2006): *Phascum vlassovii* Laz. (*Pottiaceae, Bryophytina*) in Hessen und Rheinland-Pfalz, neu für Mitteleuropa. – *Archive for Bryology* 18: 1–6.
- Oliván G., Fuertes E. & Acón M. (2007): *Hygrohypnum* (*Amblystegiaceae, Bryopsida*) in the Iberian Peninsula. – *Cryptogamie Bryologie* 28: 109–143.
- Olsson B. A. & Kellner O. (2006): Long-term effects of nitrogen fertilization on ground vegetation in coniferous forests. – *Forest Ecology and Management* 237: 458–470.
- Omar H. A., Aziz M. & Shakir K. (2007): Adsorption of U(VI) from dilute aqueous solutions onto peat moss. – *Radiochimica Acta* 95: 17–24.
- Ordóñez A., Gallego M. T., Guerra J. (2007): A preliminary account of the peristome structure and spores in six species of *Philonotis* (*Bartramiaceae, Bryophyta*). – *Cryptogamie Bryologie* 28: 15–29.
- Osyczka P., Dutkiewicz E. M. & Olech M. (2007): Trace elements concentrations in selected moss and lichen species collected within Antarctic research stations. – *Polish Journal of Ecology* 55: 39–48.
- Otte V. (2006): Erster Nachweis von *Frullania bolanderi* (*Marchantiophyta: Jubulaceae*) im Kaukasus. – *Herzogia* 19: 353–355.
- Paltto H., Nordén B., Götmark F. & Franc N. (2006): At which spatial and temporal scales does landscape context affect local density of Red Data Book and Indicator species?. – *Biological Conservation* 133: 442–454.
- Pan C., Shangguan Z. & Lei T. (2006): Influences of grass and moss on runoff and sediment yield on sloped loess surfaces under simulated rainfall. – *Hydrological Processes* 20: 3815–3824.
- Parker J. D., Burkepile D. E., Collins D. O., Kubanek J. & Hay M. E. (2007): Stream mosses as chemically-defended refugia for freshwater macroinvertebrates. – *Oikos* 116: 302–312.
- Partyka L. Ya. (2005): Brionflora Kryma. – Phytosociocentre, Kiev. [170 pp.]
- Peat H. J., Clarke A. & Convey P. (2007): Diversity and biogeography of the Antarctic flora. – *Journal of Biogeography* 34: 132–146.
- Pereira A. L. D. & Porto K. C. (2007): Patch size and isolation effects on epiphytic and epiphyllous bryophytes in the fragmented Brazilian Atlantic forest. – *Biological Conservation* 134: 415–427.
- Pesch R. & Schröder W. (2006): Integrative exposure assessment through classification and regression trees on bioaccumulation of metals, related sampling site characteristics and ecoregions. – *Ecological Informatics* 1: 55–65.
- Pesch R., Schröder W. & Schmidt G. (2007): Nitrogen accumulation in forests. Exposure monitoring by mosses. – *TheScientificWorldJOURNAL* 7: 151–158.
- Petraglia A. & Tomaselli M. (2007): Phytosociological study of the snowbed vegetation in the Northern Apennines (Northern Italy). – *Phytocoenologia* 37: 67–98.
- Pharo E. J. & Zartman C. E. (2007): Bryophytes in a changing landscape: The hierarchical effects of habitat fragmentation on ecological and evolutionary processes. – *Biological Conservation* 135: 315–325.
- Piippo S. & Kalinauskaitė N. (2006): Floristic elements of the hepatic flora from northern Finland to Lithuania. – *Memoranda Societatis pro Fauna et Flora Fennica* 82: 1–9.
- Pisarenko O. Yu. (2007) [2006]: On the variation and ecology of *Pterygoneurum subsessile* and *P. kozlovi* (*Pottiaceae, Bryophyta*). – *Arctoa* 15: 169–182.
- Pócs T. (2006): Bryophyte colonization and speciation on oceanic islands: an overview. – *Lindbergia* 31: 54–62.
- Proctor M. C. F., Ligrone R. & Duckett J. G. (2007): Desiccation tolerance in the moss *Polytrichum formosum*: Physiological and fine-structural changes during desiccation and recovery. – *Annals of Botany* 99: 75–93.
- Provan J. & Wilson P. J. (2007): Development of microsatellites for the peat moss *Sphagnum capillifolium* using ISSR cloning. – *Molecular Ecology Notes* 7: 254–256.

- Prudencio M. I. (2007): Biogeochemistry of trace and major elements in a surface environment (volcanic rock, soil, mosses, lichens) in the S. Miguel Island, Azores, Portugal. – Journal of Radioanalytical and Nuclear Chemistry 271: 431–437.
- Quatrano R. S., McDaniel S. F., Khandelwal A., Perroud P.-F. & Cove D. J. (2007): *Physcomitrella patens*: mosses enter the genomic age. – Current Opinion in Plant Biology 10: 182–189.
- Rau S., Miersch J., Neumann D., Weber E. & Krauss G.-J. (2007): Biochemical responses of the aquatic moss *Fontinalis antipyretica* to Cd, Cu, Pb and Zn determined by chlorophyll fluorescence and protein levels. – Environmental and Experimental Botany 59: 299–306.
- Renzaglia K. S., Schuette S., Duff R. J., Ligrone R., Shaw A. J., Mishler B. D. & Duckett J. G. (2007): Bryophyte phylogeny: Advancing the molecular and morphological frontiers. Bryologist 110: 179–213.
- Rice S. K. (2006): Towards an integrated understanding of bryophyte performance: the dimensions of space and time. – Lindbergia 31: 42–53.
- Rippy J. F. M. & Nelson P. V. (2007): Cation exchange capacity and base saturation variation among Alberta, Canada, moss peats. – HortScience 42: 349–352.
- Roads E. & Longton R. E. (2006): Year to year variation in the small scale distribution of shuttle mosses. – Lindbergia 31: 63–77.
- Robroek B. J. M., Limpens J., Breeuwer A. & Schouten M. G. C. (2007): Effects of water level and temperature on performance of four Sphagnum mosses. – Plant Ecology 190: 97–107.
- Ros R. M., Rams S., Holyoak D. T., Brugués M. & Cros R. M. (2007): Revision of the records of *Brachymenium commutatum* (*Bryophyta, Bryaceae*) in Europe. – Nova Hedwigia 84: 241–247.
- Ruokolainen L. & Salo K. (2006): The succession of boreal forest vegetation during ten years after slash-burning in Koli National Park, eastern Finland. – Annales Botanici Fennici 43: 363–378.
- Rydin H., Gunnarsson U. & Sundberg S. (2006): The role of *Sphagnum* in peatland development and persistence. – Ecological Studies 188: 47–65.
- Rydin H. & Jeglum J. K. (2006): The Biology of Peatlands. – Oxford University Press, Oxford. [ix + 343 pp.]
- Sabovljević M. & Natcheva R.. (2006): Check-list of the liverworts and hornworts of Southeast Europe. – Phytologia Balcanica 12: 169–180.
- Samecka-Cymerman A. & Kempers A. J. (2007): Differences in concentrations of heavy metals between native and transplanted *Pohlia nutans* (Hedw.) Lindb. – a case study from a dump exposed to industrial emissions in Poland. – Polish Journal of Environmental Studies 16: 251–258.
- Samecka-Cymerman A., Stankiewicz A., Kolon K. & Kempers A. J. (2007): Self-organizing feature map (neural networks) as a tool in classification of the relations between chemical composition of aquatic bryophytes and types of streambeds in the Tatra national park in Poland. – Chemosphere 67: 954–960.
- Schröder W. & Pesch R. (2007): Synthesizing bioaccumulation data from the German metals in mosses surveys and relating them to ecoregions. – Science of the Total Environment 374: 311–327.
- Sedia E. G. & Ehrenfeld J. G. (2006): Differential effects of lichens and mosses on soil enzyme activity and litter decomposition. – Biology and Fertility of Soils 43: 177–189.
- Selkirk P. M. & Skotnicki M. L. (2007): Measurement of moss growth in continental Antarctica. – Polar Biology 30: 407–413.
- Sérgio C., Brugués M., Cros R. M., Casas C. & Garcia C. (2006): The 2006 Red List and an updated checklist of bryophytes of the Iberian Peninsula (Portugal, Spain and Andorra). – Lindbergia 31: 109–126.
- Sérgio C., Figueira R., Draper D., Menezes R. & Sousa A. J. (2007): Modelling bryophyte distribution based on ecological information for extent of occurrence assessment. – Biological Conservation 135: 341–351.
- Shantz M. A. & Price J. S. (2006): Hydrological changes following restoration of the Bois-des-Bel Peatland, Quebec, 1999–2002. – Journal of Hydrology 331: 543–553.
- Shigyo M., Tabei N., Yoneyama T. & Yanagisawa S. (2007): Evolutionary processes during the formation of the plant-specific Dof transcription factor family. – Plant and Cell Physiology 48: 179–185.
- Shimamura M., Furuki T. & Deguchi H. (2006): Sporophyte anatomy of *Blasia pusilla* (*Blasiales, Blasiaceae*). – Hikobia 14: 399–402. [In Japanese with English abstract]
- Shotbolt L., Baker P. & Ashmore M. R. (2007): Reconstructing temporal trends in heavy metal deposition: Assessing the value of herbarium moss samples. – Environmental Pollution 147: 120–130.
- Singh M., Rawat A. K. S. & Govindarajan R. (2007): Antimicrobial activity of some Indian mosses. – Fitoterapia 78: 156–158.

- Smykla J., Wótek J. & Barcikowski A. (2007): Zonation of vegetation related to penguin rookeries on King George Island, Maritime Antarctic. – Arctic Antarctic and Alpine Research 39: 143–151.
- Söderström L. (2006): Conservation biology of Bryophytes. – Lindbergia 31: 24–32.
- Söderström L., Séneca A. & Santos M. (2007): Rarity patterns in members of the *Lophoziaceae/Scapaniaceae* complex occurring North of the Tropics – Implications for conservation. – Biological Conservation 135: 352–359.
- Solga A. (2007): Seasonal variation in the nitrogen concentration and  $^{15}\text{N}$  natural abundance of a pleurocarpous moss species in dependence on nitrogen deposition dynamics. Cryptogamie Bryologie 28: 93–102.
- Solga A., Eichert T. & Frahm J.-P. (2006): Historical alteration in the nitrogen concentration and  $^{15}\text{N}$  natural abundance of mosses in Germany: Indication for regionally varying changes in atmospheric nitrogen deposition within the last 140 years. – Atmospheric Environment 40: 8044–8055.
- Sotiaux A., Pioli A., Royaud A., Schumacker R. & Vanderpoorten A. (2007): A checklist of the bryophytes of Corsica (France): new records and a review of the literature. – Journal of Bryology 29: 41–53.
- Spagnuolo V., Muscariello L., Cozzolino S., Cobianchi R. C. & Giordano S. (2007): Ubiquitous genetic diversity in ISSR markers between and within populations of the asexually producing moss *Pleurochaete squarrosa*. – Plant Ecology 188: 91–101.
- Standovár T., Ódor P., Aszalós R. & Gálhidy L. (2006): Sensitivity of ground layer vegetation diversity descriptors in indicating forest naturalness. – Community Ecology 7: 199–209.
- Stark L. R., Oliver M. J., Mishler B. D. & McLetchie D. N. (2007): Generational differences in response to desiccation stress in the desert moss *Tortula inermis*. – Annals of Botany 99: 53–60.
- Startsev N. A. & Lieffers V. J. (2006): Dynamics of mineral nitrogen released from feathermosses after dehydration or handling stress. – Bryologist 109: 551–559.
- Startsev N. A., Lieffers V. J. & McNabb D. H. (2007): Effects of feathermoss removal, thinning and fertilization on lodgepole pine growth, soil microclimate and stand nitrogen dynamics. – Forest Ecology and Management 240: 79–86.
- Starzomski B. M. & Srivastava D. S. (2007): Landscape geometry determines community response to disturbance. – Oikos 116: 690–699.
- Stevens M. I., Hunger S. A., Hills S. F. K. & Gemmill C. E. C. (2007): Phantom hitch-hikers mislead estimates of genetic variation in Antarctic mosses. – Plant Systematics and Evolution 263: 191–201.
- Szczepaniak K., Astel A., Bode P., Sarbu C., Biziuk M., Rainska E. & Gos K. (2006): Assessment of atmospheric inorganic pollution in the urban region of Gdańsk, Northern Poland. – Journal of Radioanalytical and Nuclear Chemistry 270: 35–42.
- Szövényi P., Hock Z., Urmi E. & Schneller J. (2006): New primers for amplifying the GapC gene in bryophytes and its utility in intraspecific phylogenies in the genus *Spagnum*. – Lindbergia 31: 78–84.
- Štechová T. & Kučera J. 2007. The requirements of the rare moss, *Hamatocaulis vernicosus* (*Calliergonaceae, Musci*), in the Czech Republic in relation to vegetation, water chemistry and management. – Biological Conservation 135: 443–449.
- Tan B. C., Koponen T. & Norris D. H. (2007): Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXX. *Sematophyllaceae* (*Musci*) 1. *Acanthorrhynchium*, *Acroporium*, *Clastobryophilum*, *Pseudopilocium*, *Radulina* and *Trichosteleum*. — Annales Botanici Fennici 44 (Suppl. A): 35–78.
- Terasawa K., Odahara M., Kabeya Y., Kikugawa T., Sekine Y., Fujiwara M. & Sato N. (2007): The mitochondrial genome of the moss *Physcomitrella patens* sheds new light on mitochondrial evolution in land plants. – Molecular Biology and Evolution 24: 699–709.
- Tretiach M., Adamo P., Bargagli R., Baruffo L., Carletti L., Crisafulli P., Giordano S., Modenesi P., Orlando S. & Pittao E. (2007): Lichen and moss bags as monitoring devices in urban areas. Part I: Influence of exposure on sample vitality. – Environmental Pollution 146: 380–391.
- Ueno T., Bekku Y., Uchida M. & Kanda H. (2006): Photosynthetic light responses of a widespread moss, *Sanionia uncinata*, from contrasting water regimes in the high Arctic tundra, Svalbard, Norway. – Journal of Bryology 28: 345–349.
- Uyar G., Ören M. & Ince M. (2007): Atmospheric heavy metal deposition in Duzce province by using mosses as biomonitoring. – Fresenius Environmental Bulletin 16: 145–153.
- Uyar G., Ören M., Yıldırım Y. & Ince M. (2007): Mosses as indicators of atmospheric heavy metal deposition around a coal-fired power plant in Turkey. – Fresenius Environmental Bulletin 16: 182–192.

- Van Sandt V. S. T., Stiepераere H., Guisez Y., Verbelen J.-P. & Vissenberg K. (2007): XET activity is found near sites of growth and cell elongation in bryophytes and some green algae: New insights into the evolution of primary cell wall elongation. – Annals of Botany 99: 39–51.
- Vanderpoorten A. & Goffinet B. (2006): Mapping uncertainty and phylogenetic uncertainty in ancestral character state reconstruction: An example in the moss genus *Brachytheciastrum*. – Systematic Biology 55: 957–971.
- Vellak K., Vellak A. & Ingerpuu N. (2007): Reasons for moss rarity: Study in three neighbouring countries. – Biological Conservation 135: 360–368.
- Villaseñor J. L., Delgadillo C. & Ortiz E. (2006): Biodiversity hotspots from a multigroup perspective: Mosses and Senecios in the Transmexican Volcanic Belt. – Biodiversity and Conservation 15: 4045–4058.
- Virtanen R. & Oksanen J. (2007): The effects of habitat connectivity on cryptogam richness in boulder metacommunity. – Biological Conservation 135: 415–422.
- Vitt D. H. (2006): Bryophyte community ecology: going beyond description. – Lindbergia 31: 33–41.
- von Stackelberg M., Rensing S. A. & Reski R. (2006): Identification of genic moss SSR markers and a comparative analysis of twenty-four algal and plant gene indices reveal species-specific rather than group-specific characteristics of microsatellites. – BMC Plant Biology 6: –.
- Wallrup E., Saetre P. & Rydin H. (2006): Deciduous trees affect small-scale floristic diversity and tree regeneration in conifer forests. – Scandinavian Journal of Forest Research 21: 399–404.
- Werner O., Ros R. M., Goffinet B. (2007): A reconsideration of the systematic position of *Goniomitrium* (*Funariaceae*) based on chloroplast sequence markers. – Bryologist 110: 108–114.
- Wood A. J. (2007): Invited essay: New frontiers in bryology and lichenology. The nature and distribution of vegetative desiccation-tolerance in hornworts, liverworts and mosses. – Bryologist 110: 163–177.
- Zander R. H., Toren D. & Eckel P. M. (2007): *Gymnostomum aeruginosum*, *G. calcareum* and *G. viridulum* (*Pottiaceae*, *Bryopsida*) in California. – Journal of Bryology 29: 27–32.
- Zechmeister H. G., Dullinger S., Hohenwallner D., Riss A., Hanus-Illnar A. & Scharf S. (2006): Pilot study on road traffic emissions (PAHs, heavy metals) measured by using mosses in a tunnel experiment in Vienna, Austria. – Environmental Science and Pollution Research 13: 398–405.
- Zechmeister H. G., Hagendorfer H., Hohenwallner D., Hanus-Illnar A. & Riss A. (2006): Analyses of platinum group elements in mosses as indicators of road traffic emissions in Austria. – Atmospheric Environment 40: 7720–7732.
- Zippel E. (2006): Type material in the moss herbarium of Samuel Elisée von Bridel, 1. – Willdenowia 36: 913–935.
- Zolotov V. I. (2007 [2006]): On systematics and distribution of some species of *Bryum* (*Bryaceae*, *Bryophyta*) in Russia. – Arctoa 15: 155–162.

## **Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]**

### **1. Články a knižní publikace [Papers and books]**

- Hájek P. (2006): Contribution to the knowledge of the bryophyta (*Bryophyta*) in locality Dubno near Česká Skalice. – Acta Musei Reginae-Hradecensis, Ser. A, Scientiae Naturales 31: 29–31.
- Holá E. (2006): Bryoflóra horního toku Křemelné na Šumavě [Bryoflora of the Upper Křemelná River in the Bohemian Forest]. – Silva Gabreta 12: 109–131.
- Kubešová S. (2006): Zelení hosté v historickém podzemí měst. – Veronica 20: 25–26.
- Kučera J. (2006): Nová bryologická literatura XVI. [New bryological literature, XVI]. – Bryonora 38: 64–72.
- Kučera J. (2006): Úmrtí [Obituaries]. – Bryonora 38: 56. [S. M. Såstad]
- Kučera J. (ed.) (2006): Zajímavé bryofloristické nálezy VIII. [Interesting bryofloristic records, VIII]. – Bryonora 38: 47–52.
- Kučera J., Müller F. & Marková I. (2006): Mechorsty zaznamenané v průběhu 19. podzimního setkání Bryologicko-lichenologické Sekce v CHKO Kokořínsko [Bryophytes recorded during the 19th Autumn Meeting of the Bryological and Lichenological Section in the Kokořínsko region (Central Bohemia)]. – Bryonora 38: 18–25.

- Löbel S., Dengler J. & Hobohm C. (2006): Species richness of vascular plants, bryophytes and lichens in dry grasslands: The effects of environment, landscape structure and competition. – *Folia Geobotanica* 41: 377–393.
- Mikulášková E.: Vývoj rozšíření neofytického mechu *Campylopus introflexus* v České republice [Development in distribution of the neophytic moss *Campylopus introflexus* in the Czech Republic]. – *Bryonora* 38: 1–10.
- Navrátilová J., Navrátil J. & Hájek M. (2006): Relationships between environmental factors and vegetation in nutrient-enriched fens at fishpond margins. – *Folia Geobotanica* 41: 353–376.
- Plášek V., Zmrhalová M., Kubešová S., Szczepański M. & Loskotová E. (2006): Příspěvek k bryoflóře Tematínských vrchů. – In: Rajcová K., ed., *Najvzácnější přírodné hodnoty Tematínských vrchov, Trenčín*, p. 31–34.
- Váňa J. (2006): Rozšíření druhů rodů *Amblyodon* P. Beauv. a *Meesia* Hedw. v České republice [Distribution of the genera *Amblyodon* P. Beauv. and *Meesia* Hedw. in the Czech Republic]. – *Bryonora* 38: 10–18.

## 2. Recenze [Reviews]

- Kučera J. (2006): Recenze [Reviews]. – *Bryonora* 38: 57–58. [Frey W. & al. (2006): The Liverworts, Mosses and Ferns of Europe. – Harley Books, Colchester.]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 58–59. [Nebel M. & Philippi G. (eds.) (2005): Die Moose Baden-Württembergs, Bd. 3. – Eugen Ulmer, Stuttgart.]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 59–60. [Hallingbäck T. & al. (2006): Nationalnyckeln till Sveriges flora och fauna. Bladmossor: Sköldmossor – blåmossor. *Bryophyta: Buxbaumia – Leucobryum*. – ArtDatabanken, SLU, Uppsala]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 61–61. [Szwejkowski J. (2006): An Annotated Checklist of Polish Liverworts and Hornworts. – Biodiversity of Poland, vol. 4, W. Szafer Institute of Botany, Kraków.]

## 3. Bryologické rukopisné práce [Bryological manuscripts]

- Jandová J. (2007): Bryologické zhodnocení pramenišť Luzenského údolí (Šumava) [Bryological research results of spring area in Luzenské valley (Bohemian Forest)]. – Ms., 40 pp. [Bc. thesis, University of South Bohemia, České Budějovice]
- Košnar J. (2007): Taxonomická studie okruhu *Tortula muralis* ve střední Evropě [Taxonomic study of *Tortula muralis* complex in Central Europe]. – Ms., 46 pp. + App. [MSc. thesis, University of South Bohemia, České Budějovice]

## **NOVÁ LICHENOLOGICKÁ LITERATURA XVI.**

### **New lichenological literature, XVI**

Zdeněk Palice

*Botanický ústav AV ČR, CZ-252 43 Prihovice, e-mail: palice@ibot.cas.cz*

- Aguirre-Hudson M. B. & Spooner B. M. (eds.) (2005): New, rare and interesting lichens. – *British Lichen Society Bulletin* 97: 69–79.
- Aptroot A. (2006): Three new species of *Lithothelium* (*Pyrenulaceae*) from China and Thailand, with a revised world key and annotated list of species. – *Lichenologist* 38: 541–548.
- Aptroot A. & Honegger R. (2006): Lichens in the new botanical garden of the University of Zürich, Switzerland. – *Botanica Helvetica* 116: 135–148.
- Aptroot A., John V. & Wirth V. (2006): Flechten und lichenicole Pilze im Dreiländereck bei der Saarschleife mit Neufunden aus Lothringen, Saarland und Rheinland-Pfalz (BLAM-Exkursion 2005). – *Herzogia* 19: 63–76.