

- Löbel S., Dengler J. & Hobohm C. (2006): Species richness of vascular plants, bryophytes and lichens in dry grasslands: The effects of environment, landscape structure and competition. – *Folia Geobotanica* 41: 377–393.
- Mikulášková E.: Vývoj rozšíření neofytického mechu *Campylopus introflexus* v České republice [Development in distribution of the neophytic moss *Campylopus introflexus* in the Czech Republic]. – *Bryonora* 38: 1–10.
- Navrátilová J., Navrátil J. & Hájek M. (2006): Relationships between environmental factors and vegetation in nutrient-enriched fens at fishpond margins. – *Folia Geobotanica* 41: 353–376.
- Plášek V., Zmrhalová M., Kubešová S., Szczepański M. & Loskotová E. (2006): Příspěvek k bryoflóře Tematínských vrchů. – In: Rajcová K., ed., Najvzácnejšie prírodné hodnoty Tematínských vrchov, Trenčín, p. 31–34.
- Váňa J. (2006): Rozšíření druhů rodů *Amblyodon* P. Beauv. a *Meesia* Hedw. v České republice [Distribution of the genera *Amblyodon* P. Beauv. and *Meesia* Hedw. in the Czech Republic]. – *Bryonora* 38: 10–18.

2. Recenze [Reviews]

- Kučera J. (2006): Recenze [Reviews]. – *Bryonora* 38: 57–58. [Frey W. & al. (2006): The Liverworts, Mosses and Ferns of Europe. – Harley Books, Colchester.]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 58–59. [Nebel M. & Philippi G. (eds.) (2005): Die Moose Baden-Württembergs, Bd. 3. – Eugen Ulmer, Stuttgart.]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 59–60. [Hallingbäck T. & al. (2006): Nationalnyckeln till Sveriges flora och fauna. Bladmossor: Sköldmossor – blåmossor. *Bryophyta: Buxbaumia – Leucobryum*. – ArtDatabanken, SLU, Uppsala]
- Váňa J. (2006): Recenze [Reviews]. – *Bryonora* 38: 61–61. [Szweykowski J. (2006): An Annotated Checklist of Polish Liverworts and Hornworts. – Biodiversity of Poland, vol. 4, W. Szafer Institute of Botany, Kraków.]

3. Bryologické rukopisné práce [Bryological manuscripts]

- Jandová J. (2007): Bryologické zhodnocení pramenišť Luzenského údolí (Šumava) [Bryological research results of spring area in Luzenské valley (Bohemian Forest)]. – Ms., 40 pp. [Bc. thesis, University of South Bohemia, České Budějovice]
- Košnar J. (2007): Taxonomická studie okruhu *Tortula muralis* ve střední Evropě [Taxonomic study of *Tortula muralis* complex in Central Europe]. – Ms., 46 pp. + App. [MSc. thesis, University of South Bohemia, České Budějovice]

NOVÁ LICHENOLOGICKÁ LITERATURA XVI.

New lichenological literature, XVI

Zdeněk P a l i c e

Botanický ústav AV ČR, CZ-252 43 Průhonice, e-mail: palice@ibot.cas.cz

- Aguirre-Hudson M. B. & Spooner B. M. (eds.) (2005): New, rare and interesting lichens. – *British Lichen Society Bulletin* 97: 69–79.
- Aptroot A. (2006): Three new species of *Lithothelium* (Pyrenulaceae) from China and Thailand, with a revised world key and annotated list of species. – *Lichenologist* 38: 541–548.
- Aptroot A. & Honegger R. (2006): Lichens in the new botanical garden of the University of Zürich, Switzerland. – *Botanica Helvetica* 116: 135–148.
- Aptroot A., John V. & Wirth V. (2006): Flechten und lichenicole Pilze im Dreiländereck bei der Saarschleife mit Neufunden aus Lothringen, Saarland und Rheinland-Pfalz (BLAM-Exkursion 2005). – *Herzogia* 19: 63–76.

- Aptroot A. & Sparrius L. B. (2006): Additions to the lichen flora of Vietnam, with an annotated checklist and bibliography. – *Bryologist* 109: 358–371.
- Aptroot A., Umaña L., Chaves J. L. & Trest M. T. (2006): A first assessment of the Ticolichen biodiversity inventory in Costa Rica: three new squamulose genera (*Lecanorales*: *Ramalinaceae* and *Pilocarpaceae*). – *Journal of the Hattori Botanical Laboratory* 100: 617–623.
- Aragón G., García A. & López R. (2006): Aportación al conocimiento de los líquenes rupícolas que colonizan las pedrizas del Parque Nacional de Cabañeros. – *Botanica Complutensis* 30: 53–59.
- Aragón G., Martínez I. & Belinchón R. (2006): Aportación al conocimiento de los líquenes epífitos de *Pinus halepensis*, *P. nigra*, *P. pinaster* y *P. sylvestris* en la Península Ibérica. – *Botanica Complutensis* 30: 61–70.
- Aragón G., Rico V. J. & Belinchón R. (2006): Lichen diversity from Cazorla, Segura and Las Villas Biosphere Reserve (SE Spain). – *Nova Hedwigia* 82: 31–50.
- Archer A. W. (2006): The lichen family *Graphidaceae* in Australia. – *Bibliotheca Lichenologica* 94: 1–191.
- Arup U. (2006): A new taxonomy of the *Caloplaca citrina* group in the Nordic countries, except Iceland. – *Lichenologist* 38: 1–20.
- Arup U. (2006): *Caloplaca sorediella* Arup, a new sorediate species from western Britain. – *Lichenologist* 38: 499–502.
- Arup U. (2006): The status of *Caloplaca havaasii*. – *Graphis Scripta* 18: 33–37.
- Asta J., Gavériaux J.-P., Moreau P.-A., Rémy C., Sussey J.-M. & Van Haluwyn C. (2005): Numéro spécial lichens. – *Bulletin Mycologique et Botanique Dauphiné-Savoie* 178: 5–93.
- Averis B. (2006): Lichen communities in relation to bryophyte communities and the National Vegetation Classification. – *British Lichen Society Bulletin*. 98: 11–13.
- Baloch E. & Grube M. (2006): Evolution and phylogenetic relationships within *Porinaceae* (*Ostropomycetidae*), focusing on foliicolous species. – *Mycological Research* 110: 125–136.
- Barbero A., Giralt M., Elix J. A., Gómez-Bolea A. & Llimona X. (2006): A taxonomic study of *Protoparmelia montagnei* (syn. *P. psarophana*) centered in the Eastern Iberian Peninsula. – *Mycotaxon* 97: 299–320.
- Baruffo L., Zedda L., Elix J. A. & Tretiach M. (2006): A revision of the lichen genus *Lepraria* s.lat. in Italy. – *Nova Hedwigia* 83: 387–430.
- Beckett R. P., Mayaba N., Minibayeva F. V. & Alyabyev A. J. (2005): Hardening by partial dehydration and ABA increase desiccation tolerance in the cyanobacterial lichen *Peltigera polydactylon*. – *Annals of Botany* 96: 109–115.
- Behera B. C., Adawadkar B. & Makhija U. (2006): Tissue-culture of selected species of the *Graphis* lichen and their biological activities. – *Fitoterapia* 77: 208–215.
- Berryman S. & McCune B. (2006): Epiphytic lichens along gradients in topography and stand structure in western Oregon, USA. – *Pacific Northwest Fungi* 1: 1–38.
- Berryman S. & McCune B. (2006): Estimating epiphytic macrolichen biomass from topography, stand structure and lichen community data. – *Journal of Vegetation Science* 17: 157–170.
- Berthold S. & Otto P. (2005): Untersuchungen zur Diversität und Standortwahl von Pilzen und Flechten auf vegetationsbedeckten Flachdächern in Chemnitz (Sachsen). – *Botetus* 28: 37–47.
- Białońska D. & Dayan F. E. (2005): Chemistry of the lichen *Hypogymnia physodes* transplanted to an industrial region. – *Journal of Chemical Ecology* 31: 2975–2991.
- Bielczyk U. (2006): Distribution of calciphilous lichens in Poland. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 153–164, Mycotaxon Ltd., Ithaca.
- Bielczyk U. (2006): The lichen biota of Polish Carpathians – general characteristic. – *Polish Botanical Journal* 51: 1–24.
- Bielczyk U., Bylińska E., Czarnota P., Czyżewska K., Guzow-Krzemińska B., Hachułka M., Kiszka J., Kowalewska A., Krzewicka B., Kukwa M., Leśniański G., Śliwa L. & Zalewska A. (2005): Contribution to the knowledge of lichens and lichenicolous fungi of Western Ukraine. – *Polish Botanical Journal* 50: 39–64.
- Bjerke J. W., Elvebakk A. & Elverland E. (2006): The lichen genus *Usnea* in Norway north of the Arctic Circle: biogeography and ecology. – *Nova Hedwigia* 83: 293–310.
- Bjerke J. W. & Timdal E. (2006): The distribution of *Menegazzia subsimilis* in Norway. – *Graphis Scripta* 18: 58–61.

- Bjerke J. W., Tømmervik H. A., Finne T. E., Jensen H., Lukina N. & Bakkestuen V. (2006): Epiphytic lichen distribution and plant leaf heavy metal concentrations in Russian-Norwegian boreal forests influenced by air pollution from nickel-copper smelters. – *Boreal Environment Research* 11: 441–450.
- Blaha J., Baloch E. & Grube M. (2006): High photobiont diversity associated with the euryoecious lichen-forming ascomycete *Lecanora rupicola* (*Lecanoraceae*, *Ascomycota*). – *Biological Journal of the Linnean Society* 88: 283–293.
- Blanco O., Crespo A., Ree R. H. & Lumbsch H. T. (2006): Major clades of parmelioid lichens (*Parmeliaceae*, *Ascomycota*) and the evolution of their morphological and chemical diversity. – *Molecular Phylogenetics and Evolution* 39: 52–69.
- Boch S. & Sparrius L. B. (2006): Neue und interessante Flechtenfunde aus den Landkreisen Lüneburg und Lüchow-Dannenberg (Nordost-Niedersachsen, Deutschland). – *Herzogia* 19: 77–83.
- Bradwell T. & Armstrong R. A. (2006): Growth rates of *Rhizocarpon geographicum* lichens: a review with new data from Iceland. – *Journal of Quaternary Science* 22: 311–320.
- Breuss O. (2006): Bemerkenswerte Flechtenfunde aus Niederösterreich und der Steiermark 2. – *Österreichische Zeitschrift für Pilzkunde* 15: 121–126.
- Brunauer G., Hager A., Krautgartner W. D., Türk R. & Stocker-Wörgötter E. (2006): Experimental studies on *Lecanora rupicola* (L.) Zahlbr.: chemical and microscopical investigations of the mycobiont and re-synthesis stages. – *Lichenologist* 38: 577–585.
- Bültmann H. (2005): Strategien und Artenreichtum von Erdflechten in Sandtrockenrasen. – *Tuexenia* 25: 425–443.
- Bültmann H. (2005): Syntaxonomy of arctic terricolous lichen vegetation, including a case study from Southeast Greenland. – *Phytocoenologia* 35: 909–949.
- Burgaz A. R. (2006): Check-list of lichenized and lichenicolous fungi of Madrid Community (Spain). – *Flora Mediterranea* 16: 57–110.
- Buschbom J. & Barker D. (2006): Evolutionary history of vegetative reproduction in *Porpidia* s.l. (lichen-forming *Ascomycota*). – *Systematic Biology* 55: 471–484.
- Buschbom J. & Mueller G. M. (2006): Testing "species pair" hypotheses: evolutionary processes in the lichen-forming species complex *Porpidia flavocoerulescens* and *Porpidia melinodes*. – *Molecular Biology and Evolution* 23: 574–586.
- Cardinale M., Puglia A. M. & Grube M. (2006): Molecular analysis of lichen-associated bacterial communities. – *FEMS Microbiology Ecology* 57: 484–495.
- Ceynowa-Giełdon M. & Adamska E. (2005): *Caloplaca ruderum*, a lichen species new to Poland. – *Polish Botanical Journal* 50: 75–76.
- Chaujar R. K. (2006): Lichenometry of yellow *Rhizocarpon geographicum* as database for the recent geological activities in Himachal Pradesh. – *Current Science* 90: 1552–1554.
- Ciaciura M., Przybył I. & Bosiacka B. (2006): Porosty projektowanego rezerwatu krajobrazowego "Diabelskie Pustacie" na Pojezierzu Szczesineckim. – *Fragmenta Floristica et Geobotanica Polonica* 13: 183–190.
- Cieśliński S. & Czyżewska K. (2006): Changes in the lichen biota of the "Skalki Piekło pod Niekłanem" Nature Reserve and its surroundings (Central Poland) during the past 100 years. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 259–270, Mycotaxon Ltd., Ithaca.
- Clerc P. (2006): *Parmelia afrorevoluta* et *P. britannica*: deux nouvelles espèces de macrolichens pour la Suisse: une redéfinition de *Parmelia revoluta*! – *Meylania* 35: 6–15.
- Clerc P. (2006): Synopsis of *Usnea* (lichenized *Ascomycetes*) from the Azores with additional information on the species in Macaronesia. – *Lichenologist* 38: 191–212.
- Cooley D., Naveau P., Jomelli V., Rabatel A. & Grancher D. (2006): A Bayesian hierarchical extreme value model for lichenometry. – *Environmetrics* 17: 555–574.
- Coppins B. J. & Coppins A. M. (2005): Lichens - the biodiversity value of western woodlands. – *Botanical Journal of Scotland* 57: 141–153.
- Coppins B. J. & Coppins A. M. (2006): The lichens of the Scottish native pinewoods. – *Forestry* 79: 249–259.
- Coppins B. J. & Fryday A. M. (2006): New or previously misunderstood species of *Lithographa* and *Rimularia* (*Agyriaceae*) from the southern subpolar region and western Canada. – *Lichenologist* 38: 93–107.

- Coppins B. J. & Fryday A. M. (2006): Reassessment of some lichen species described by Josiah Lowe, and notes on some other North American lecideoid lichens. – *Bryologist* 109: 9–17.
- Coutin R. (2004): Chenilles mangeuses de lichens, de mousses et d'hépatiques. 1^{ère} partie. – *Insectes* 133: 33–37.
- Coutin R. (2004): Chenilles mangeuses de lichens, de mousses et d'hépatiques. 2^e partie. – *Insectes* 134: 35–37.
- Crespo A., Arguello A., Lumbsch H. T., Llimona X. & Tønsberg T. (2006): A new species of *Lepraria* (*Lecanorales: Stereocaulaceae*) from the Canary Islands and the typification of *Lepraria isidiata*. – *Lichenologist* 38: 213–221.
- Crewe A. T., Purvis O. W. & Wedin M. (2006): Molecular phylogeny of *Acarosporaceae* (*Ascomycota*) with focus on the proposed genus *Polysporinopsis*. – *Mycological Research* 110: 521–526.
- Czarnota P. & Coppins B. J. (2005): A second *Micareia* with a hypothecial K+ violet pigment. – *Lichenologist* 37: 477–479.
- Czarnota P. & Coppins B. J. (2006): A new *Bacidia* with long-necked pycnidia from Central Europe. – *Lichenologist* 38: 407–410.
- Czarnota P., Glanc K. & Nowak J. (2005): Materiały do bioty porostów Gorców ze zbiorów Herbarium Instytutu Botaniki Polskiej Akademii Nauk w Krakowie. – *Fragmenta Floristica et Geobotanica Polonica* 12: 327–370.
- Czyżewska K., Motiejūnaitė J. & Cieśliński S. (2005): New and noteworthy species of lichens and allied fungi from North-Eastern Poland. – *Acta Mycologica* 40: 277–291.
- Davydov E. A. (2006): *Umbilicaria altaiensis* (*Umbilicariaceae*) - novyj vid lishainika dlja Rossii, Mongolii i Gruzii. – *Botanicheskij Zhurnal* 91: 1260–1267.
- De Graef B., De Windt W., Dick J., Verstraete W. & De Belie N. (2005): Cleaning of concrete fouled by lichens with the aid of *Thiobacilli*. – *Materials and Structures* 38: 875–882.
- Diederich P., Van den Broeck D., Ertz D., Signoret J., Aptroot A., Sparrius L., Jordaens D. & Sérusiaux E. (2006): Contribution to the knowledge of lichens in northern France. – *Bulletin de la Société des Naturalistes Luxembourgeois* 106: 53–62.
- Dietrich M. (2006): Die mittelalterliche Museggmauer in Luzern (Zentralschweiz) und ihre Vielfalt an gesteinsbewohnenden Flechten. – *Meylania* 35: 16–21.
- Dolnik C. (2005): *Agonimia allobata* und Nachweise andere seltener Flechten aus Schleswig-Holstein. – *Kieler Notizen zur Pflanzenkunde* 33: 90–97.
- Dolnik C. (2005): *Bacidia etayana* on the German Baltic coast. – *Herzogia* 18: 219–222.
- Dolnik C., Zimmer D., Stolley G. & Boch S. (2006): *Fellhanera viridisorediata* und andere interessante Flechtenfunde aus Schleswig-Holstein. – *Kieler Notizen zur Pflanzenkunde* 34: 63–73.
- Domeño C., Blasco M., Sánchez C. & Nerín C. (2006): A fast extraction technique for extracting polycyclic aromatic hydrocarbons (PAHs) from lichens samples used as biomonitors of air pollution: dynamic sonication versus other methods. – *Analytica Chimica Acta* 569: 103–112.
- Dunford J. S., McLoughlin P. D., Dalerum F. & Boutin S. (2006): Lichen abundance in the peatlands of northern Alberta: Implications for boreal caribou. – *Ecoscience* 13: 469–474.
- Elix J. A. & Tønsberg T. (2005): On the chemistry of *Ramalina thrausta*. – *Graphis Scripta* 17: 35–36.
- Elix J. A. & Tønsberg T. (2006): Notes on the chemistry of Scandinavian *Lobaria* species. – *Graphis Scripta* 18: 27–28.
- Ellis C. J. & Coppins B. J. (2006): Contrasting functional traits maintain lichen epiphyte diversity in response to climate and autogenic succession. – *Journal of Biogeography* 33: 1643–1656.
- Elvebakk A. & Bjerke J. W. (2006): The Skibotn area in North Norway – An example of very high lichen species richness far to the north: A supplement with an annotated list of species. – *Mycotaxon* 96: 141–146.
- Esseen P.-A. (2006): Edge influence on the old-growth forest indicator lichen *Alectoria sarmentosa* in natural ecotones. – *Journal of Vegetation Science* 17: 185–194.
- Favero-Longo S. E., Turci F., Tomatis M., Castelli D., Bonfante P., Hochella M. F. Piervittori R. & Fubini B. (2005): Chrysotile asbestos is progressively converted into a non-fibrous amorphous material by the chelating action of lichen metabolites. – *Journal of Environmental Biomonitoring* 7: 764–766.
- Favero-Longo S. E., Siniscalco C. & Piervittori R. (2006): Plant and lichen colonization in an asbestos mine: Spontaneous bioattenuation limits air dispersion of fibres. – *Plant Biosystems* 140: 190–205.

- Fernández-Salegui A. B., Calatayud Á., Terrón A. & Barreno E. M. (2006): Chlorophyll *a* fluorescence in transplants of *Parmelia sulcata* Taylor near a power station (La Robla, León, Spain). – *Lichenologist* 38: 457–468.
- Flakus A. & Bielczyk U. (2006): New and interesting records of lichens from the Tatry Mountains. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 271–281, Mycotaxon Ltd., Ithaca.
- Follmann G. (2006): Two new roccelloid *Physciaceae* from Pacific South America and a synopsis of the genus *Santessonia*. – *Journal of the Hattori Botanical Laboratory* 100: 651–670.
- Fрати L. & Brunialti G. (2006): Long-term biomonitoring with lichens: comparing data from different sampling procedures. – *Environmental Monitoring and Assessment* 119: 391–404.
- Fрати L., Brunialti G. & Loppi S. (2005): Problems related to lichen transplants to monitor trace element deposition in repeated surveys: A case study from central Italy. – *Journal of Atmospheric Chemistry* 52: 221–230.
- Fremstad E., Paal J. & Möls T. (2005): Impacts of increased nitrogen supply on Norwegian lichen-rich alpine communities: a 10-year experiment. – *Journal of Ecology* 93: 471–481.
- Friedel A., Oheimb G. v., Dengler J. & Härdtle W. (2006): Species diversity and species composition of epiphytic bryophytes and lichens - a comparison of managed and unmanaged beech forests in NE Germany. – *Feddes Repertorium* 117: 172–185.
- Frisch A. & Kalb K. (2006): The lichen genus *Topeliopsis*, additions and corrections. – *Lichenologist* 38: 37–45.
- Frisch A., Kalb K. & Grube M. (2006): Contributions towards a new systematics of the lichen family *Thelotre mataceae*. – *Bibliotheca Lichenologica* 92: 1–556.
- Fröberg L., Baur A. & Baur B. (2006): Field study on the regenerative capacity of three calcicolous lichen species damaged by snail grazing. – *Lichenologist* 38: 491–493.
- Fryday A. M. (2006): New and interesting North American lichen records from the alpine and sub-alpine zones of Mt. Katahdin, Maine. – *Bryologist* 109: 570–578.
- Gaio-Oliveira G., Moen J., Danell Ö. & Palmqvist K. (2006): Effect of simulated reindeer grazing on the re-growth capacity of mat-forming lichens. – *Basic and Applied Ecology* 7: 109–121.
- Gauslaa Y. (2006): Trade-off between reproduction and growth in the foliose old forest lichen *Lobaria pulmonaria*. – *Basic and Applied Ecology* 7: 455–460.
- Gauslaa Y., Holien H., Ohlson M. & Solhøy T. (2006): Does snail grazing affect growth of the old forest lichen *Lobaria pulmonaria*? – *Lichenologist* 38: 587–593.
- Gauslaa Y., Lie M., Solgaug K. A. & Ohlson M. (2006): Growth and ecophysiological acclimation of the foliose lichen *Lobaria pulmonaria* in forests with contrasting light climates. – *Oecologia* 147: 406–416.
- Geebelen W., Bijmens O., Claeys N., Adriaenssens E., Makkonen S., Ruttens A., Carleer R., Yperman J., Clijsters H., Holopainen T. & Vangronsveld J. (2006): Transplanted lichens as biomonitors for atmospheric fluoride pollution near two fluoride point sources in Flanders (Belgium). – *Belgian Journal of Botany* 138: 141–151.
- Giordani P. (2006): Variables influencing the distribution of epiphytic lichens in heterogeneous areas: A case study for Liguria, NW Italy. – *Journal of Vegetation Science* 17: 195–206.
- Gombert S., Asta J. & Seaward M. R. D. (2006): Lichens and tobacco plants as complementary biomonitors of air pollution in the Grenoble area (Isère, southeast France). – *Ecological Indicators* 6: 429–443.
- Gorbushina A. A., Beck A. & Schulte A. (2005): Microcolonial rock inhabiting fungi and lichen photobionts: evidence for mutualistic interactions. – *Mycological Research* 109: 1288–1296.
- Gregersen I. K., Hegseth M. N., Hestmark G., Kongsbak R. H. & Moe T. F. (2006): The relationship between thallus mass, surface area and apothecium production in *Umbilicaria rigida*. – *Nova Hedwigia* 82: 115–121.
- Groner U. (2006): The genus *Chaenothecopsis* (*Mycocaliciaceae*) in Switzerland, and a key to the European species. – *Lichenologist* 38: 395–406.
- Grube M. & Kantvilas G. (2006): *Siphula* represents a remarkable case of morphological convergence in sterile lichens. – *Lichenologist* 38: 241–249.
- Gulluce M., Aslan A., Sokmen M., Sahin F., Adiguzel A., Agar G. & Sokmen A. (2006): Screening the antioxidant and antimicrobial properties of the lichens *Parmelia saxatilis*, *Platismatia glauca*, *Ramalina pollinaria*, *Ramalina polymorpha* and *Umbilicaria nylanderiana*. – *Phytomedicine* 13: 515–521.
- Guschina I. A. & Harwood J. L. (2006): Lead and copper effects on lipid metabolism in cultured lichen photobionts with different phosphorus status. – *Phytochemistry* 67: 1731–1739.

- Guzow-Krzemińska B. (2006): Photobiont flexibility in the lichen *Protoparmeliopsis muralis* as revealed by ITS rDNA analyses. – *Lichenologist* 38: 469–476.
- Hafellner J. (2006): Lecideoid lecanoralean ascomycetes invading *Rhizocarpon* subgen. *Rhizocarpon* taxa, with special emphasis on cryptothalline species. – *Fritschiana* 52: 31–48.
- Hafellner J. (2006): *Protoblastenia szaferi* (lichenized *Ascomycotina*) – new to the Alps. – *Herzogia* 19: 23–33.
- Hafellner J. & John V. (2006): Über Funde lichenicoler nicht-lichenisierter Pilze in der Türkei, mit einer Synopsis der bisher im Land nachgewiesenen Taxa. – *Herzogia* 19: 155–176.
- Hafellner J. & Muggia L. (2006): Über Vorkommen von *Caloplaca erodens* in der Steiermark (Österreich). – *Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark* 135: 33–49.
- Hafellner J., Petutschnig W., Taurer-Zeiner C. & Mayrhofer H. (2005): Über einige bemerkenswerte Flechtenfunde in Kärnten, hauptsächlich in den Gurktaler Alpen. – *Carinthia II* 195/115: 423–439.
- Hansen E. S. (2006): New or interesting Greenland lichens X. – *Cryptogamie, Mycologie* 27: 271–277.
- Hansen E. S. (2006): Notes on some new and interesting Greenland lichens IX. – *Graphis Scripta* 18: 1–5.
- Hansen E. S., Dawes P. R. & Thomassen B. (2006): Epilithic lichen communities in high Arctic Greenland: Physical, environmental, and geological aspects of their ecology in Inglefield Land (78°–79°N). – *Arctic, Antarctic, and Alpine Research* 38: 72–81.
- Harris R. C. (2006): A preliminary glance at *Maronea* (*Fuscideaceae*) in North America. – *Opuscula Philolichenum* 3: 65–68.
- Harris R. C. & Knudsen K. (2006): The genus *Myriospora*. – *Opuscula Philolichenum* 3: 1–4.
- Hauck M., Hofmann E. & Schmall M. (2006): Site factors determining epiphytic lichen distribution in a dieback-affected spruce–fir forest on Whiteface Mountain, New York: microclimate. – *Annales Botanici Fennici* 43: 1–12.
- Hauck M. & Javkhlán S. (2006): Additions to the lichen flora of Mongolia: records from Khentey and Khangay. – *Willdenowia* 36: 895–912.
- Hauck M., Paul A. & Spribille T. (2006): Uptake and toxicity of manganese in epiphytic cyanolichens. – *Environmental and Experimental Botany* 56: 216–224.
- Hawksworth D. L. (2006): Misunderstanding the status of Ciferri & Tomaselli's generic names necessitates *Peterjamesia* gen. nov. for *Sclerophyton circumscriptum* and an additional species. – *Lichenologist* 38: 187–190.
- Hawksworth D. L., Earland-Bennett P. M. & Coppins B. J. (2006): *Lichenophoma haematommatis*, a previously overlooked European sorediate species of *Cliostomum* (*Lecanorales*, *Ramalinaceae*). – *Herzogia* 19: 5–10.
- Hébert F., Boucher J.-F., Bernier P. Y. & Lord D. (2006): Growth response and water relations of 3-year-old planted black spruce and jack pine seedlings in site prepared lichen woodlands. – *Forest Ecology and Management* 223: 226–236.
- Hedenås H., Lundin K. & Ericson L. (2006): Interaction between a lichen and a fungal parasite in a successional community: implications for conservation. – *Journal of Vegetation Science* 17: 207–216.
- Hertel H. (2006): World distribution of species of *Lecidea* (*Lecanorales*) occurring in Central Europe. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 19–73, Mycotaxon Ltd., Ithaca.
- Hilmo O., Holien H. & Hytteborn H. (2005): Logging strategy influences colonization of common chlorolichens on branches of *Picea abies*. – *Ecological Applications* 15: 983–996
- Hobohm C., Litterski B., Boch S. & Schiefelbein U. (2005): Flechten in der Góhrde. – *Jahresberichte des Naturwissenschaftlichen Vereines Lüneburg* 43: 165–174.
- Högnabba F. (2006): Molecular phylogeny of the genus *Stereocaulon* (*Stereocaulaceae*, lichenized ascomycetes). – *Mycological Research* 110: 1080–1092.
- Holien H. (2005): Additions to the Norwegian flora of lichens and lichenicolous fungi III. – *Graphis Scripta* 17: 62–64.
- Holien H. & Tønsberg T. (2006): *Norsk lavflora*. – Tapir Akademisk Forlag, Trondheim. [224 pp.]
- Holt E. A., McCune B. & Neitlich P. (2006): Defining a successional metric for lichen communities in the Arctic Tundra. – *Arctic, Antarctic, and Alpine Research* 38: 373–377.
- Holtan-Hartwig J. (2005): *Peltigera latiloba*, a new lichen species from Norway and USA (Alaska). – *Graphis Scripta* 17: 34.
- Honegger R. (2006): Water relations in lichens. – In: Gadd G. M., Watkinson S. C. & Dyer P. (eds.), *Fungi in the environment*, p. 185–200, Cambridge University Press.

- Huneck S. & Schmidt J. (2006): Phenolische Verbindungen einiger Flechten aus der Familie *Physciaceae*. – *Herzogia* 19: 199–203.
- Incerti G. & Nimis P. L. (2006): Biogeographical outline of epiphytic lichens in a Mediterranean area: Calabria (S Italy). – *Lichenologist* 38: 355–371.
- Jackson H. B., St. Clair L. L. & Eggett D. L. (2006): Size is not a reliable measure of sexual fecundity in two species of lichenized fungi. – *Bryologist* 109: 157–165.
- Jastrzębska B. (2005): Porosty cmentarza w Ugoszczy oraz jego okolicy (Nadbużański Park Krajobrazowy). – *Fragmenta Floristica et Geobotanica Polonica* 12: 194–197.
- Johansson P. & Reich P. B. (2005): Population size and fire intensity determine post-fire abundance in grassland lichens. – *Applied Vegetation Science* 8: 193–198.
- Johansson P., Wetmore C. M., Carlson D. J., Reich P. B. & Thor G. (2006): Habitat preference, growth form, vegetative dispersal and population size of lichens along a wildfire severity gradient. – *Bryologist* 109: 527–540.
- John V. (2006): Die Schlackenhalde bei Hostenbach im mittleren Saartal als Sekundärbiotop für Flechten. – *Herzogia* 19: 49–61.
- John V. (2006): Flechten auf den Halde und im Industriegelände der ehemaligen Grube Reden. – *Abh. Delattinia* 30: 191–195.
- John V. & Türk A. (2006): Species/area curves for lichens on gypsum in Turkey. – *Mycologia Balcanica* 3: 55–60.
- Jørgensen P. M. (2005): *Placynthium garovaglioii* not present in Scandinavia. – *Graphis Scripta* 17: 3–7.
- Jørgensen P. M. (2006): *Leptochidium crenulatum*, a misunderstood arctic-alpine lichen. – *Graphis Scripta* 18: 19–22.
- Jørgensen P. M. & Johnsen J. I. (2006): *Fuscopannaria atlantica* new to Scandinavia. – *Graphis Scripta* 18: 16–18.
- Jovan S. & McCune B. (2006): Using epiphytic macrolichen communities for biomonitoring ammonia in forests of the greater Sierra Nevada, California. – *Water, Air and Soil Pollution* 170: 69–93.
- Jüriado I., Suija A. & Liira J. (2006): Biogeographical determinants of lichen species diversity on islets in the West-Estonian Archipelago. – *Journal of Vegetation Science* 17: 125–134.
- Kaschik M. (2006): Taxonomic studies on saxicolous species of the genus *Rinodina* (lichenized Ascomycetes, *Physciaceae*) in the Southern Hemisphere with emphasis in Australia and New Zealand. – *Bibliotheca Lichenologica* 93: 1–162.
- Ketner-Oostra R., van der Peijl M. J. & Sýkora K. V. (2006): Restoration of lichen diversity in grass-dominated vegetation of coastal dunes after wildfire. – *Journal of Vegetation Science* 17: 147–156.
- Kirschbaum U., Windisch U., Vorbeck A. & Hanewald K. (2006): Mapping lichen diversity in Wetzlar and Giessen as indicator of air quality. (Comparison between the surveys of 1970, 1985, 1995 and 2005). – *Gefahrstoffe - Reinhaltung der Luft* 66: 272–280.
- Kiszka J. (2005): Lichens of the Biała Woda nature reserve in the Małe Pieniny Mts (Western Carpathians). – *Polish Botanical Studies* 19: 177–188.
- Kiszka J. & Kościelniak R. (2006): Localities of the high-mountain species *Leucocarpia biatorella* in the Silesian Upland (southern Poland). – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 325–329, Mycotaxon Ltd., Ithaca.
- Kiurski J. S., Ranogajec J. G., Ujhelji A. L., Radeka M. M. & Bokorov M. T. (2005): Evaluation of the effect of lichens on ceramic roofing tiles by scanning electron microscopy and energy-dispersive spectroscopy analyses. – *Scanning* 27: 113–119.
- Kłos A., Rajfur M., Waclawek M. & Waclawek W. (2006): Determination of the atmospheric precipitation pH value on the basis of the analysis of lichen cationoactive layer constitution. – *Electrochimica Acta* 51: 5053–5061.
- Knowles R. D., Pastor J. & Biesboer D. D. (2006): Increased soil nitrogen associated with dinitrogen-fixing, terricolous lichens of the genus *Peltigera* in northern Minnesota. – *Oikos* 114: 37–48.
- Kochaniewicz G. & Wirth V. (2006): Untersuchung zur Diversität von Epiphyten unter Lokalbedingungen des Südschwarzwaldes. – *Herzogia* 19: 215–227.
- Kondratyuk S., Khodosovtsev A. & Kärnefelt I. (2006): *Llimoniella caloplacae* sp. nova (*Leothiales*), a new lichenicolous fungus on *Caloplaca borysthenica* sp. nova (*Lecanorales*, *Ascomycotina*). – *Mycologia Balcanica* 3: 95–98.
- Kościelniak R. (2004): Porosty (*Lichenes*) Bieszczadów Niskich. – *Fragmenta Floristica et Geobotanica Polonica*, Suppl. 5: 3–164.

- Kościelniak R. & Kiszka J. (2005): A supplement to the lichen checklist of the Eastern Carpathians. – *Roczniki Bieszczadzkie* 13: 235–244.
- Kossowska M. (2006): Checklist of lichens and allied fungi of the Polish Karkonosze Mts. – W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków. [131 pp.]
- Krzewicka B. & Galas J. (2006): Ecological notes on *Verrucaria aquatilis* and *V. hydrela* in the Polish Tatry Mountains. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), Central European lichens – diversity and threat, p. 193–203, Mycotaxon Ltd., Ithaca.
- Kukwa M. (2005): *Lecanora thysanophora* (Lecanoraceae, zlichenizowane Ascomycota) w Polsce. – *Fragmenta Floristica et Geobotanica Polonica* 12: 385–391.
- Kukwa M. (2005): Notes on the chemistry and distribution of *Lecanora rouxii* (lichenized Ascomycota) in Poland. – *Botanica Lithuanica* 11: 161–166.
- Kukwa M. (2005): The lichen species *Cladonia incrassata* (Lecanorales, Ascomycota lichenisati) in Poland, and notes on *C. anitae*. – *Polish Botanical Journal* 50: 69–73.
- Kukwa M. (2006): Notes on taxonomy and distribution of the lichen species *Lepraria ecorticata* comb. nov. – *Mycotaxon* 97: 63–66.
- Kukwa M. (2006): The lichen genus *Lepraria* in Poland. – *Lichenologist* 38: 293–305.
- Kukwa M. & Czarnota P. (2006): New or interesting records of lichenicolous fungi from Poland IV. – *Herzogia* 19: 111–123.
- Kukwa M. & Śliwa L. (2005): The genus *Lepraria* (lichenized Ascomycota) in the Beskid Sądecki Mts (West Carpathians, s Poland). – *Polish Botanical Journal* 50: 163–184.
- Kurokawa S. & Kashiwadani H. (2006): Checklist of Japanese lichens and allied fungi. – *National Science Museum Tokyo* 33: 1–157.
- LaGreca S. (2006): Notes on the chemistry of *Maronea constans* and *Maronea polyphaea* (Fuscideaceae). – *Lichenologist* 38: 595–598.
- LaGreca S. & Stutzman B. W. (2006): Distribution and ecology of *Lecanora conizaeoides* (Lecanoraceae) in eastern Massachusetts. – *Bryologist* 109: 335–347.
- Lakatos M., Rascher U. & Büdel B. (2006): Functional characteristics of corticolous lichens in the understory of a tropical lowland rain forest. – *New Phytologist* 172: 679–695.
- Lalley J. S. & Viles H. A. (2006): Do vehicle track disturbances affect the productivity of soil-growing lichens in a fog desert? – *Functional Ecology* 20: 548–556.
- Lalley J. S., Viles H. A., Copeman N. & Cowley C. (2006): The influence of multi-scale environmental variables on the distribution of terricolous lichens in a fog desert. – *Journal of Vegetation Science* 17: 831–838.
- Lalley J. S., Viles H. A., Henschel J. R. & Lalley V. (2006): Lichen-dominated soil crusts as arthropod habitat in warm deserts. – *Journal of Arid Environments* 67: 579–593.
- Lange O. L. & Green T. G. A. (2006): Nocturnal respiration of lichens in their natural habitat is not affected by preceding diurnal net photosynthesis. – *Oecologia* 148: 396–404.
- Lange O. L., Green T. G. A., Melzer B., Meyer A. & Zellner H. (2006): Water relations and CO₂ exchange of the terrestrial lichen *Teloschistes capensis* in the Namib fog desert: Measurements during two seasons in the field and under controlled conditions. – *Flora* 201: 268–280.
- Laufer Z., Beckett R. P., Minibayeva F. V., Lüthje S. & Böttger M. (2006): Occurrence of laccases in lichenized Ascomycetes of the *Peltigerineae*. – *Mycological Research* 110: 846–853.
- Liba C. M., Ferrara F. I. S., Manfio G. P., Fantinatti-Garboggini F., Albuquerque R. C., Pavan C., Ramos P. L., Moreira-Filho C. A. & Barbosa H. R. (2006): Nitrogen-fixing chemo-organotrophic bacteria isolated from cyanobacteria-deprived lichens and their ability to solubilize phosphate and to release amino acids and phytohormones. – *Journal of Applied Microbiology* 101: 1076–1086.
- Lindblom L. (2006): *Xanthomendoza galericulata*, a new sorediate lichen species, with notes on similar species in North America. – *Bryologist* 109: 1–8.
- Lindblom L. & Ekman S. (2006): Genetic variation and population differentiation in the lichen-forming ascomycete *Xanthoria parietina* on the island Storfosna, central Norway. – *Molecular Ecology* 15: 1545–1559.
- Lipnicki L. (2006): Protected lichens in The Bory Tucholskie Forest (N Poland) and threats to them. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), Central European lichens – diversity and threat, p. 331–336, Mycotaxon Ltd., Ithaca.
- Litterski B. (2006): Foliose and fruticose lichens of Kyrgyzstan. – *Herzogia* 19: 177–190.

- Llop E. & Gómez-Bolea A. (2006): Foliicolous lichens and associated lichenicolous fungi in the north-eastern Iberian Peninsula: the effect of environmental factors on distribution. – *Lichenologist* 38: 55–65.
- Lõhmus P. (2005): Forest lichens and their substrata in Estonia. – Tartu University, Tartu. [156 pp.]
- Loppi S. & Frati L. (2006): Lichen diversity and lichen transplants as monitors of air pollution in a rural area of central Italy. – *Environmental Monitoring and Assessment* 114: 361–375.
- Loppi S., Paoli L. & Gaggi C. (2006): Diversity of epiphytic lichens and Hg contents of *Xanthoria parietina* thalli as monitors of geothermal air pollution in the Mt. Amiata area (central Italy). – *Journal of Atmospheric Chemistry* 53: 93–105.
- Loso M. G. & Doak D. F. (2006): The biology behind lichenometric dating curves. – *Oecologia* 147: 223–229.
- Lücking R., Aptroot A., Umaña L., Chaves J. L., Sipman H. J. M. & Nelsen M. P. (2006): A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Gyalideopsis* and its segregates (*Ostropales: Gomphillaceae*), with a world-wide key and name status checklist. – *Lichenologist* 38: 131–160.
- Lumbsch H. T., Schmitt I., Barker D. & Pagel M. (2006): Evolution of micromorphological and chemical characters in the lichen-forming fungal family *Pertusariaceae*. – *Biological Journal of the Linnean Society* 89: 615–626.
- Machado A., Šlejkovec Z., Van Elteren J. T., Freitas M. C. & Baptista M. S. (2006): Arsenic speciation in transplanted lichens and tree bark in the framework of a biomonitoring scenario. – *Journal of Atmospheric Chemistry* 53: 237–249.
- Marsh J. E. & Timoney K. P. (2005): How long must northern saxicolous lichens be immersed to form a waterbody trimline? – *Wetlands* 25: 495–499.
- Martínez I., Carreño F., Escudero A. & Rubio A. (2006): Are threatened lichen species well-protected in Spain? Effectiveness of a protected areas network. – *Biological Conservation* 133: 500–511.
- Mattsson J.-E., Lättman H. & Milberg P. (2006): Rapid changes in the epiphytic macrolichen flora on sites in southern Sweden. – *Lichenologist* 38: 323–329.
- Mayer W., Pfefferkorn-Dellali V. & Türk R. (2006): Immissionsökologische Flechtenkartierung am Zöbelboden im Reichraminger Hintergebirge, Oberösterreich. – *Beiträge zur Naturkunde Oberösterreichs* 16: 443–446.
- Mayrhofer H., Kalb K. & Harutyunyan S. (2006): Belege zur „Flora von Istrien“ aus anderen Herbarien. Flechten. – In: Starmühler W. (ed.), Vorarbeiten zu einer „Flora von Istrien“ Teil IX. – *Carinthia II* 196/116: 573–581.
- Mayrhofer H., Mešl R., Batič F. & Bilovitz P. O. (2006): Remarkable records of lichenized and lichenicolous fungi from Slovenia. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 165–178, Mycotaxon Ltd., Ithaca.
- Mayrhofer H., Mešl R. & Bilovitz P. O. (2005): Belege zur „Flora von Istrien“ aus anderen Herbarien. Flechten. – In: Starmühler W. (ed.), Vorarbeiten zu einer „Flora von Istrien“ Teil VIII. – *Carinthia II* 195/115: 640–646.
- McCune B., Ahti T. & Duncan C. M. (2006): *Hypogymnia incurvodes* (*Parmeliaceae*), a little-known Russian species discovered in eastern North America. – *Bryologist* 109: 80–84.
- McEvoy M., Nybakken L., Solhaug K. A. & Gauslaa Y. (2006): UV triggers the synthesis of the widely distributed secondary lichen compound usnic acid. – *Mycological Progress* 5: 221–229.
- Monnet F., Bordas F., Deluchat V. & Baudu M. (2006): Toxicity of copper excess on the lichen *Dermatocarpon luridum*: Antioxidant enzyme activities. – *Chemosphere* 65: 1806–1813.
- Motiejūnaitė J. & Fałtynowicz W. (2005): Effect of land-use on lichen diversity in the transboundary region of Lithuania and northeastern Poland. – *Ekologija* 2005/3: 34–43.
- Motiejūnaitė J. & Jucevičienė N. (2005): Epidemiology of the fungus *Athelia arachnoidea* in epiphytic communities of broadleaved forests under strong anthropogenic impact. – *Ekologija* 2005/4: 28–34.
- Motiejūnaitė J., Suija A., Lõhmus P., Kuznetsova E., Tõrra T., Prigodina-Lukošienė I. & Piterāns A. (2006): New or noteworthy lichens, lichenicolous and allied fungi found during the 16th Symposium of Mycologists and Lichenologists in Latvia. – *Botanica Lithuanica* 12: 113–119.
- Mrak T., Šlejkovec Z. & Jeran Z. (2006): Extraction of arsenic compounds from lichens. – *Talanta* 69: 251–258.
- Myllys L., Halonen P. & Velmala S. (2006): Notes on some rare species of *Bryoria* from Finland. – *Graphis Scripta* 18: 23–26.
- Nascimbene J. (2006): Lichenological studies in N-Italy : new records for Lombardy. – *Cryptogamie, Mycologie* 27: 79–82.

- Nascimbene J., Caniglia G. & Dalle Vedove M. (2006): Lichen diversity and ecology in five EU habitats of interest of the Sexten Dolomiten Natural Park (S Tyrol - NE Italy). – *Cryptogamie, Mycologie* 27: 185–193.
- Nascimbene J., Caniglia G., Nicli M. & Dalle Vedove M. (2006): Populations of *Lobaria pulmonaria* (L.) Hoffm. in the Cansiglio Regional Forest (Veneto, Pre-Alps, north-east Italy): Distribution, diversity and conservation issues. – *Plant Biosystems* 140: 34–42.
- Nascimbene J., Isocrono D., Marini L., Caniglia G. & Piervittori R. (2006): Epiphytic lichen vegetation on *Larix* in the Italian Alps. – *Plant Biosystems* 140: 132–137.
- Nascimbene J., Martellos S. & Nimis P. L. (2006): Epiphytic lichens of tree-line forests in the Central-Eastern Italian Alps and their importance for conservation. – *Lichenologist* 38: 373–382.
- Nelsen M. P., Lücking R., Chaves J. L., Sipman H. J. M., Umaña L. & Navarro E. (2006): A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Haematomma* (*Lecanorales, Lecanoraceae*). – *Lichenologist* 38: 251–262.
- Neuwirth G., Breuss O. & Hammerl R. (2006): Beitrag zur Flechtenflora des Nationalparks Bayerischer Wald. Die Buchberger Leite. – *Österreichische Zeitschrift für Pilzkunde* 15: 105–120.
- Nimis P. L. & Skert N. (2006): Lichen chemistry and selective grazing by the coleopteran *Lasioderma serricornis*. – *Environmental and Experimental Botany* 55: 175–182.
- Nordin A. (2005): *Rhizocarpon umense*, a distinct but originally misplaced species. – *Graphis Scripta* 17: 37–38.
- Nordin A. & Tibell L. (2005): Additional species in *Tetramelas*. – *Lichenologist* 37: 491–498.
- Nybakken L. & Julkunen-Tiitto R. (2006): UV-B induces usnic acid in reindeer lichens. – *Lichenologist* 38: 477–485.
- Ohmura Y., Kawachi M., Kasai F. & Watanabe M. (2006): Genetic combinations of symbionts in a vegetatively reproducing lichen, *Parmotrema tinctorum*, based on ITS rDNA sequences. – *Bryologist* 109: 43–59.
- Oksanen I. (2006): Ecological and biotechnological aspects of lichens. – *Applied Microbiology and Biotechnology* 73: 723–734.
- Omarsdóttir S., Ólafsdóttir E. S. & Freysdóttir J. (2006): Immunomodulating effects of lichen-derived polysaccharides on monocyte-derived dendritic cells. – *International Immunopharmacology* 6: 1642–1650.
- Omarsdóttir S., Petersen B. O., Paulsen B. S., Togola A., Duus J. Ø. & Ólafsdóttir E. S. (2006): Structural characterization of novel lichen heteroglycans by NMR spectroscopy and methylation analysis. – *Carbohydrate Research* 341: 2449–2455.
- Opanowicz M., Blaha J. & Grube M. (2006): Detection of paralogous polyketide synthase genes in *Parmeliaceae* by specific primers. – *Lichenologist* 38: 47–54.
- Osyczka P. (2006): The lichen genus *Cladonia* (*Cladoniaceae*, lichenized *Ascomycota*) from Spitsbergen. – *Polish Polar Research* 27: 207–242.
- Osyczka P. & Stolarczyk P. (2005): Porosty z rodzaju *Lepraria* w rejonie Dołów Jasielsko-Sanockich (Karpaty Zachodnie). – *Fragmenta Floristica et Geobotanica Polonica* 12: 371–383.
- Osyczka P. & Węgrzyn M. (2005): Arktyczno-alpejski gatunek porostu *Cladonia amaurocraea* w Tatrach. – *Fragmenta Floristica et Geobotanica Polonica* 12: 157–163.
- Otte V. (2005): Noteworthy lichen records for Bulgaria. – *Abhandlungen und Berichte des Naturkundemuseums Görlitz* 77: 77–86.
- Otte V. (2006): Herbarnachweis von *Vulpicida juniperinus* für die deutsche Flechtenflora. – *Herzogia* 19: 43–48.
- Otte V., van den Boom P. & Rätzel S. (2006): Bemerkenswerte Funde von Flechten und lichenicolen Pilzen aus Brandenburg XI. – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 139: 275–291.
- Øvstedal D. O. & Gremmen N. J. M. (2006): Lichens of sub-Antarctic Heard Island. – *South African Journal of Botany* 72: 353–366.
- Palka K. & Śliwa L. (2006): Lichens of Oran Dosh protected area in the Lake Khubsugul National Park (Mongolia). Part 1: corticolous and lignicolous species. – *Nova Hedwigia* 82: 51–68.
- Palmqvist K. & Dahlman L. (2006): Responses of the green algal foliose lichen *Platismatia glauca* to increased nitrogen supply. – *New Phytologist* 171: 343–356.
- Paul A. & Hauck M. (2006): Effects of manganese on chlorophyll fluorescence in epiphytic cyano- and chlorolichens. – *Flora* 201: 451–460.

- Pérez-Ortega S. & Álvarez-Lafuente A. (2006): Primer catálogo de líquenes y hongos liquenícolas de la Comunidad Autónoma de Cantabria. – *Botanica Complutensis* 30: 5–16.
- Pérez-Ortega S. & Álvarez-Lafuente A. (2006): Primer catálogo de líquenes y hongos liquenícolas de la Comunidad Autónoma de Castilla y León (España). – *Botanica Complutensis* 30: 17–52.
- Perlmutter G. B. (2006): *Flakea papillata* in North America. – *Bryologist* 109: 566–569.
- Piercey-Normore M. D. (2006): Lichens from the Hudson Bay Lowlands: diversity in the southeastern peatlands of Wapusk National Park, Manitoba. – *Canadian Journal of Botany* 84: 1781–1793.
- Piercey-Normore M. D. (2006): The lichen-forming ascomycete *Evernia mesomorpha* associates with multiple genotypes of *Trebouxia jamesii*. – *New Phytologist* 169: 331–344.
- Pirintsos S. A., Matsi T., Vokou D., Gaggi C. & Loppi S. (2006): Vertical distribution patterns of trace elements in an urban environment as reflected by their accumulation in lichen transplants. – *Journal of Atmospheric Chemistry* 54: 121–131.
- Pöykkö H. (2006): Females and larvae of a geometrid moth, *Cleorodes lichenaria*, prefer a lichen host that assures shortest larval period. – *Environmental Entomology* 35: 1669–1676.
- Printzen C. & Otte V. (2005): *Biatora longispora*, new to Europe, and a revised key to European and Macaronesian *Biatora* species. – *Graphis Scripta* 17: 56–61.
- Printzen C., Tønsberg T. & Ekman S. (2005): Jetsetter oder Säulenheilige. Wie entstehen Verbreitungsmuster bei Flechten? – *Natur und Museum* 135: 16–25.
- Pykälä J. (2006): Additions to the lichen flora of Finland. – *Graphis Scripta* 18: 41–48.
- Pykälä J., Heikkinen R. K., Toivonen H. & Jääskeläinen K. (2006): Importance of forest act habitats for epiphytic lichens in Finnish managed forests. – *Forest Ecology and Management* 223: 84–92.
- Randlane T. (2006): Genus *Xylographa* in Estonia. – *Folia Cryptogamica Estonica* 42: 105–107.
- Randlane T. & Saag A. (2006): Cetrarioid lichens in Europe – an identification key for the species. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 75–84, Mycotaxon Ltd., Ithaca.
- Rätzel F., Otte V., de Bruyn W. & Sipman H. J. M. (2006): Bemerkenswerte Flechtenfunde aus Brandenburg (incl. lichenicoler und lichenoider Pilze) X. – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 138: 83–105.
- Ravera S. (2006): Two new species of *Arthopyrenia* from Italy. – *Lichenologist* 38: 21–26.
- Ravera S., Genovesi V. & Massari G. (2006): Phytoclimatic characterization of lichen habitats in central Italy. – *Nova Hedwigia* 82: 143–165.
- Redhead S. A. & Jørgensen P. M. (2006): Proposal to conserve the name *Hygrophorus hudsonianus* against *Endocarpon viride* and *Normandina davidis* (lichenized *Basidiomycota*). – *Taxon* 55: 1036–1037.
- Rico V. J., Calatayud V. & Giralt M. (2006): The lichen genus *Dimelaena* (*Physciaceae*, *Lecanorales*) in the Iberian Peninsula. – *Cryptogamie, Mycologie* 27: 149–166.
- Rivas Plata E., Lücking R., Aptroot A., Sipman H. J. M., Chaves J. L., Umaña L. & Lizano D. (2006): A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Coenogonium* (*Ostropales*: *Coenogoniaceae*), with a world-wide key and checklist and a phenotype-based cladistic analysis. – *Fungal Diversity* 23: 255–321.
- Roszbach M. & Lambrecht S. (2006): Lichens as biomonitors: Global, regional and local aspects. – *Croatica Chemica Acta* 79: 119–124.
- Roux C., Bricaud O. & Coste C. (2005): Lichens et champignons lichénicoles du parc national des Cévennes (France) 1 - Le mont Lozère. – *Bulletin de la Société Linnéenne de Provence* 56: 201–224.
- Roux C. & Coste C. (2005): Lichens et champignons lichénicoles du parc national des Cévennes (France) 2 - Le causse Méjean. – *Bulletin de la Société Linnéenne de Provence* 56: 225–241.
- Roux C., Coste C., Bricaud O. & Masson D. (2006): Catalogue des lichens et des champignons lichénicoles de la région Languedoc-Roussillon (France méridionale). – *Bulletin de la Société Linnéenne de Provence* 57: 85–200.
- Roux C., Coste C., Masson D. & Bauvet C. (2006): Lichens et champignons lichénicoles du parc national des Cévennes (France) 3 - Les basses Cévennes. – *Bulletin de la Société Linnéenne de Provence* 57: 59–84.
- Rusu A.-M., Jones G. C., Chimonides P. D. J. & Purvis O. W. (2006): Biomonitoring using the lichen *Hypogymnia physodes* and bark samples near Zlatna, Romania immediately following closure of a copper ore-processing plant. – *Environmental Pollution* 143: 81–88.
- Sanders W. B. (2006): A feeling for the superorganism: expression of plant form in the lichen thallus. – *Botanical Journal of the Linnean Society* 150: 89–99.

- Sanderson N. (2006): *Cliostomum flavidulum* Haf. & Kalb, yet another yellow soresiate crust. – British Lichen Society Bulletin 98: 29–35
- Sanderson N. (2006): *Pyrenula hibernica* at Ceunant Llennyrch, Meirionnydd. – British Lichen Society Bulletin 98: 23–27.
- Sanità di Toppi L., Musetti R., Vattuone Z., Pawlik-Skowrońska B., Fossati F., Bertoli L., Badiani M. & Favali M. A. (2005): Cadmium distribution and effects on ultrastructure and chlorophyll status in photobionts and mycobionts of *Xanthoria parietina*. – Microscopy Research and Technique 66: 229–238.
- Savić S. & Tibell L. (2006): Checklist of the lichens of Serbia. – Mycologia Balcanica 3: 187–215.
- Savić S., Tibell L. & Andreev M. (2006): New and interesting lichenized and lichenicolous fungi from Serbia. – Mycologia Balcanica 3: 99–106.
- Schiefelbein U. (2006): Ökologische und naturschutzfachliche Aspekte der Flechtenflora des Landkreises Uecker-Randow (Mecklenburg-Vorpommern). – Archiv Naturwissenschaftlicher Dissertationen 16: 1–216.
- Schmitt I., Yamamoto Y. & Lumbsch H. T. (2006): Phylogeny of *Pertusariales* (*Ascomycotina*): resurrection of *Ochrolechiaceae* and new circumscription of *Megasporaceae*. – Journal of the Hattori Botanical Laboratory 100: 753–764.
- Schultz M. (2006): *Cryptothete rhodosticta* new to Scandinavia from Norway. – Graphis Scripta 18: 49–51.
- Seaward M. R. D. & Sipman H. J. M. (2006): An updated checklist of lichenized and lichenicolous fungi for Egypt. – Willdenowia 36: 537–555.
- Sedia E. G. & Ehrenfeld J. G. (2006): Differential effects of lichens and mosses on soil enzyme activity and litter decomposition. – Biology and Fertility of Soils 43: 177–189.
- Seneviratne G. & Indrasena I. K. (2006): Nitrogen fixation in lichens is important for improved rock weathering. – Journal of Biosciences 31: 639–643.
- Sérusiaux E., Berger F., Coppins B. J. & Roux C. (2005): A further new species of *Strigula* from Europe. – Lichenologist 37: 481–483.
- Sérusiaux E., Diederich P., Ertz D., Brand M. & van den Boom P. (2006): New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. X. – Bulletin de la Société des Naturalistes Luxembourgeois 107: 63–74.
- Seymour F. A., Crittenden P. D. & Dyer P. S. (2005): Sex in the extremes: lichen-forming fungi. – Mycologist 19: 51–58.
- Shirtcliffe N. J., Pyatt F. B., Newton M. I. & McHale G. (2006): A lichen protected by a super-hydrophobic and breathable structure. – Journal of Plant Physiology 163: 1193–1197.
- Sipman H. J. M. (2005): Líquenes de los Páramos de Costa Rica. – In: Kappelle M. & Horn S. P. (eds.), Páramos de Costa Rica, p. 242–360, Santo Domingo de Heredia, Costa Rica.
- Śliwa L. (2006): Additions to the lichen flora of the Tatry National Park and its surroundings (Polish Carpathians). – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), Central European lichens – diversity and threat, p. 305–313, Mycotaxon Ltd., Ithaca.
- Śliwa L. (2006): The typification of *Lecanora dispersa* and *L. albescens*. – Mycotaxon 97: 291–297.
- Śliwa L. & Hawksworth D. L. (2006): Proposal to conserve the name *Lichen hagenii* (*Lecanora hagenii*) with a conserved type (lichenized *Ascomycota*). – Taxon 55: 1038–1039.
- Slonov L. K. & Slonov T. Kh. (2006): Radionuclide accumulation in thalli of lichens. – Russian Journal of Ecology 37: 141–142.
- Spribile T., Schultz M., Breuss O. & Bergmeier E. (2006): Notes on the lichens and lichenicolous fungi of western Crete (Greece). – Herzogia 19: 125–148.
- Staiger B., Kalb K. & Grube M. (2006): Phylogeny and phenotypic variation in the lichen family *Graphidaceae* (*Ostropomycetidae*, *Ascomycota*). – Mycological Research 110: 765–772.
- Stenroos S., Högnabba F., Myllys L., Hyvönen J. & Thell A. (2006): High selectivity in symbiotic associations of lichenized ascomycetes and cyanobacteria. – Cladistics 22: 232–238.
- Stocker-Wörgötter E. & Elix J. A. (2006): Morphogenetic strategies and induction of secondary metabolite biosynthesis in cultured lichen-forming *Ascomycota*, as exemplified by *Cladia retipora* (Labill.) Nyl. and *Dactylina arctica* (Richards) Nyl. – Symbiosis 41: 9–20.
- Stofer S., Bergamini A., Aragón G., Carvalho P., Coppins B. J., Davey S., Dietrich M., Farkas E., Kärkkäinen K., Keller C., Lökös L., Lommi S., Máguas C., Mitchell R., Pinho P., Rico V. J., Truscott A.-M., Wolseley P. A., Watt A. & Scheidegger C. (2006): Species richness of lichen functional groups in relation to land use intensity. – Lichenologist 38: 331–353.

- Suija A. (2005): Lichens and lichenicolous fungi in Estonia: Diversity, distribution patterns, taxonomy. – Tartu University, Tartu. [156 pp.]
- Suija A., Jürriado I., Leppik E. & Randlane T. (2006): New Estonian records: New lichens and lichenicolous fungi. – *Folia Cryptogamica Estonica* 42: 103–105.
- Szczepańska K. (2005): Porosty grupy skalnej "Trzy Siostry" v Górach Bialskich (Sudety). – *Fragmenta Floristica et Geobotanica Polonica* 12: 393–397.
- Szczepańska K. & Szczepański A. (2006): Porosty z rodzaju *Bryoria* v Karkonoszach. – *Fragmenta Floristica et Geobotanica Polonica* 13: 191–195.
- Taylor T. N. & Krings M. (2005): Fungi and lichens. – In: Selley R. C., Cocks L. R. M. & Pilmer I. R. (eds.), *Encyclopedia of Geology* Vol. 2., p. 436–443, Amsterdam.
- Thell A., Feuerer T., Elix J. A. & Kärnelfelt I. (2006): A contribution to the phylogeny and taxonomy of *Xanthoparmelia* (Ascomycota, Parmeliaceae). – *Journal of the Hattori Botanical Laboratory* 100: 797–807.
- Thormann M. N. (2006): Lichens as indicators of forest health in Canada. – *Forestry Chronicle* 82: 335–343.
- Thüs H. & Licht W. (2006): Oceanic and hygrophytic lichens from the Gargano-Peninsula (Puglia / South-Eastern-Italy). – *Herzogia* 19: 149–153.
- Tindal E. & Tønsberg T. (2006): *Psoroma paleaceum* comb. nov. the only hairy *Psoroma* in northern Europe. – *Graphis Scripta* 18: 54–57.
- Tønsberg T. (2006): *Pertusaria flavocorallina* new to Norway. – *Graphis Scripta* 18: 62–63.
- Tønsberg T. & Zhurbenko M. (2006): *Lepraria gelida*, a new species from the Arctic. – *Graphis Scripta* 18: 64.
- Tozer W. C., Hackell D., Miers D. B. & Silvester W. B. (2005): Extreme isotopic depletion of nitrogen in New Zealand lithophytes and epiphytes; the result of diffusive uptake of atmospheric ammonia? – *Oecologia* 144: 628–635.
- Tretiach M. & Muggia L. (2006): *Caloplaca badioreagens*, a new calcicolous, endolithic lichen from Italy. – *Lichenologist* 38: 223–229.
- Tretiach M. & Rinino S. (2006): Critical notes on the calcicolous endolithic lichen *Strigula endolitheae* (Strigulaceae, Ascomycota). – *Nova Hedwigia* 83: 293–310.
- Trinkaus P. (2004): Wiederbesiedelung durch *Xanthoria parietina* als Indiz für eine merkliche Verbesserung der Luftqualität in Klagenfurt. – *Carinthia* II 194/114: 517–522.
- Tucker S. C. & Ryan B. D. (2006): Revised catalog of lichens, lichenicoles, and allied fungi in California. – *Constancea* 84: 1–275.
- Türk R. (2006): Die Flechtenflora in Wien – Veränderungen im Zeitraum 1853 bis 2004. – *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich* 143: 113–118.
- Türk R. & Obermayer W. (2006): The distribution of the species of the genera *Physcia* and *Physconia* (Physciaceae) in Austria. – In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), *Central European lichens – diversity and threat*, p. 119–152, Mycotaxon Ltd., Ithaca.
- Uchida M., Nakatsubo T., Kanda H. & Koizumi H. (2006): Estimation of the annual primary production of the lichen *Cetrariella delisei* in a glacier foreland in the High Arctic, Ny-Ålesund, Svalbard. – *Polar Research* 25: 39–49.
- Urbanavichus G. P. & Melekhin A. B. (2006): Interesnye nachodky lishainikov *Maronella laricina* i *Micarea lutulata* v Permskom krae. – *Botanicheskij Zhurnal* 91: 963–966.
- van den Boom P. P. G. (2006): Contribution to the flora of Portugal: lichens and lichenicolous fungi 5. – *Österreichische Zeitschrift für Pilzkunde* 15: 11–19.
- van den Boom P. P. G. & Etayo J. (2006): New records of lichens and lichenicolous fungi from Fuerteventura (Canary Islands), with descriptions of some new species. – *Cryptogamie, Mycologie* 27: 341–374.
- van den Boom P. P. G. & Rico V. J. (2006): *Caloplaca squamuloisidiata*, a new lichen species from Portugal and Spain. – *Lichenologist* 38: 529–535.
- van den Boom P. & van den Boom B. (2006): An inventory of lichens and lichenicolous fungi in northern Belgium (area of Stramprooierbroek). – *Bulletin de la Société des Naturalistes Luxembourgeois* 106: 39–52.
- Wagner H. H., Werth S., Kalwij J. M., Bolli J. & Scheidegger C. (2006): Modelling forest recolonization by an epiphytic lichen using a landscape genetic approach. – *Landscape Ecology* 21: 849–865.

- Walker T. R., Crittenden P. D., Young S. D. & Prystina T. N. (2006): An assessment of pollution impacts due to the oil and gas industries in the Pechora basin, north-eastern European Russia. – *Ecological Indicators* 6: 369–387.
- Walker T. R. & Pystina T. N. (2006): The use of lichens to monitor terrestrial pollution and ecological impacts caused by oil and gas industries in the Pechora Basin, NW Russia. – *Herzogia* 19: 229–238.
- Washburn S. J. & Culley T. M. (2006): Epiphytic macrolichens of the greater Cincinnati metropolitan area—part II: distribution, diversity and urban ecology. – *Bryologist* 109: 516–526.
- Wedin M., Döring H. & Gilenstam G. (2006): *Stictis* s. lat. (*Ostropales*, *Ascomycota*) in northern Scandinavia, with a key and notes on morphological variation in relation to lifestyle. – *Mycological Research* 110: 773–789.
- Weiss M., Hobbie S. E. & Gettel G. M. (2005): Contrasting responses of nitrogen-fixation in Arctic lichens to experimental and ambient nitrogen and phosphorus availability. – *Arctic, Antarctic and Alpine Research* 37: 396–401.
- Weissman L., Fraiberg M., Shine L., Garty J. & Hochman A. (2006): Responses of antioxidants in the lichen *Ramalina lacera* may serve as an early-warning bioindicator system for the detection of air pollution stress. – *FEMS Microbiology Ecology* 58: 41–53.
- Welch A. R., Gillman M. P. & John E. A. (2006): Effect of nutrient application on growth rate and competitive ability of three foliose lichen species. – *Lichenologist* 38: 177–186.
- Werth S., Wagner H. H., Gugerli F., Holderegger R., Csencsics D., Kalwij J. M. & Scheidegger C. (2006): Quantifying dispersal and establishment limitation in a population of an epiphytic lichen. – *Ecology* 87: 2037–2046.
- Werth S., Wagner H. H., Holderegger R., Kalwij J. M. & Scheidegger C. (2006): Effect of disturbances on the genetic diversity of an old-forest associated lichen. – *Molecular Ecology* 15: 911–921.
- Westberg M. (2006): *Squamarina scopulorum* new to Sweden. – *Graphis Scripta* 18: 52–53.
- Wieczorek A. (2005): Biota porostów Szczecińskiego Parku Krajobrazowego. – *Fragmenta Floristica et Geobotanica Polonica* 12: 143–156.
- Wilk K. & Flakus A. (2006): Four species of *Caloplaca* (*Teloschistaceae*, lichenized *Ascomycota*) new to Poland. – *Mycotaxon* 96: 61–71.
- Will-Wolf S., Geiser L. H., Neitlich P. & Reis A. H. (2006): Linking variation in lichen community composition to environmental variables at regional and subregional geographic scales – how consistent is the story? – *Journal of Vegetation Science* 17: 171–184.
- Wirtz N., Printzen C., Sancho L. G. & Lumbsch H. T. (2006): The phylogeny and classification of *Neuropogon* and *Usnea* (*Parmeliaceae*, *Ascomycota*) revisited. – *Taxon* 55: 367–376.
- Wolseley P. A., James P. W., Theobald M. R. & Sutton M. A. (2006): Detecting changes in epiphytic lichen communities at sites affected by atmospheric ammonia from agricultural sources. – *Lichenologist* 38: 161–176.
- Wolseley P. A., Stofer S., Mitchell R., Truscott A.-M., Vanbergen A., Chimonides J. & Scheidegger C. (2006): Variation of lichen communities with landuse in Aberdeenshire, UK. – *Lichenologist* 38: 307–322.
- Woods R. (2006): *Pertusaria albescens* var. *corallina* as a temporarily dominant epiphytic lichen in mid Wales. – *British Lichen Society Bulletin* 98: 28–29.
- Yahr R., Vilgalys R. & DePriest P. T. (2006): Geographic variation in algal partners of *Cladonia subtenuis* (*Cladoniaceae*) highlights the dynamic nature of a lichen symbiosis. – *New Phytologist* 171: 847–860.
- Zavarzina A. G. & Zavarzin A. A. (2006): Laccase and tyrosinase activities in lichens. – *Microbiology* 75: 546–556.
- Zhou Q.-M., Guo S.-Y., Huang M.-R. & Wei J.-C. (2006): A study of the genetic variability of *Rhizoplaca chrysoleuca* using DNA sequences and secondary metabolic substances. – *Mycologia* 98: 57–67.
- Zhou Q.-M., Wei J.-C., Ahti T., Stenroos S. & Högnabba F. (2006): The systematic position of *Gymnoderma* and *Cetradonia* based on SSU rDNA sequences. – *Journal of the Hattori Botanical Laboratory* 100: 871–880.