

- Školek J. (2003): *Geo rivali-Caricetum paniculatae* ass. nova from the West Carpathians. – *Thaiszia – Journal of Botany* 13: 31–66.
- Šomšák L., Šimonovič V. & Kollár J. (2004): Phytocenoses of pine forests in the central part of the Záhorská nížina lowland. – *Biologia, Bratislava* 59: 101–113.
- Špryňar P. (2004): Poznámky k překvapivému výskytu podmrvky jižní (*Notholaena maranthae*) a sleziníku hadcového (*Asplenium cuneifolium*) na ultrabazickém pikritu v Českém krasu. – *Zprávy České Botanické Společnosti* 39: 321–338.
- Špryňar P. & Kubát K. (2004): *Eragrostis albensis* a *E. pectinacea*, dva nové cizí druhy trav pro květenu České republiky (Poaceae). – *Zprávy České Botanické Společnosti* 39: 1–24.
- Těšitel J. (2004): *Epipactis atrorubens* a *Lithospermum officinale* v Železných horách. – *Zprávy České Botanické Společnosti* 39: 233–235.
- Uhlířová J. & Bernátová D. (2003): Príspevok k flóre a vegetácii skalných stanovišť Muránskej planiny. – *Zborník Slovenského Národného Múzea, Prír. Vedy*, 49: 55–67.
- Valachovič M. (2004): Spoločenstvá lemov na Borskej nížine – príklad edaficky vyvolanej variability. – *Bulletin Slovenskej Botanickej Spoločnosti* 26: 193–200.
- Višňák R. (2003): Jizerskohorská rašeliniště po 40 letech a jejich botanická charakteristika. – *Sborník Severočeského Muzea, Prír. Vědy*, 23: 11–84.
- Višňák R. (2004): *Sesleria uliginosa* – nový druh pro západní Čechy. – *Zprávy České Botanické Společnosti* 39: 397–404.
- Zákravský P., Hroudová Z. & Rauch O. (2004): Response of *Phragmites australis* to heavy metal loaded habitats. – In: Kovář P. (ed.), *Natural recovery of human-made deposits in landscape (Biotic interactions and ore/ash-slag artificial ecosystems)*, p. 83–97, Academia, Prague.

NOVÁ LICHENOLOGICKÁ LITERATURA XIV.

New lichenological literature, XIV

Zdeněk P a l i c e

Botanický ústav AV ČR, CZ–252 43 Průhonice, e-mail: palice@ibot.cas.cz

- Adamo P., Arienzo M., Pugliese M., Roca V. & Violante P. (2004): Accumulation history of radionuclides in the lichen *Stereocaulon vesuvianum* from Mt. Vesuvius (south Italy). – *Environmental Pollution* 127: 455–461.
- Alstrup V. (2004): New records in distribution of lichens and lichenicolous fungi. – *Graphis Scripta* 16: 46–57.
- Andersen H. L. & Ekman S. (2004): Phylogeny of the *Micareaceae* inferred from nrSSU DNA sequences. – *Lichenologist* 36: 27–35.
- Andreev M. P. (2004): Notes on the lichen genus *Miriquidica* (*Lecanorales*, *Lecanoraceae*) in Russia. – *Bibliotheca Lichenologica* 88: 15–42.
- Antoine M. E. (2004): An ecophysiological approach to quantifying nitrogen fixation by *Lobaria oregana*. – *Bryologist* 107: 82–87.
- Antoine M. E. & McCune B. (2004): Contrasting fundamental and realized ecological niches with epiphytic lichen transplants in an old-growth *Pseudotsuga* forest. – *Bryologist* 107: 163–173.
- Aptroot A. (2004): Lichenen in Veere: 115 soorten in een kilometerhok. – *Buxbaumiella* 66: 50–52.
- Aptroot A. & Berg M. P. (2004): *Collembola* help lichens in competition with algae. – *Lichenologist* 36: 167–169.
- Aptroot A. & Seaward M. R. D. (2003): Freshwater lichens. – In: Tsui C. K. M. & Hyde K. D. (eds.), *Freshwater Mycology, Fungal Diversity Research Series* 10, p. 101–110, Fungal Diversity Press, Hong Kong.
- Aptroot A., van Herk C. M., Sparrius L. B. & Spier J. L. (2004): Standaardlijst van Nederlandse korstmossen en korstmosparasieten. – *Buxbaumiella* 69: 17–55.

- Aptroot A. & Zielmann R. (2003): *Lobaria amplissima* and other rare epiphytes on lava rock outcrop in the Eifel (Rhineland-Palatinate, Germany). – *Herzogia* 17: 87–93.
- Aragón G. & Otálora M. A. G. (2004): Ecological and chorological novelties of the genus *Leptogium* in the Iberian Peninsula. – *Nova Hedwigia* 78: 353–366.
- Aragón G. & Sarrión F. J. (2003): Epiphytic *Verrucariaceae* in Mediterranean Spain. – *Nova Hedwigia* 77: 169–187.
- Aragón G., Sarrión F. J. & Martínez I. (2004): Epiphytic lichens on *Juniperus oxycedrus* in the Iberian Peninsula. – *Nova Hedwigia* 78: 45–56.
- Armstrong R. A. (2003): Lobe growth in foliose lichen thalli. – *Current Topics in Plant Biology* 4: 129–138.
- Armstrong R. A. (2004): Lichens, lichenometry and global warming. – *Microbiologist* 5: 32–35.
- Arnerup J., Höglberg N. & Thor G. (2004): Phylogenetic analysis of multiple loci reveal the population structure within *Letharia* in the Caucasus and Morocco. – *Mycological Research* 108: 311–316.
- Articus K. (2004): *Neuropogon* and the phylogeny of *Usnea* s. l. (*Parmeliaceae*, lichenized ascomycetes). – *Taxon* 53: 925–934.
- Ask P. & Nilsson S. G. (2004): Stand characteristics in colour-infrared aerial photographs as indicators of epiphytic lichens. – *Biodiversity and Conservation* 13: 529–542.
- Aslan A., Budak G. & Karabulut A. (2004): The amounts Fe, Ba, Sr, K, Ca and Ti in some lichens growing in Erzurum province (Turkey). – *Journal of Quantitative Spectroscopy and Radiative Transfer* 88: 423–431.
- Aslan A., Budak G., Tirasoglu E., Karabulut A., Karagöz Y., Apaydin G., Ertugral B. & Çevik U. (2004): Analysis of elements in some lichens by radioisotope X-ray fluorescence spectrometry. – *Fresenius Environmental Bulletin* 13: 740–747.
- Augusto S., Pinho P., Branquinho C., Pereira M. J., Soares A. & Catarino F. (2004): Atmospheric dioxin and furan deposition in relation to land-use and other pollutants: a survey with lichens. – *Journal of Atmospheric Chemistry* 49: 53–65.
- Batts J. E., Calder L. J. & Batts B. D. (2004): Utilizing stable isotope abundances of lichens to monitor environmental change. – *Chemical Geology* 204: 345–368.
- Batty K., Bates J. W. & Bell J. N. B. (2003): A transplant experiment on the factors preventing lichen colonization of oak bark in southeast England under declining SO₂ pollution. – *Canadian Journal of Botany* 81: 439–451.
- Benesperi R. & Tretiach M. (2004): Differential land snail damage to selected species of the lichen genus *Peltigera*. – *Biochemical Systematics and Ecology* 32: 127–138.
- Bergamaschi L., Rizzio E., Giaveri G., Profumo A., Loppi S. & Gallorini M. (2004): Determination of baseline element composition of lichens using samples from high elevations. – *Chemosphere* 55: 933–939.
- Berger F. & Aptroot A. (2002): Further contributions to the flora of lichens and lichenicolous fungi of the Azores. – *Arquipélago. Life and Marine Sciences* 19A: 1–12.
- Bezivin C., Tomasi S., Rouaud I., Delcros J. G. & Boustie J. (2004): Cytotoxic activity of compounds from the lichen: *Cladonia convoluta*. – *Planta Medica* 70: 874–877.
- Bhattacharya D., Friedl T. & Helms G. (2002): Vertical evolution and intragenic spread of lichen-fungal group I introns. – *Journal of Molecular Evolution* 55: 74–84.
- Bielczyk U. & Betleja L. (2003): The lichens of the ‘Bór na Czerwonem’ raised peat-bog in the Orawa-Nowy Targ Basin (southern Poland). – *Polish Botanical Journal* 48: 69–75.
- Bjerke J. W. & Elvebakk A. (2004): Distribution of the lichen genus *Flavocetraria* (*Parmeliaceae*, *Ascomycota*) in the Southern Hemisphere. – *New Zealand Journal of Botany* 42: 647–656.
- Bjerke J. W., Joly D., Nilsen L. & Brossard T. (2004): Spatial trends in usnic acid concentrations of the lichen *Flavocetraria nivalis* along local climatic gradients in the Arctic (Kongsfjorden, Svalbard). – *Polar Biology* 27: 409–417.
- Blanco O., Crespo A., Divakar P. K., Esslinger T. L., Hawksworth D. L. & Lumbsch H. T. (2004): *Melanelixia* and *Melanohalea*, two new genera segregated from *Melanelia* (*Parmeliaceae*) based on molecular and morphological data. – *Mycological Research* 108: 873–884.
- Blanco O., Crespo A., Elix J. A., Hawksworth D. L. & Lumbsch H. T. (2004): A molecular phylogeny and a new classification of parmelioid lichens containing *Xanthoparmelia*-type lichenan (*Ascomycota*: *Lecanorales*). – *Taxon* 53: 959–975.

- Borg L. & Aronsson M. (2004): Population size and ecology of *Pleurosticta acetabulum*, *Anaptychia ciliaris* and *Ramalina fraxinea* in avenues and churchyards in Kalmar county, SE Sweden. – *Graphis Scripta* 16: 37–42.
- Bradwell T. (2004): Lichenometric dating in southeast Iceland: the size frequency approach. – *Geografiska Annaler, Series A*, 86: 31–41.
- Breuss O. (2004): Neue Flechtenfunde, vorwiegend pyrenocarper Arten, aus Oberösterreich. – *Österreichische Zeitschrift für Pilzkunde* 13: 267–275.
- Bucar F., Schneider I., Ögmundsdóttir H. & Ingólfssdóttir K. (2004): Anti-proliferative lichen compounds with inhibitory activity on 12(S)-HETE production in human platelets. – *Phytomedicine* 11: 602–606.
- Budka D., Mesjasz-Przybyłowicz J. & Przybyłowicz W. J. (2004): Environmental pollution monitoring using lichens as bioindicators: a micro-PIXE study. – *Radiation Physics and Chemistry* 71: 783–784.
- Bukhov N. G., Govindachary S., Egorova E. A. & Carpentier R. (2004): Recovery of photosystem I and II activities during re-hydration of lichen *Hypogymnia physodes* thalli. – *Planta* 219: 110–120.
- Bungartz F., Elix J. A. & Nash T. H. (2004): The genus *Buellia* sensu lato in the Greater Sonoran Desert Region: saxicolous species with one-septate ascospores containing xanthenes. – *Bryologist* 107: 441–458.
- Bungartz F., Garvie L. A. J. & Nash T. H. (2004): Anatomy of the endolithic Sonoran Desert lichen *Verrucaria rubrocincta* Breuss: implications for biodeterioration and biomineralization. – *Lichenologist* 36: 55–73.
- Bungartz F. & Nash T. H. (2004): The *Buellia aethalea*-group in the Greater Sonoran Desert region with reference to similar species in North America. – *Bryologist* 107: 441–458.
- Bungartz F., Nash T. H. & Ryan B. D. (2004): Morphology and anatomy of chasmolithic versus epilithic growth: a taxonomic revision of inconspicuous saxicolous *Buellia* species from the Sonoran Desert Region generally ascribed to the "*Buellia punctata*" group. – *Canadian Journal of Botany* 82: 540–562.
- Burgaz A. R. & Martínez I. (2003): Flora Liquenológica Ibérica. *Peltigerales: Lobariaceae, Nephromataceae, Peltigeraceae*. – Sociedad Española de Liquenología (SEL), Murcia. [61 pp.]
- Buschbom J. & Mueller G. (2004): Resolving evolutionary relationships in the lichen-forming genus *Porpidia* and related allies (*Porpidiaceae, Ascomycota*). – *Molecular Phylogenetics and Evolution* 32: 66–82.
- Cabral J. P. (2003): Copper toxicity to five *Parmelia* lichens in vitro. – *Environmental and Experimental Botany* 49: 237–250.
- Campbell J. & Fredeen A. L. (2004): *Lobaria pulmonaria* abundance as an indicator of macrolichen diversity in Interior Cedar–Hemlock forests of east-central British Columbia. – *Canadian Journal of Botany* 82: 970–982.
- Carter N. E. A. & Viles H. A. (2003): Experimental investigations into the interactions between moisture, rock surface temperatures and an epilithic lichen cover in the bioprotection of limestone. – *Building and Environment* 38: 1225–1234.
- Carter N. E. A. & Viles H. A. (2004): Lichen hotspots: raised rock temperatures beneath *Verrucaria nigrescens* on limestone. – *Geomorphology* 62: 1–16.
- Caviglia A. M., Nicora P., Giordani P., Brunialti G. & Modenesi P. (2001): Oxidative stress and usnic acid content in *Parmelia caperata* and *Parmelia soledians* (*Lichenes*). – *Il Farmaco* 56: 379–382.
- Cezanne R., Eichler M., Lumbsch H. T. & Zimmermann D. G. (2003): *Moelleropsis humida* – eine übersehene Flechte? – *Herzogia*. 16: 161–166.
- Chaves J. L., Lücking R., Sipman H. J. M., Umaña L. & Navarro E. (2004): A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Dictyonema* (*Polyporales: Atheliaceae*). – *Bryologist* 107: 242–249.
- Christensen S. N. (2004): Epilithic lichen vegetation of corrugated asbestos-cement roof tiles in the Copenhagen area. – *Graphis Scripta* 15: 7–13.
- Clayden S. R., Pentecost A. & Dawson R. J. M. (2004): Growth of the lichen *Rhizocarpon lecanorinum*, with comments on Aplin-Hill and lichenometric curves. – *Symbiosis* 37: 379–393.
- Clerc P. (2004): Les champignons lichénisés de Suisse. Catalogue bibliographique complété par des données sur la distribution et l'écologie des espèces. – *Cryptogamica Helvetica* 19: 1–320.
- Clerc P. (2004): Notes on the genus *Usnea* Adanson. II. – *Bibliotheca Lichenologica* 88: 79–90.
- Cordeiro L. M. C., Iacomini M. & Stocker-Wörgötter E. (2004): Culture studies and secondary compounds of six *Ramalina* species. – *Mycological Research* 108: 489–497.

- Cordeiro L. M. C., Stocker-Wörgötter E., Gorin P. A. J. & Iacomini M. (2003): Comparative studies of the polysaccharides from species of the genus *Ramalina* – lichenized fungi – of three distinct habitats. – *Phytochemistry* 63: 967–975.
- Coxson D., Stevenson S. & Campbell J. (2003): Short-term impacts of partial cutting on lichen retention and canopy microclimate in an Engelmann spruce-subalpine fir forest in northcentral British Columbia. – *Canadian Journal of Forest Research* 33: 830–841.
- Coxson D. S. & Coyle M. (2003): Niche partitioning and photosynthetic response of alectorioid lichens from subalpine spruce-fir forest in north-central British Columbia, Canada: the role of canopy microclimate gradients. – *Lichenologist* 35: 157–175.
- Coxson D. S. & Curteanu M. (2002): Decomposition of hair lichens (*Alectoria sarmentosa* and *Bryoria* spp.) under snowpack in montane forest, Cariboo Mountains, British Columbia. – *Lichenologist* 34: 395–402.
- Crespo A., Divakar P. K., Argüello A., Gasca C. & Hawksworth D. L. (2004): Molecular studies on *Punctelia* species of the Iberian Peninsula, with an emphasis on specimens newly colonizing Madrid. – *Lichenologist* 36: 299–308.
- Cubero O. F., Crespo A., Esslinger T. L. & Lumbsch H. T. (2004): Molecular phylogeny of the genus *Physconia* (*Ascomycota*, *Lecanorales*) inferred from a Bayesian analysis of nuclear ITS rDNA sequences. – *Mycological Research* 108: 498–505.
- Cuny D., Denayer F. O., de Foucault B., Schumacker R., Colein P. & Van Haluwyn C. (2004): Patterns of metal soil contamination and changes in terrestrial cryptogamic communities. – *Environmental Pollution* 129: 289–297.
- Cuny D., Van Haluwyn C., Shirali P., Zerimech F., Jérôme L. & Haguenoer J. M. (2004): Cellular impact of metal trace elements in terricolous lichen *Diploschistes muscorum* (Scop.) R. Sant. – identification of oxidative stress biomarkers. – *Water, Air and Soil Pollution* 152: 55–69.
- Czarnota P. & Kukwa M. (2004): Some sorediate lichens and lichenicolous fungi new to Poland. – *Graphis Scripta* 15: 24–32.
- Czczuga B., Czczuga-Semeniuk E., Galun M. & Mukhtar A. (2004): Pigmentation changes in *Xanthoria parietina* (L.) Th. Fr. – *Journal of the Hattori Botanical Laboratory* 95: 293–300.
- Czczuga B., Czczuga-Semeniuk E. & Hammer S. (2004): Carotenoids in the thalli of *Cladonia grayi* Merrill from sites of varied degrees of insolation. – *Journal of the Hattori Botanical Laboratory* 95: 285–291.
- Czczuga B., Czczuga-Semeniuk E., Vitikainen O. & Ahti T. (2004): Carotenoids in species of the *Peltigera* genus from green-algae and cyanobacteria as phyto- and photobiont. – *Journal of the Hattori Botanical Laboratory* 96: 281–290.
- Dahlman L., Persson J., Näsholm T. & Palmqvist K. (2003): Carbon and nitrogen distribution in the green algal lichens *Hypogymnia physodes* and *Platismatia glauca* in relation to nutrient supply. – *Planta* 217: 41–48.
- Dahlman L., Persson J., Palmqvist K. & Näsholm T. (2004): Organic and inorganic nitrogen uptake in lichens. – *Planta* 219: 459–467.
- Daillant O., Kirchner G., Pigrée G. & Porstendörfer J. (2004): Lichens as indicators of tritium and radiocarbon contamination. – *Science of the Total Environment* 323: 253–262.
- de Vera J. P., Horneck G., Rettberg B. & Ott S. (2004): The potential of the lichen symbiosis to cope with the extreme conditions of outer space II: germination capacity of lichen ascospores in response to simulated space conditions. – *Advances in Space Research* 33: 1236–1243.
- den Herder M., Kitöviita M.-M. & Niemelä P. (2003): Growth of reindeer lichens and effect of reindeer grazing on ground cover vegetation in a Scots pine forest and a subarctic heathland in Finnish Lapland. – *Ecography* 26: 3–12.
- DePriest P. (2004): Early molecular investigations of lichen-forming symbionts: 1986–2001. – *Annual Review of Microbiology* 58: 273–301.
- Desbenoit B., Galin E. & Akkouche S. (2004): Simulating and modeling lichen growth. – *Computer Graphics Forum* 23: 341–350.
- Dettki H. & Esseen P.-A. (2003): Modelling long-term effects of forest management on epiphytic lichens in northern Sweden. – *Forest Ecology and Management* 175: 223–238.
- Di Lella L. A., Frati L., Loppi S., Protano G. & Riccobono F. (2003): Lichens as biomonitors of uranium and other trace elements in an area of Kosovo heavily shelled with depleted uranium rounds. – *Atmospheric Environment* 37: 5445–5449.

- Dolnik C. (2003): Artenzahl-Areal-Beziehungen von Wald- und Offenlandgesellschaften – ein Beitrag zur Erfassung der botanischen Artenvielfalt unter besonderer Berücksichtigung der Flechten und Moose am Beispiel des Nationalparks Kurische Nehrung (Russland). – *Mitteilungen der Arbeitsgemeinschaft Geobotanik in Schleswig-Holstein und Hamburg* 62: 1–183.
- Edwards H. G. M., Newton E. M. & Wynn-Williams D. D. (2003): Molecular structural studies of lichen substances II: atranorin, gyrophoric acid, fumarprotocetraric acid, rhizocarpic acid, calycin, pulvinic dilactone and usnic acid. – *Journal of Molecular Structure* 651–653: 27–37.
- Edwards H. G. M., Newton E. M., Wynn-Williams D. D. & Coombes S. R. (2003): Molecular structural studies of lichen substances I: parietin and emodin. – *Journal of Molecular Structure* 648: 49–59.
- Edwards H. G. M., Seaward M. R. D., Attwood S. J., Little S. J., de Oliveira L. F. C. & Tretiach M. (2003): FT-Raman spectroscopy of lichens on dolomitic rocks: an assessment of metal oxalate formation. – *Analyst* 128: 1218–1221.
- Elix J. A. & Tønsberg T. (2004): Notes on the chemistry of some lichens, including four species of *Lepraria*. – *Graphis Scripta* 16: 43–45.
- Ellis C. (2004): The lichen ecology of aspen woods – a preliminary analysis. – *British Lichen Society Bulletin* 94: 23–27.
- Ellis C. J., Crittenden P. D. & Scrimgeour C. M. (2004): Soil as a potential source of nitrogen for mat-forming lichens. – *Canadian Journal of Botany* 82: 145–149.
- Ellis C. J., Crittenden P. D., Scrimgeour C. M. & Ashcroft C. (2003): The natural abundance of ^{15}N in mat-forming lichens. – *Oecologia* 136: 115–123.
- Emmerer B. & Hafellner J. (2003): Zur aktuellen Vegetation auf Halden historischer Bergbaue in den Hochlagen der Niederen Tauern. – *Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark* 133: 37–79.
- Ertz D. & Diederich P. (2004): Revision of *Trimmatothele* (*Verrucariaceae*), and description of *Oevstedalia* for *Trimmatothelopsis antarctica*, a new lichen genus with true ascoconidia. – *Mycological Progress* 3: 229–236.
- Etayo J. & Marbach B. (2003): *Hafellia alisioae* and *H. gomerana* (lichenized *Ascomycetes*, *Physciaceae*), two new species from the Canary Islands, with a key to all known corticolous species. – *Lichenologist* 35: 369–375.
- Etayo J. & Mayrhofer H. (2003): *Thelenella melanospora* (*Thelenellaceae*, lichenized *Ascomycetes*), a new species from the Mediterranean region. – *Nova Hedwigia* 77: 109–114.
- Eversman S. & Horton D. (2004): Recolonization of burned substrates by lichens and mosses in Yellowstone National Park. – *Northwest Science* 78: 85–92.
- Fałtynowicz W. (2003): The lichens, lichenicolous and allied fungi of Poland. An annotated checklist. – W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków. [435 pp.]
- Favero-Longo S. E., Isocrono D. & Piervittori R. (2004): Lichens and ultramafic rocks: a review. – *Lichenologist* 36: 391–404.
- Ferraro L. I. (2004): Morphological diversity in the hyphophores of *Gomphillaceae* (*Ostropales*, lichenized *Ascomycetes*). – *Fungal Diversity* 15: 153–169.
- Ferretti M., Brambilla E., Brunialti G., Fornasier F., Mazzali C., Giordani P. & Nimis P. L. (2004): Reliability of different sampling densities for estimating and mapping lichen diversity in biomonitoring studies. – *Environmental Pollution* 127: 249–256.
- Francolini I., Norris P., Piozzi A., Donelli G. & Stoodley P. (2004): Usnic acid, a natural antimicrobial agent able to inhibit bacterial biofilm formation on polymer surfaces. – *Antimicrobial Agents and Chemotherapy* 48: 4360–4365.
- Frödén P. & Lassen P. (2004): Typification and emendation of *Seiropora* Poelt to include species segregated from *Teloschistes* Norman. – *Lichenologist* 36: 289–298.
- Fryday A. (2004): Lichens of the Scottish mountains. – *British Lichen Society Bulletin* 94: 4–7.
- Fryday A. M. (2004): A new species of *Fuscopannaria* with a green photobiont, and other taxonomic innovations and new records of lichenized-fungi from Alaska. – *Bryologist* 107: 173–179.
- Fryday A. M. (2004): A re-assessment of the genus *Gongylia* in North America, and a new species of *Protothelenella* from Colorado. – *Bryologist* 107: 232–233.
- Fryday A. M. & Coppins B. J. (2004): A reassessment of the genera *Chromatochlamys* and *Thelenella*, and a new species of *Strigula* from the British Isles. – *Lichenologist* 36: 89–95.

- Gaio-Oliveira G., Dahlman L., Máguas C. & Palmqvist K. (2004): Growth in relation to microclimatic conditions and physiological characteristics of four *Lobaria pulmonaria* populations in two contrasting habitats. – *Ecography* 27: 13–28.
- Gaio-Oliveira G., Dahlman L., Palmqvist K. & Máguas C. (2004): Ammonium uptake in the nitrophytic lichen *Xanthoria parietina* and its effects on vitality and balance between symbionts. – *Lichenologist* 36: 75–86.
- Garty J. (2003): About the applicability of the lichen *Ramalina maciformis* (lichenized *Ascomycetes*, *Ramalinaceae*) as a biomonitor of air pollution in the Negev Desert, Israel. – *Bibliotheca Lichenologica* 86: 243–255.
- Garty J., Tomer S., Levin T. & Lehr H. (2003): Lichens as biomonitors around a coal-fired power station in Israel. – *Environmental Research*: 91: 186–198.
- Gauslaa Y. & Solhaug K. A. (2004): Photoinhibition in lichens depends on cortical characteristics and hydration. – *Lichenologist* 36: 133–143.
- Gilbert O. L. (2003): The lichen flora of unprotected soft sea cliffs and slopes. – *Lichenologist* 35: 245–254.
- Gilbert O. L. (2004): The phenology of *Sarcosagium campestre* observed over three years. – *Lichenologist* 36: 159–161.
- Giordani P. & Brunialti G. (2002): Anatomical and histochemical differentiation in lobes of the lichen *Lobaria pulmonaria*. – *Mycological Progress* 1: 131–136.
- Gombert S., Asta J. & Seaward M. R. D. (2004): Assessment of lichen diversity by index of atmospheric purity (IAP), index of human impact (IHI) and other environmental factors in an urban area (Grenoble, southeast France). – *Science of the Total Environment* 324: 183–199.
- González C. M., Pignata M. L. & Orellana L. (2003): Applications of redundancy analysis for the detection of chemical response patterns to air pollution in lichen. – *Science of the Total Environment* 312: 245–253.
- Grube M., Baloch E. & Arup U. (2004): A phylogenetic study of the *Lecanora rupicola* group (*Lecanoraceae*, *Ascomycota*). – *Mycological Research* 108: 506–514.
- Grube M., Baloch E. & Lumbsch H. T. (2004): The phylogeny of *Porinaceae* (*Ostropomycetidae*) suggests a neotenic origin of perithecia in *Lecanoromycetes*. – *Mycological Research* 108: 1111–1118.
- Gunnarsson B., Hake M. & Hultengren S. (2004): A functional relationship between species richness of spiders and lichens in spruce. – *Biodiversity and Conservation* 13: 685–693.
- Guschina I. A., Dobson G. & Harwood J. L. (2003): Lipid metabolism in cultured lichen photobionts with different phosphorus status. – *Phytochemistry* 64: 209–217.
- Hafellner J. (2003): Ein Beitrag zur Flechtenflora der Fischbacher Alpen (Steiermark). – *Fritschiana* 41: 21–40.
- Hafellner J. (2003): Ein Beitrag zur Flechtenflora des Jogllandes (Steiermark). – *Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark* 133: 81–97.
- Hafellner J. (2003): Ein Beitrag zur Flechtenflora für die Naturräume Weststeirisches Hügelland, Sausal und Windische Bühel (Steiermark). – *Fritschiana* 43: 47–63.
- Hafellner J. (2004): A further evolutionary lineage to lichenicolous growth in *Physciaceae* (*Lecanorales*). – *Bibliotheca Lichenologica* 88: 175–186.
- Hafellner J. (2004): A revision of *Maronella laricina* and *Piccolia ochrophora*. – *Symbolae Botanicae Upsalienses* 34: 87–96.
- Hafellner J. (2004): Notes on *Scoliciosporum intrusum*. – *Fritschiana* 49: 29–41.
- Hafellner J. & Obermayer W. (2004): Beobachtungen an einer sorediösen Population von *Pseudevernia furfuracea*. – *Herzogia* 17: 45–50.
- Hafellner J., Obermayer W., Breuss O. & Türk R. (2003): Flechtenfunde in den Schladminger Tauern in der Steiermark (BLAM-Exkursion 2001). – *Herzogia* 16: 187–206.
- Hansen E. S. (2004): An initial study of lichen growth on boulders and rocks near the Mittivakkat Gletscher, South East Greenland. – *Graphis Scripta* 15: 33–38.
- Harada H., Okamoto T. & Yoshimura I. (2004): A checklist of lichens and lichen-allies of Japan. – *Lichenology* 2/2: 1–165.
- Härkönen M. & Vänskä H. (2004): Corticolous myxomycetes and lichens in the Botanical Garden in Helsinki, Finland: a comparison after decades of recovering from air pollution. – *Systematics and Geography of Plants* 74: 183–187.

- Hauck M. & Zoller T. (2003): Copper sensitivity of soredia of the epiphytic lichen *Hypogymnia physodes*. – *Lichenologist* 35: 271–274.
- Hawksworth D. L. (2003): The lichenicolous fungi of Great Britain and Ireland: an overview and annotated checklist. – *Lichenologist* 35: 191–232.
- Hawksworth D. L. (2004): Fungi living on lichens: a source of unexplored diversity. – *British Wildlife* 15: 192–199.
- Hawksworth D. L. (2004): Rediscovery of the original material of Osbeck's „*Lichen chinensis*“ and the reinstatement of the name *Parmotrema perlatum* (*Parmeliaceae*). – *Herzogia* 17: 37–44.
- Hawksworth D. L. & Coppins B. J. (2003): *Bryoria tenuis* (*Parmeliaceae*) new to the British Isles, and either awaiting rediscovery or extinct. – *Lichenologist* 35: 361–364.
- Hedenås H. & Ericson L. (2004): Aspen lichens in agricultural and forest landscapes: the importance of habitat quality. – *Ecography* 27: 521–531.
- Hermansson J. & Thor G. (2004): *Bryoria tortuosa* discovered as new to Sweden. – *Graphis Scripta* 15: 39–41.
- Hermansson J. & Thor G. (2004): *Byssoloma subdiscordans* and *Usnea substerilis* new to Sweden. – *Graphis Scripta* 15: 42–44.
- Herrera-Campos M. L. Á., Lücking R., Pérez R.-E., Campos Á., Colín P. M. & Peña A. B. (2004): The foliicolous lichen flora of Mexico. V. Biogeographical affinities, altitudinal preferences, and an updated checklist of 293 species. – *Lichenologist* 36: 309–327.
- Hestmark G., Skogedal O. & Skullerud Ø. (2004): Growth in the alpine saxicolous lichens *Allantoparmelia alpicola* and *Melanelia stygia*. – *Nova Hedwigia* 78: 301–309.
- Hestmark G., Skogedal O. & Skullerud Ø. (2004): Growth, reproduction, and population structure in four alpine lichens during 240 years of primary colonization. – *Canadian Journal of Botany* 82: 1356–1362.
- Holub S. M. & Lajtha K. (2003): Mass loss and nitrogen dynamics during the decomposition of a ¹⁵N-labeled N₂-fixing epiphytic lichen, *Lobaria oregana*. – *Canadian Journal of Botany* 81: 698–705.
- Honegger R. (2004): Fine structure of the interaction of *Leprocaulon microscopicum* with its green algal photobiont, *Dictyochooropsis symbiontica*. – *Bibliotheca Lichenologica* 88: 201–210.
- Honegger R., Zippler U., Gansner H. & Scherrer S. (2004): Mating systems in the genus *Xanthoria*, (lichen-forming *Ascomycetes*). – *Mycological Research* 108: 480–488.
- Honegger R., Zippler U., Scherrer S. & Dyer P. S. (2004): Genetic diversity in *Xanthoria parietina* (L.) Th. Fr. (lichen-forming ascomycete) from worldwide locations. – *Lichenologist* 36: 381–390.
- Hultengren S. & Arvidsson L. (2003): Jordbrukslandskapets lavar – habitat och livsvillkor [Lichens in the agricultural landscape – habitats and threats]. – *Fauna och Flora* 98: 32–39.
- Hultengren S., Gralén H. & Pleijel H. (2004): Recovery of the epiphytic lichen flora following air quality improvement in south-west Sweden. – *Water, Air, and Soil Pollution* 154: 203–211.
- Huneck S., Feige G. B. & Schmidt J. (2004): Chemie von *Cladonia furcata* und *Cladonia rangiformis*. – *Herzogia* 17: 51–58.
- Huneck S., Lumbsch H. T., Porzel A. & Schmidt J. (2004): Die Verteilung von Flechteninhaltsstoffen in *Lecanora muralis* und *Lecidea inops* und die Abhängigkeit der Usninsäure-Konzentration vom Substrat und von den Jahreszeiten bei *Lecanora muralis*. – *Bibliotheca Lichenologica* 88: 211–221.
- Hyvärinen M., Härdling R. & Tuomi J. (2002): Cyanobacterial lichen symbiosis: the fungal partner as an optimal harvester. – *Oikos* 98: 498–504.
- Ihlen P. G. (2004): Taxonomy of the non-yellow species of *Rhizocarpon* (*Rhizocarpaceae*, lichenized *Ascomycota*) in the Nordic countries, with hyaline and muriform ascospores. – *Mycological Research* 108: 533–570.
- Ingólfssdóttir K. (2002): Molecules of interest: Usnic acid. – *Phytochemistry* 61: 729–736.
- Ingólfssdóttir K., Guðmundsdóttir G. F., Ögmundsdóttir H. M., Paulus K., Haraldsdóttir S., Kristinsson H. & Bauer R. (2002): Effects of tenuiorin and methyl orsellinate from the lichen *Peltigera leucophlebia* on 5-/15-lipoxygenases and proliferation of malignant cell lines in vitro. – *Phytomedicine* 9: 654–658.
- Jahns H. M., Hardt K. & Ott S. (2004): Sexual reproduction and growth-pattern in *Cladonia rangiferina*. – *Bibliotheca Lichenologica* 88: 223–228.
- Jahren A. H., Porter S. & Kuglitsch J. J. (2004): Lichen metabolism identified in Early Devonian terrestrial organisms. – *Geology* 31: 99–102.
- James P. W. (2003): Aide mémoire: *Usnea*. – British Lichen Society, London. [32 pp.]
- John V. & Breuss O. (2004): Flechten der östlichen Schwarzmeer-Region in der Türkei (BLAM Exkursion 1997). – *Herzogia* 17: 137–156.

- John V., Seaward M. R. D., Sipman H. J. M. & Zedda L. (2004): Lichens and lichenicolous fungi from Syria, including a first checklist. – *Herzogia* 17: 157–177.
- Joneson S., Kashiwadani H., Tschabanenko S. & Gage S. (2004): *Ramalina* of the Kuril Islands. – *Bryologist* 107: 98–106.
- Jørgensen P. M. (2004): *Psoroma tenue* var. *boreale*, an overlooked, widespread, arctic-alpine lichen. – *Graphis Scripta* 15: 60–64.
- Jüriado I., Paal J. & Liira J. (2003): Epiphytic and epixylic lichen species diversity in Estonian natural forests. – *Biodiversity and Conservation* 12: 1587–1607.
- Kainz C. & Rambold G. (2004): A phylogenetic study of the lichen genus *Protoblastenia* (*Lecanorales*, *Psoraceae*) in Central Europe. – *Bibliotheca Lichenologica* 88: 267–299.
- Kapusta P., Szarek-Lukaszewska G. & Kiszka J. (2004): Spatial analysis of lichen species richness in a disturbed ecosystem (Niepołomice Forest, S Poland). – *Lichenologist* 36: 249–260.
- Khodosovtsev A., Kondratyuk S. & Kärnefelt I. (2004): *Candelariella boikoi*, a new lichen species from Eurasia. – *Graphis Scripta* 16: 11–15.
- Kiszka J. & Grodzińska K. (2004): Lichen flora and air pollution in the Niepołomice Forest (S Poland) in 1960–2000. – *Biologia, Bratislava* 59: 25–37.
- Kohlmeyer J., Hawksworth D. L. & Volkmann-Kohlmeyer B. (2004): Observations on two marine and maritime “borderline” lichens: *Mastodia tessellata* and *Collempsidium pelvetiae*. – *Mycological Progress* 3: 51–56.
- Kossowska M. (2003): Lichens growing on *Abies alba* in the Karkonosze National Park (South West Poland) – the results of the first stage of the conservation program. – *Botanica Lithuanica* 9: 65–70.
- Kossowska M. & Fabiszewski J. (2004): Threatened lichens of Lower Silesia, Poland. – *Acta Societatis Botanicorum Poloniae* 73: 139–150.
- Kowalewska A. & Kukwa M. (2003): Preliminary studies of the *Cladonia chlorophaea* group (*Cladoniaceae*, *Ascomycota Lichenisati*) in northern Poland. – *Botanica Lithuanica* 9: 135–143.
- Kowalewska A. & Kukwa M. (2004): *Cladonia metacorallifera* (lichenized *Ascomycota*, *Cladoniaceae*) new to Poland and additional record from Slovakia. – *Biologia, Bratislava* 59: 433–434.
- Kowalewska A. & Kukwa M. (2004): New records of *Cladonia mesomorpha* (*Cladoniaceae*, lichenized *Ascomycota*) from Europe. – *Herzogia* 17: 103–105.
- Krzewicka B. (2004): The lichen genera *Lasallia* and *Umbilicaria* in the Polish Tatra Mts. – *Polish Botanical Studies* 17: 1–88.
- Kubiak D. & Sparrius L. B. (2004): *Bacidia adastrae*, *B. brandii* and *B. neosquamulosa* found in North-Eastern Poland. – *Graphis Scripta* 16: 61–64.
- Lange O. L. & Green T. G. A. (2004): Lichens show that fungi can acclimate their respiration to seasonal changes in temperature. – *Oecologia* 142: 11–19.
- Lange O. L. & Green T. G. A. (2004): Photosynthetic performance of the squamulose soil-crust lichen *Squamarina lentigera*: laboratory measurements and long-term monitoring of CO₂ exchange in the field. – *Bibliotheca Lichenologica* 88: 363–390.
- Larsen R. & Søchting U. (2003): *Ramonia chrysophaea* new to Denmark. – *Graphis Scripta* 14: 7–9.
- Legaz M. E., Fontaniella B., Millanes A. M. & Vicente C. (2004): Secreted arginases from phylogenetically far-related lichen species act as cross-recognition factors for two different algal cells. – *European Journal of Cell Biology* 83: 435–446.
- Leuckert C., Wirth V., Kümmerling H. & Heklau M. (2004): Chemical lichen analyses XIV. *Lepraria nivalis* J. R. Laundon and *Lepraria flavescens* Cl. Roux & Tønsberg. – *Bibliotheca Lichenologica* 88: 393–407.
- Lidén M., Pettersson M., Bergsten U. & Lundmark T. (2004): Artificial dispersal of endangered epiphytic lichens: a tool for conservation in boreal forest landscapes. – *Biological Conservation* 118: 431–442.
- Lindahl P., Roos P., Eriksson M. & Holm E. (2004): Distribution of Np and Pu in Swedish lichen samples (*Cladonia stellaris*) contaminated by atmospheric fallout. – *Journal of Environmental Radioactivity* 73: 73–85.
- Lõhmus P. (2003): Composition and substrata of forest lichens in Estonia: a meta-analysis. – *Folia Cryptogamica Estonica* 40: 19–38.
- Loppi S., Boscagli A. & De Dominicis V. (2004): Ecology of soil lichens from Pliocene clay badlands of central Italy in relation to geomorphology and vascular vegetation. – *Catena* 55: 1–15.
- Loppi S. & Frati L. (2004): Influence of tree substrate on the diversity of epiphytic lichens: comparison between *Tilia platyphyllos* and *Quercus ilex* (central Italy). – *Bryologist* 107: 340–344.

- Loppi S., Frati L., Paoli L., Bigagli V., Rossetti C., Bruscoli C. & Corsini A. (2004): Biodiversity of epiphytic lichens and heavy metal contents of *Flavoparmelia caperata* thalli as indicators of temporal variations of air pollution in the town of Montecatini Terme (central Italy). – *Science of the Total Environment* 326: 113–122.
- Loppi S. & Pirintsos S. A. (2003): Epiphytic lichens as sentinels for heavy metal pollution at forest ecosystems (central Italy). – *Environmental Pollution* 121: 327–332.
- Loppi S., Riccobono F., Zhang Z. H., Savic S., Ivanov D. & Pirintsos S. A. (2003): Lichens as biomonitors of uranium in the Balkan area. – *Environmental Pollution* 125: 277–280.
- Lücking R. (2004): A revised key to foliicolous *Porinaceae* (*Ascomycota: Trichotheliales*). – *Bibliotheca Lichenologica* 88: 409–426.
- Lücking R., Stuart B. L. & Lumbsch H. T. (2004): Phylogenetic relationships of *Gomphillaceae* and *Asterothyriaceae*: evidence from a combined Bayesian analysis of nuclear and mitochondrial sequences. – *Mycologia* 96: 283–294.
- Lumbsch H. T. (2004): Comparison of homoplasy supports that characters have no a priori value in phylogenetic analyses of lichenized ascomycetes. – *Bibliotheca Lichenologica* 88: 427–431.
- Lumbsch H. T., Mangold A., Lücking R., García M. A. & Martín M. P. (2004): Phylogenetic position of the genera *Nadvornikia* and *Pyrgillus* (*Ascomycota*) based on molecular data. – *Symbolae Botanicae Upsalienses* 34: 9–17.
- MacGillivray T. & Helleur R. (2001): Analysis of lichens under environmental stress using TMAH thermochemolysis–gas chromatography. – *Journal of Analytical and Applied Pyrolysis* 58–59: 465–480.
- Mallavadhani U. V., Sudhakar A. V. S., Mahapatra A., Narasimhan K., Thirunavokkarasu M. & Elix J. A. (2004): Phenolic and steroidal constituents of the lichen *Usnea longissima*. – *Biochemical Systematics and Ecology* 32: 95–97.
- Martín M. P., Coucheron D. H. & Johansen S. (2003): Structural features and evolutionary considerations of group IB introns in SSU rDNA of the lichen fungus *Teloschistes*. – *Fungal Genetics and Biology* 40: 252–260.
- Martínez I. & Aragón G. (2004): The *Lecanora varia* group in Spain: species with amphithecial cortex. – *Bryologist* 107: 222–230.
- Martínez I., Burgaz A. R., Vitikainen O. & Escudero A. (2003): Distribution patterns in the genus *Peltigera* Willd. – *Lichenologist* 35: 301–323.
- Mattsson J.-E. & Articus K. (2004): The monophyletic groups of cetrarioid lichens. – *Symbolae Botanicae Upsalienses* 34: 237–244.
- McKinze K. M., Orwin J. F. & Bradwell T. (2004): Re-dating the moraines at Skálafellsjökull and Heinabergsjökull using different lichenometric methods: Implications for the timing of the Icelandic Little Ice Age maximum. – *Geografiska Annaler: Series A, Physical Geography* 86A: 319–335.
- Miadlikowska J. & Lutzoni F. (2004): Phylogenetic classification of peltigeralean fungi (*Peltigerales, Ascomycota*) based on ribosomal RNA small and large subunits. – *American Journal of Botany* 91: 449–464.
- Millanes A. M., Fontaniella B., Legaz M. E. & Vicente C. (2003): Histochemical detection of an haematommoyl alcohol dehydrogenase in the lichen *Evernia prunastri*. – *Plant Physiology and Biochemistry* 41: 786–791.
- Millanes A. M., Fontaniella B., García M. L., Solas M. T., Vicente C. & Legaz M. E. (2004): Cytochemical location of urease in the cell wall of two different lichen phycobionts. – *Tissue and Cell* 36: 373–377.
- Moberg R. (2004): The lichen genus *Heterodermia* in Europe and the Macaronesian Islands. – *Bibliotheca Lichenologica* 88: 453–463.
- Mohr F., Ekman S. & Heegard E. (2004): Evolution and taxonomy of the marine *Collemopsisidium* species (lichenized *Ascomycota*) in north-west Europe. – *Mycological Research* 108: 515–532.
- Molina M. C., Crespo A., Blanco O., Lumbsch H. T. & Hawksworth D. L. (2004): Phylogenetic relationships and species concepts in *Parmelia* s.str. (*Parmeliaceae*) inferred from nuclear ITS rDNA and beta-tubulin sequences. – *Lichenologist* 36: 37–54.
- Molina M. C., Crespo A., Vicente C. & Elix J. A. (2003): Differences in the composition of phenolics and fatty acids of cultured mycobiont and thallus of *Physconia distorta*. – *Plant Physiology and Biochemistry* 41: 175–180.
- Motiejūnaitė J. (2003): Aquatic lichens in Lithuania. Lichens on submerged alder roots. – *Herzogia* 16: 113–121.

- Motiejūnaitė J. & Andersson L. (2003): Contribution to the Lithuanian flora of lichens and allied fungi. – *Botanica Lithuanica* 9: 71–88.
- Motiejūnaitė J., Kukwa M., Czarnota P., Prigodina-Lukošienė I., Himelbrant D., Kuznetsova E. & Kowalewska A. (2003): Lichens and allied fungi collected during the 15th symposium of Baltic mycologists and lichenologists in Birštonas, Lithuania. – *Botanica Lithuanica* 9: 109–119.
- Mrak T., Mayrhofer H. & Batič F. (2004): Contributions to the lichen flora of Slovenia XI. Lichens from the surrounding area of Bohinj Lake (Julian Alps). – *Herzogia* 17: 107–127.
- Myllys L., Stenroos S., Thell A. & Ahti T. (2003): Phylogeny of bipolar *Cladonia arbuscula* and *Cladonia mitis* (Lecanorales, Euascomycetes). – *Molecular Phylogenetics and Evolution* 27: 58–69.
- Nash T. H., Ryan B. D., Diederich P., Gries C. & Bungartz F. (eds.) (2004): Lichen flora of the Sonoran Desert Region, Vol. 2. – *Lichens Unlimited*, Arizona State University, Tempe. [742 pp.]
- Nimis P. L. & Martellos S. (2003): A second checklist of Italy with a thesaurus of synonyms. – *Museo Regionale di Scienze Naturali Saint-Pierre – Valle d'Aosta*. [192 pp.]
- Nimis P. L. & Tretiach M. (2004): Delimiting Tyrrhenian Italy: A lichen foray in the SW part of the peninsula. – *Bibliotheca Lichenologica* 88: 465–478.
- Nordin A. (2004): New species in *Tetramelas*. – *Lichenologist* 36: 355–359.
- Nordin A. (2004): Three species new to Sweden in material collected by Du Rietz. – *Graphis Scripta* 16: 20–22.
- Nybakken L., Solhaug K. A., Bilger W. & Gauslaa Y. (2004): The lichens *Xanthoria elegans* and *Cetraria islandica* maintain a high protection against UV-B radiation in Arctic habitats. – *Oecologia* 140: 211–216.
- Ohmura Y. & Kanda H. (2004): Taxonomic status of section *Neuropogon* in the genus *Usnea* elucidated by morphological comparisons and ITS rDNA sequences. – *Lichenologist* 36: 217–225.
- Ohnuki T., Aoyagi H., Kitatsuji Y., Samadfam M., Kimura Y. & Purvis O. W. (2004): Plutonium(VI) accumulation and reduction by lichen biomass: correlation with U(VI). – *Journal of Environmental Radioactivity* 77: 339–353.
- Oksanen I., Jokela J., Fewer D. P., Wahlsten M., Rikkinen J. & Sivonen K. (2004): Discovery of rare and highly toxic microcystins from lichen-associated cyanobacterium *Nostoc* sp. strain IO-102-I. – *Applied and Environmental Microbiology* 70: 5756–5763.
- O'Neal M. A. & Schoenenberger K. R. (2003): A *Rhizocarpon geographicum* growth curve for the Cascade Range of Washington and northern Oregon, USA. – *Quaternary Research* 60: 233–241.
- Opanowicz M. & Grube M. (2004): Photobiont genetic variation in *Flavocetraria nivalis* from Poland (Parmeliaceae, lichenized Ascomycota). – *Lichenologist* 36: 125–131.
- Orange A. (2003): *Cryptothele* in the British Isles. – *British Lichen Society Bulletin* 92: 32–34.
- Orange A. (2004): A remarkable new freshwater *Verrucaria* from Europe. – *Lichenologist* 36: 349–354.
- Orange A. (2004): The *Verrucaria fuscella* group in Great Britain and Ireland. – *Lichenologist* 36: 173–182.
- Orange A. (2004): *Verrucaria papillosa* is a synonym of *V. viridula*. – *Lichenologist* 36: 445–447.
- Ott S., Etges S., Rogge E. & Jahns H. M. (2004): Regeneration and determination in the peltigeralean lichens *Nephroma arcticum* and *Peltigera aphthosa*. – *Bibliotheca Lichenologica* 88: 527–538.
- Otte V. (2003): On C. G. Mosig's lichen herbarium, with emphasis on type material of lichens and fungi. – *Mycotaxon* 88: 393–399.
- Otte V. (2004): Flechten, Moose und lichenicole Pilze aus dem nordwestlichen Kaukasus – erster Nachtrag. – *Feddes Repertorium* 115: 155–163.
- Otte V. & Rätzel S. (2004): Kommentiertes Verzeichnis der Flechten und flechtenbewohnenden Pilze Brandenburgs, Deutschland – zweite Fassung. – *Feddes Repertorium* 115: 134–154.
- Palmqvist K., Dahlman L., Valladares F., Tehler A., Sancho L. G. & Mattsson J.-E. (2002): CO₂ exchange and thallus nitrogen across 75 contrasting lichen associations from different climate zones. – *Oecologia* 133: 295–306.
- Pannowitz S., Schlenz M., Green T. G. A., Sancho L. G. & Schroeter B. (2003): Are lichens active under snow in continental Antarctica? – *Oecologia* 135: 30–38.
- Paul A., Hauck M. & Langenfeld-Heyser R. (2004): Ultrastructural changes in soredia of the epiphytic lichen *Hypogymnia physodes* cultivated with manganese. – *Environmental and Experimental Botany* 52: 139–147.

- Pech P., Jomelli V., Baumgart-Kotarba M., Bravard J. P., Chardon M., Jacob N., Kedzia S., Kotarba A., Raczkowska Z. & Tsao C. (2003): A lichenometric growth curve in the French Alps: Ailefroide and Veneon valleys; Massif des Ecrins. – *Geodinamica Acta* 16: 187–193.
- Pentecost A. (2003): A preliminary investigation into ascospore characters in the *Graphidaceae* and *Opegraphaceae*. – *British Lichen Society Bulletin* 92: 35–39.
- Pereyra M. T., Prieto A., Bernabé M. & Leal J. A. (2003): Studies of new polysaccharides from *Lasallia pustulata* (L.) Hoffm. – *Lichenologist* 35: 177–185.
- Peršoh D., Beck A. & Rambold G. (2004): The distribution of ascus types and photobiontal selection in *Lecanoromycetes* (Ascomycota) against the background of a revised SSU nrDNA phylogeny. – *Mycological Progress* 3: 103–121.
- Piercey-Normore M. D. (2004): Selection of algal genotypes by three species of lichen fungi in the genus *Cladonia*. – *Canadian Journal of Botany* 82: 947–961.
- Piercey-Normore M. D., Hausner G. & Gibb E. A. (2004): Group I intron-like insertions in SSU rDNA of *Cladonia gracilis* and *C. rangiferina*. – *Lichenologist* 36: 365–380.
- Pöykkö H. & Hyvärinen M. (2003): Host preference and performance of lichenivorous *Eilema* spp. larvae in relation to lichen secondary metabolites. – *Journal of Animal Ecology* 72: 383–390.
- Pramyothin P., Janthasoot W., Pongnimitprasert N., Phrukudom S. & Ruangrunsi N. (2004): Hepatotoxic effect of (+)usnic acid from *Usnea siamensis* Wainio in rats, isolated rat hepatocytes and isolated rat liver mitochondria. – *Journal of Ethnopharmacology* 90: 381–387.
- Printzen C. & Kantvilas G. (2004): *Hertelidea*, genus novum Stereocaulacearum (*Ascomycetes lichenisati*). – *Bibliotheca Lichenologica* 88: 539–553.
- Printzen C. & Tønsberg T. (2004): New and interesting *Biatora*-species, mainly from North America. – *Symbolae Botanicae Upsalienses* 34: 343–357.
- Purvis O. W., Bailey E. H., McLean J., Kasama T. & Williamson B. J. (2004): Uranium biosorption by the lichen *Trapelia involuta* at a uranium mine. – *Geomicrobiology Journal* 21: 159–167.
- Purvis O. W., Chimonides J., Din V., Erotokritou L., Jeffries T., Jones G. C., Louwhoff S., Read H. & Spiro B. (2003): Which factors are responsible for the changing lichen floras of London? – *Science of the Total Environment* 310: 179–189.
- Pykälä J. (2004): Effects of new forestry practices on rare epiphytic macrolichens. – *Conservation Biology* 18: 831–838.
- Pykälä J. (2004): New finds of red-listed, rare and overlooked calicioid lichens and fungi in southern Finland. – *Graphis Scripta* 15: 17–23.
- Rätzel S., Otte V., Sipman H. J. M. & Fürstenow J. (2004): Bemerkenswerte Flechtenfunde aus Brandenburg VIII (incl. lichenicoler und lichenoider Pilze). – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 136: 321–340.
- Reeb V., Lutzoni F. & Roux C. (2004): Contribution of RPB2 to multilocus phylogenetic studies of the euascomycetes (*Pezizomycotina*, *Fungi*) with special emphasis on the lichen-forming *Acarosporaceae* and evolution of polyspory. – *Molecular Phylogenetics and Evolution* 32: 1036–1060.
- Rheault H., Drapeau P., Bergeron Y. & Esseen P.-A. (2003): Edge effects on epiphytic lichens in managed black spruce forests of eastern North America. – *Canadian Journal of Forest Research* 33: 23–32.
- Rikkinen J. (2003): Ecological and evolutionary role of photobiont-mediated guilds in lichens. – *Symbiosis* 34: 99–110.
- Romagni J. G., Rosell R. C., Nanayakkara N. P. D. & Dayan F. E. (2004): Ecophysiology and potential modes of action for selected lichen secondary metabolites. – In: Macías F. A., Galindo J. C. G., Molinillo J. M. G. & Cutler H. G. (eds.), *Allelopathy. Chemistry and Mode of Action of Allelochemicals*, p. 13–33, CRC Press, Boca Raton.
- Roux C. (2004): Les pycnides et conidies de *Lecanora vaenskaei* (lichens, *Lecanoraceae*). – *Mycotaxon* 90: 77–80.
- Roux C., Bricaud O., Ménard T., Gueidan C., Coste C. & Navarro-Rosinés P. (2003): Champignons lichénisés et lichénicoles de la France méridionale (Corse comprise): espèces nouvelles et intéressantes (9). – *Bulletin de la Société Linnéenne de Provence* 54: 125–141.
- Roux C., Gueidan C. & Clerc P. (2003): Lichens et champignons lichénicoles des Alpes françaises: espèces nouvelles et intéressantes. – *Cryptogamie, Mycologie* 24: 275–286.
- Sancho L. G., Palacios D., de Marcos J. & Valladares F. (2001): Geomorphological significance of lichen colonization in a present snow hollow: Hoya del Cuchillar de las Navajas, Sierra de Gredos (Spain). – *Catena* 43: 323–340.

- Sancho L. G. & Pintado A. (2004): Evidence of high annual growth rate for lichens in the maritime Antarctic. – *Polar Biology* 27: 312–319.
- Sanders W. B., Moe R. L. & Ascaso C. (2004): The intertidal marine lichen formed by the pyrenomycete fungus *Verrucaria tavaresiae* (Ascomycotina) and the brown alga *Petroderma maculiforme* (Phaeophyceae): thallus organization and symbiont interaction. – *American Journal of Botany* 91: 511–522.
- Santesson R., Moberg R., Nordin A., Tønsberg T. & Vitikainen O. (2004): Lichen-forming and lichenicolous fungi of Fennoscandia. – Museum of Evolution, Uppsala University, Uppsala. [359 pp.]
- Scherrer S. & Honegger R. (2003): Inter- and intraspecific variation of homologous hydrophobin (H1) gene sequences among *Xanthoria* spp. (lichen-forming ascomycetes). – *New Phytologist* 158: 375–389.
- Schiefelbein U. (2003): Bemerkenswerte Funde von Flechten und flechten-bewohnenden Pilzen aus dem Landkreis Uecker-Randow (Mecklenburg-Vorpommern). – *Herzogia* 16: 173–181.
- Schiefelbein U., Rätzl S. & Litterski B. (2003): Ergebnisse des 6. Flechtenkartierungstreffen in Mecklenburg-Vorpommern. – *Botanischer Rundbrief für Mecklenburg-Vorpommern* 38: 147–155.
- Schmitt I. & Lumbsch H. T. (2004): Molecular phylogeny of the *Pertusariaceae* supports secondary chemistry as an important systematic character set in lichen-forming ascomycetes – *Molecular Phylogenetics and Evolution* 33: 43–55.
- Seaward M. R. D. & Coppins B. J. (2004): Lichens and hypertrophication. – *Bibliotheca Lichenologica* 88: 561–572.
- Sedia E. G. & Ehrenfeld J. G. (2003): Lichens and bryophytes promote alternate stable plant communities in New Jersey Pinelands. – *Oikos* 100: 447–458.
- Sérusiaux E., Diederich P. & Lambinon J. (2004): Les macrolichens de Belgique, du Luxembourg et du nord de la France. Clés de détermination. – *Ferrantia* 40: 1–192.
- Signoret J. (2003): Suivi des populations de *Lobaria pulmonaria* (L.) Hoffm. en Lorraine. Résultats préliminaires et premières applications à leur gestion conservatoire. – *Bulletin d'Information de l'Association Française de Lichénologie* 28: 7–15.
- Signoret J. (2004): Les lichens – la diversité lichénique, une richesse à sauvegarder. – *Espaces Naturels* 6: 27–29.
- Signoret J. & Diederich P. (2003): Inventaire des champignons lichénisés et lichénicoles de la Réserve Naturelle des rochers et tourbières du Pays de Bitche. – *Annales Scientifiques de la Réserve de Biosphère Transfrontalière Vosges du Nord-Pfälzerwald* 11: 193–222.
- Signoret J., Van Haluwyn C., Ayrault S., Muller S. & Ferard J. F. (2002): Polluants atmosphériques et bioaccumulations métalliques: une hypothèse concernant l'impact de l'ozone. – *Air Pur* 63: 33–37.
- Sipman H., Leuckert C., Otte V., Knoph J.-G. & Rux K.-D. (2004): Die Flechten in Willdenows "Florae Berolinensis Prodrromus" und ihr Vorkommen im heutigen Berlin. – *Feddes Repertorium* 115: 121–133.
- Søchting U. (2004): *Flavoparmelia caperata* – a probable indicator of increased temperatures in Denmark. – *Graphis Scripta* 15: 53–56.
- Søchting U. & Frøberg L. (2003): *Caloplaca asserigena*, the correct name for a small lichen with a unique chemistry. – *Bibliotheca Lichenologica* 86: 47–51.
- Søchting U., Øvstedal D. O. & Sancho L. G. (2004): The lichens of Hurd Peninsula, Livingston Island, South Shetlands, Antarctica. – *Bibliotheca Lichenologica* 88: 607–658.
- Solhaug K. A. & Gauslaa Y. (2004): Photosynthates stimulate the UV-B induced fungal anthraquinone synthesis in the foliose lichen *Xanthoria parietina*. – *Plant, Cell and Environment* 27: 167–176.
- Solomina O. & Calkin P. E. (2003): Lichenometry as applied to moraines in Alaska, U.S.A., and Kamchatka, Russia. – *Arctic, Antarctic, and Alpine Research* 35: 129–143.
- Soukup D. & Türk R. (2003): Flechten der Murberge und der NW-Ausläufer der Gurktaler Alpen (Steiermark, Österreich). – *Fritschiana* 44: 1–26.
- Sparrius L. B. (2004): A monograph of *Enterographa* and *Sclerophyton*. – *Bibliotheca Lichenologica* 89: 1–141.
- Sparrius L. B. & Vervoort M. (2003): A contribution to the lichen flora of East Jutland (Denmark). – *Graphis Scripta* 14: 59–61.
- Stenroos S., Myllys L., Thell A. & Hyvönen J. (2002): Phylogenetic hypotheses: *Cladoniaceae*, *Stereocaulaceae*, *Baeomycetaceae*, and *Icmadophilaceae* revisited. – *Mycological Progress* 1: 67–282.
- Stepanenko L. S., Krivoshechkova O. E. & Skirina I. F. (2002): Functions of phenolic secondary metabolites in lichens from Far East Russia. – *Symbiosis* 32: 119–131.

- Sterner E. & Mayrhofer H. (2003): Ein Beitrag zur Diversität von lichenisierten Pilzen im Gebiet des Lachtals (Wölzer Tauern, Steiermark, Österreich). – *Fritschiana* 41: 1–19.
- Stevenson S. K. & Coxson D. S. (2003): Litterfall, growth, and turnover of arboreal lichens after partial cutting in an Engelmann spruce - subalpine fir forest in north-central British Columbia. – *Canadian Journal of Forest Research* 33: 2306–2320.
- Stocker-Wörgötter E. & Elix J. A. (2004): Experimental studies of lichenized fungi: formation of rare depsides and dibenzofurans by the cultured mycobiont of *Bunodophoron patagonicum* (*Sphaerophoraceae*, lichenized *Ascomycota*). – *Bibliotheca Lichenologica* 88: 659–669.
- Stocker-Wörgötter E., Elix J. A. & Grube M. (2004): Secondary chemistry of lichen-forming fungi: Chemosyndromic variation and DNA-analyses of cultures and chemotypes in the *Ramalina farinacea* complex. – *Bryologist* 107: 152–162.
- Stretch R. C. & Viles H. A. (2002): The nature and rate of weathering by lichens on lava flows on Lanzarote. – *Geomorphology* 47: 87–94.
- Süleyman H., Odabasoglu F., Aslan A., Cakir A., Karagöz Y., Gocer F., Halıcı M. & Bayir Y. (2003): Anti-inflammatory and antiulcerogenic effects of the aqueous extract of *Lobaria pulmonaria* (L.) Hoffm. – *Phytomedicine* 10: 552–557.
- Szczepańska K. (2004): Porosty rezerwatu „Nowa Morawa“ v Górach Bialskich (Sudety). – *Parki Narodowe i Rezerваты Przyrody* 23: 13–19.
- Takenaka Y., Tanahashi T., Nagakura N. & Hamada N. (2002): Production of anthraquinones by the cultured mycobionts of *Xanthoria polycarpa* and *X. elegans*. – *Lichenology* 1: 7–10.
- Thell A., Feuerer T., Kärnefelt I., Myllys L. & Stenroos S. (2004): Monophyletic groups within the *Parmeliaceae* identified by ITS rDNA, β -tubulin and GAPDH sequences. – *Mycological Progress* 3: 297–314.
- Thell A., Stenroos S., Feuerer T., Kärnefelt I., Myllys L. & Hyvönen J. (2002): Phylogeny of cetrarioid lichens (*Parmeliaceae*) inferred from ITS and β -tubulin sequences, morphology, anatomy and secondary chemistry. – *Mycological Progress* 1: 335–354.
- Thüs H. (2003): Flechten und die Stadtbiotopkartierung von Mainz. – *Mainzer Naturwissenschaftliches Archiv, Beihefte* 22: 218–236.
- Thüs H. & Dornes P. (2003): Neu- und Wiederfunde von Flechten in Hessen. – *Hessische Floristische Briefe* 52: 62–67.
- Tormo R., Recio D., Silva I. & Muñoz A. F. (2001): A quantitative investigation of airborne algae and lichen soredia obtained from pollen traps in south-west Spain. – *European Journal of Phycology* 36: 385–390.
- Torres A., Hochberg M., Pergament I., Smoum R., Niddam V., Dembitsky V. M., Temina M., Dor I., Lev O., Srebnik M. & Enk C. D. (2004): A new UV-B absorbing mycosporine with photo protective activity from the lichenized ascomycete *Collema cristatum*. – *European Journal of Biochemistry* 271: 780–784.
- Tretiach M., Crisafulli P., Virgilio D., Baruffo L. & Jensen M. (2003): Seasonal variation of photoinhibition in an epiphytic population of the lichen *Parmelia sulcata*. – *Bibliotheca Lichenologica* 86: 313–327.
- Tretiach M., Pinna D. & Grube M. (2003): *Caloplaca erodens* [sect. *Pyrenodesmia*], a new lichen species from Italy with an unusual thallus type. – *Mycological Progress* 2: 127–136.
- Türk R., Hafellner J. & Taurer-Zeiner C. (2004): Die Flechten Kärntens. Eine Bestandsaufnahme nach mehr als einem Jahrhundert lichenologischer Forschungen. – Verlag Naturwissenschaftlicher Verein für Kärnten, Sonderreihe Natur Kärnten 2, Klagenfurt. [333 pp.]
- Türk R. & Pfefferkorn-Dellali V. (2004): Über die Flechten im Naturschutzgebiet Rheindelta sowie neue und seltene Flechten in Vorarlberg (Austria). – *Linzer Biologische Beiträge* 36: 485–491.
- Türk R. & Uhl A. (2004): Die Verbreitung und Ökologie einiger ausgewählter Arten der Gattung *Lecidea* und von *Tremolecia atrata* in Österreich. – *Bibliotheca Lichenologica* 88: 691–699.
- van den Boom P. P. G. (2003): Contribution to the flora of Portugal, lichens and lichenicolous fungi III. – *Nova Hedwigia* 76: 157–171.
- van den Boom P. P. G. (2003): Some interesting records of lichens and lichenicolous fungi from The Netherlands. – *Österreichische Zeitschrift für Pilzkunde* 12: 123–127.
- van den Boom P. P. G. (2004): A long-term inventory of lichens and lichenicolous fungi of the Strabrechtse Heide and Lieropse Heide in Noord-Brabant, The Netherlands. – *Österreichische Zeitschrift für Pilzkunde* 13: 131–151.

- van den Boom P. P. G. & Brand A. M. (2004): *Micarea subcinerea*, an additional species of the lichen flora from Western Europe. – *Österreichische Zeitschrift für Pilzkunde* 13: 125–129.
- van den Boom P. P. G. & Khodosovtsev A. (2004): Notes on *Lecania* in eastern Europe and central Asia. – *Graphis Scripta* 16: 1–10.
- van Herk C. M., Mathijssen-Spiekman E. A. M. & de Zwart D. (2003): Long distance nitrogen air pollution effects on lichens in Europe. – *Lichenologist* 35: 347–359.
- van Herk K. & Aptroot A. (2004): Verspreidingspatronen en ecologie van Nederlandse korstmossen [Distribution patterns and ecology of lichens in the Netherlands]. – *Gorteria* 30: 77–91.
- Veldkamp J. F. (2004): *Bilimbia (Lichenes)* resurrected. – *Lichenologist* 36: 191–195.
- Villar S. E. J., Edwards H. G. M. & Seaward M. R. D. (2004): Lichen biodeterioration of ecclesiastical monuments in northern Spain. – *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 60: 1229–1237.
- Waldleigh M. A. (2003): Lichens and atmospheric sulphur: what stable isotopes reveal. – *Environmental Pollution* 126: 345–351.
- Walser J.-C. (2004): Molecular evidence for limited dispersal of vegetative propagules in the epiphytic lichen *Lobaria pulmonaria*. – *American Journal of Botany* 91: 1273–1276.
- Walser J.-C., Gugerli F., Holderegger R., Kuonen D. & Scheidegger C. (2004): Recombination and clonal propagation in different populations of the lichen *Lobaria pulmonaria*. – *Heredity* 93: 322–329.
- Warren J. (2003): Isolation by distance in the crustose lichens *Candelariella vitellina* and *Placynthium nigrum* colonising gravestones in northeast Scotland. – *Biodiversity and Conservation* 12: 217–224.
- Weckesser M. (2004): Stammepiphyten in einem Karpatenbirken-Fichten-Moorwald im Hochharz. – *Tuexenia* 24: 207–213.
- Wedin M., Döring H. & Gilenstam G. (2004): Saprotrophy and lichenization as options for the same fungal species on different substrata: environmental plasticity and fungal lifestyles in the *Stictis–Conotrema* complex. – *New Phytologist* 164: 459–465.
- Westberg M., Kärnefelt I. & Thell A. (2004): *Melanelia agnata*, an overlooked species, new to Sweden. – *Graphis Scripta* 16: 23–27.
- Wetmore C. M. (2004): The isidiate corticolous *Caloplaca* species in North and Central America. – *Bryologist* 107: 284–292.
- Wetmore C. M. (2004): The sorediate corticolous *Caloplaca* species in North and Central America. – *Bryologist* 107: 505–520.
- Winchester V. & Chaujar R. K. (2002): Lichenometric dating of slope movements, Nant Ffrancon, North Wales. – *Geomorphology* 47: 61–74.
- Wolinski H. (2003): Confocal imaging reveals structural detail of the cell nucleus and ascospore formation in lichenized fungi. – *Mycological Research* 107: 989–995.
- Yahr R., Vilgalys R. & DePriest P. T. (2004): Strong fungal specificity and selectivity for algal symbionts in Florida scrub *Cladonia* lichens. – *Molecular Ecology* 13: 3367–3378.
- Yenisoy Karakaş S. & Tuncel S. G. (2004): Comparison of accumulation capacities of two lichen species analyzed by instrumental neutron activation analysis. – *Journal of Radioanalytical and Nuclear Chemistry* 259: 113–118.
- Yenisoy-Karakaş S. & Tuncel S. G. (2004): Geographic patterns of elemental deposition in the Aegean region of Turkey indicated by the lichen, *Xanthoria parietina* (L.) Th. Fr. – *Science of the Total Environment* 329: 43–60.
- Yun M., Wadleigh M. A. & Pye A. (2004): Direct measurement of sulphur isotopic composition in lichens by continuous flow-isotope ratio mass spectrometry. – *Chemical Geology* 204: 369–376.
- Zalewska A. & Fałtynowicz W. (eds.) (2004): Lichens of the protected areas in the Euroregion Niemen. – „Man and Nature“ Association, Helios, Suwalki. [110 pp.]
- Zavada M. S., DiMichele L. & Toth C. R. (2004): The demi-lichenization of *Trametes versicolor* (L.:Fries) Pilat (*Polyporaceae*): the transfer of fixed ¹⁴CO₂ from epiphytic algae to *T. versicolor*. – *Northeastern Naturalist* 11: 33–40.
- Zhurbenko M. P. (2004): Lichenicolous and some interesting lichenized fungi from the Northern Ural, Komi Republic of Russia. – *Herzogia* 17: 77–86.