

- Rašeliniská Slovenska, p. 153-155, Daphne - Inštitút aplikovanej ekológie, Bratislava.
- Valachovič M. et Hájek M. (2000): Poznámky k výskytu a cenológii dvoch druhov rodu *Montia* na Slovensku. – Bull. Slov. Bot. Spoločn., Bratislava, 22: 165-169.
- Valachovič M. et Kochjarová J. (2000): *Cochlearia pyrenaica* – nový druh v Západných Karpatoch. – Preslia, Praha, 72: 475-493.
- Váňa J. (2001): K otázce biodiverzity bezečvných rastlín na Šumavě. – In: Mánek J., ed., Aktuality šumavského výzkumu, p. 27-28, ed. Správa NP a CHKO Šumava, Vimperk.
- Viceníková A. (2000): Ekologická charakteristika a klasifikácia vrchovísk. – In: Stanová V., ed., Rašeliniská Slovenska, p. 11-15, Daphne - Inštitút aplikovanej ekológie, Bratislava.
- Watzka R. (1999): Spoločstvá lužných jelšíň Lúbochnianskej doliny vo Veľkej Fatre. – Bull. Slov. Bot. Spoločn., Bratislava, 21: 151-160.
- Zlinská J. (2000): Vegetácia Holubyho lesostepi pri Vinosadoch v Malých Karpatoch. – Acta Envir. Univ. Comenianae, Bratislava, 10: 139-152.

NOVÁ LICHENOLOGICKÁ LITERATURA X.

sestavil Z. Palice

- Ahti T. (2000). Cladoniaceae. Flora Neotropica. – p. 78.-362, New York Bot. Garden, Bronx.
- Ahti T. et al. [eds.] (1999): Nordic Lichen Flora Vol. 1. Introductory parts, Calicioid lichens and fungi. – 94 p., Nordic Lichen Society, Uddevalla.
- Aptroot A. et van Herk C.M. (1999): *Bacidia neosquamulosa*, a new and rapidly spreading corticolous lichen species from western Europe. – Lichenologist 31: 121-127.
- Armstrong R.A. (2000): Competitive interactions between four foliose lichen species with and without nutrient enrichment. – Symbiosis 28: 323-335.
- Armstrong R.A. et Smith S.N. (1999): Factors associated with lobe division in the lichen *Parmelia conspersa* (Ehrh. ex Ach.) Ach. – Symbiosis 26: 265-277.
- Arseneau M.-J., Ouellet J.-P. et Sirois L. (1998): Fruticose arboreal lichen biomass accumulation in an old-growth balsam fir forest. – Can. J. Bot. 76: 1669-1676.
- Arseneau M.-J., Sirois L. et Ouellet J.-P. (1997): Effects of altitude and tree height on the distribution and biomass of fruticose arboreal lichens in an old growth balsam fir forest. – Ecoscience 4: 206-213.
- Articus K. (2000): Artbestämning av långa hängande skäggglavar (släktet *Usnea*) i Sverige [The long and pendant *Usnea* species in Sweden]. – Svensk Bot. Tidskr. 94: 81-100.
- Arvidsson L. (1999): A survey of lichenology in Sweden during the 19th century. – Symb. Bot. Upsal. 32/2: 23-60.
- Ascaso C. (2000): Lichens on rock substrates: observation of the biomineralization processes. – Bibl. Lichenol. 175: 127-135.
- Berger F. (2000): Beitrag zur Kenntnis der Flechten und lichenicolen Pilze Islands. – Acta Bot. Island. 13: 69-82.
- Berger F. et Priemetzhofer F. (2000): Neue und seltene Flechten und lichenicole Pilze aus Oberösterreich, Österreich III. – Herzogia 14: 59-84.
- Biazrov L.G. (1999): Indeks razvitiya epifitnykh lišajnikov i ocenka sostojaniya vozdušnogo bassejna territorii g. Moskvy. – Bjul. Mosk. Obšč. Ispitatelyj Prirody. Otd. Biol. 104/6: 31-40.
- Brady P.V. et al. (1999): Direct measurement of the combined effects of lichen, rainfall, and temperature on silicate weathering. – Geochim. Cosmochim. Acta 63: 3293-3300.

- Bridge P.D. et al. (1998): Applications of PCR in Mycology. – 357 p., CAB International, Wallingford.
- Bridge P.D. et Hawksworth D.L. (1998): What molecular biology has to tell us at the species level in lichenized fungi. – *Lichenologist* 30: 307-320.
- Burgaz A.R. et Martínez I. (1999): La familia Lobariaceae en la Península Ibérica. – *Bot. Complut.* 23: 59-90.
- Campanella L. et Ercole P. (1998): The lichens: from atmospheric pollution bioindicators to experimental model in order to define the tumours origin. – *Rec. Res. Developm. Pure and Appl. Analyt. Chem.* 1: 87-98.
- Chakir S. et Jensen M. (1999): How does *Lobaria pulmonaria* regulate photosystem II during progressive desiccation and osmotic water stress? A chlorophyll fluorescence study at room temperature and at 77 K. – *Physiol. Plant.* 105: 257-265.
- Cieślański S. et Fałtynowicz W. (1999): Atlas of the geographical distribution of lichens in Poland. Part 2. – 63 p., W. Szafer Institute of Botany, Polish Academy of Science, Kraków.
- Clark B.M. et al. (1999): Analysis of lichen thin sections by PIXE and STIM using a proton microprobe. – *Nucl. Instrum. Meth. Phys. Res., ser. B*, 150: 248-253.
- Clayden S.R. (1998): Thallus initiation and development in the lichen *Rhizocarpon lecanorinum*. – *New Phytol.* 139: 685-695.
- Crespo A. et al. (1999): Comparison of rRNA genotype frequencies of *Parmelia sulcata* from long established and recolonizing sites following sulphur dioxide amelioration. – *Plant Syst. Evol.* 217: 177-183.
- Crespo A. et al. (1999): Técnicas y métodos para la iniciación en el estudio de la evolución molecular con aplicaciones especiales para el análisis de los hongos liquenizados. – *Bot. Complut.* 23: 13-51.
- Cuny D., van Haluwyn C. et Caron B. (1997): Stimulation of ascospore germination by fumaroprotocetraric acid in *Diploschistes muscorum* (Scop.) R. Sant. (lichenized Ascomycotina). – *Nova Hedwigia* 64: 103-110.
- Czarnota P. (2000): Porosty Gorczafiskiego Parku Narodowego. Część I. Wykaz i rozmieszczenie gatunków. – *Parki Narod. Rezer. Przyr.* 19: 3-73.
- Czarnota P. et Coppins B.J. (2000): A new species of *Agromyia* and some interesting lichens from Gorce Mts (Western Biskidy Mts) new to Poland. – *Graphis Scripta* 11: 56-60.
- Diederich P. et Sérusiaux E. (2000): The lichens and lichenicolous fungi of Belgium and Luxembourg. An annotated checklist. – 207 p., Musée National d'Histoire Naturelle, Luxembourg.
- Dietz S. et al. (2000): Transmittance of light through the cortex of lichens from contrasting habitats. – *Bibl. Lichenol.* 75: 171-182.
- Döring H. et al. (2000): Mycobiont-specific PCR primers for the amplification of nuclear ITS and LSU rDNA from lichenized ascomycetes. – *Lichenologist* 32: 200-204.
- Ekman S. et Wedin M. (2000): The phylogeny of the families Lecanoraceae and Bacidiaceae (lichenized Ascomycota) inferred from nuclear SSU rDNA sequences. – *Plant Biol.* 2: 350-360.
- Elix J.A. (1999): Detection and identification of secondary lichen substances: contributions by the Uppsala school of lichen chemistry. – *Symb. Bot. Upsal.* 32/2: 103-121.
- Esseen P.-A. et Renhorn K.-E. (1998): Mass loss of epiphytic lichen litter in a boreal forest. – *Ann. Bot. Fenn.* 35: 211-217.
- Figueira R. et al. (1999): Space-time geostatistical modelling: a case study of sea-salt measured on lichens. – In: Gómez-Hernández J., Soares A. et Froidevaux R. (eds.), *GeoEnv II – Geostatistics for environmental applications*, p. 53-64, Kluwer Academic Publishers, Dordrecht.
- Fryday A. (2000): On *Rhizocarpon obscuratum* (Ach.) Massal., with notes on some related species in the British Isles. – *Lichenologist* 32: 207-224.
- Gauslaa Y. et Solhaug K.A. (1999): High-light damage in air-dry thalli of the old forest lichen

- Lobaria pulmonaria – interactions of irradiance, exposure duration and high temperature. – *J. Exp. Bot.* 50: 697-705.
- Gauslaa Y. et Solhaug K.A. (2000): High-light-intensity damage to the foliose lichen *Lobaria pulmonaria* within a natural forest: the applicability of chlorophyll fluorescence methods. – *Lichenologist* 32: 271-289.
- Giordano S. et al. (1999): Toxic effects of the thallus of the lichen *Cladonia foliacea* on the growth and morphogenesis of bryophytes. – *Cryptogamie, Bryol.-Lichénol.* 20: 35-41.
- Giralt M., Barbero M. et Elix J.A. (2000): Notes on some corticolous and lignicolous *Buellia* species from the Iberian Peninsula. – *Lichenologist* 32: 105-128.
- Goward T. (1998): Observations on the ecology of the lichen genus *Bryoria* in high elevation conifer forests. – *Canad. Field-Natur.* 112: 496-501.
- Goward T. (1999): The lichens of British Columbia, Illustrated Keys. Part 2, Fruticose Species. – 319 p., British Columbia Ministry of Forests, Victoria.
- Goward T. et Arsenault A. (2000): Cyanolichen distribution in young unmanaged forests: a dripzone effect? – *Bryologist* 103: 28-37.
- Grasso M.F. et al. (1999): Lichens as bioindicators in volcanic areas: Mt. Etna and Vulcano Island (Italy). – *Environm. Geology* 37: 207-217.
- Grube M. (1999): Epifluorescence studies of the ascus in *Verrucariales* (lichenized Ascomycotina). – *Nova Hedwigia* 68: 241-249.
- Grube M. et Kroken S. (2000): Molecular approaches and the concept of species and species complexes in lichenized fungi. – *Mycol. Res.* 104: 1284-1294.
- Haas J.R., Bailey E.H. et Purvis O.W. (1998): Bioaccumulation of metals by lichens: uptake of aqueous uranium by *Peltigera membranacea* as a function of time and pH. – *Amer. Mineralogist* 83: 1494-1502.
- Halonen P. et al. (1998): Synopsis of the genus *Usnea* (lichenized Ascomycetes) in British Columbia, Canada. – *Bryologist* 101: 36-60.
- Hammer S. (2000): Meristem growth dynamics and branching patterns in the Cladoniaceae. – *Amer. J. Bot.* 87: 33-47.
- Hazell P. et Gustafsson L. (1999): Retention of trees at final harvest – evaluation of a conservation technique using epiphytic bryophyte and lichen transplants. – *Biol. Conserv.* 90: 133-142.
- Hertel H., Schwaiger J. et Vorwerk B. (2000): Die Flechtenflora der Staatsforste am Südrand Münchens, einst und jetzt. - *Hoppea* 61: 445-524.
- Hyvärinen M. et Crittenden P.D. (1998): Growth of the cushion-forming lichen, *Cladonia portentosa*, at nitrogen-polluted and unpolluted heathland sites. – *Environm. Exp. Bot.* 40: 67-76.
- Hyvärinen M., Halonen P. et Kauppi M. (1999): Habitat type and primary colonisation of annual shoots of conifer samplings by epiphytic lichens. – *Nordic J. Bot.* 19: 505-511.
- Insarov G.E., Semenov S.M. et Insarova I.D. (1999): A system to monitor climate change with epilithic lichens. – *Environm. Monitor. Assessm.* 55: 279-298.
- Kivistö L. (1998): Taxonomy of *Stereocaulon paschale* and allied species in Finland. – *Sauteria* 9: 25-36.
- Kivistö L. et Kuusinen M. (2000): Edge effects on the epiphytic lichen flora of *Picea abies* in middle boreal Finland. – *Lichenologist* 32: 37-398.
- Krüger O. et Daniels F.J.A. (1998): A short account on lichen succession on cut surfaces of Scots pines. – *Herzogia* 13: 231-233.
- Lange O.L. (2000): Die Lebensbedingungen von Bodenkrusten-Organismen: Tagesverlauf der Photosynthese einheimischer Erdflechten. – *Hoppea* 61: 423-443.
- Lee M.R. et Parsons I. (1999): Biomechanical and biochemical weathering of lichen-encrusted granite: textural controls on organic-mineral interactions and deposition of silica-rich layers. –

- Chem. Geol. 161: 385-397.
- Loppi S. et Pirintos S.A. (2000): Effect of dust on epiphytic lichen vegetation in the Mediterranean area (Italy and Greece). – Israel J. Plant Sci. 4: 91-95.
- Lumbsch H.T., Lindemuth R. et Schmitt I. (2000): Evolution of filamentous ascomycetes inferred from LSU rDNA sequence data. – Plant Biol. 2: 525-529.
- Marbach B. (2000): Corticole und lignicole Arten der Flechtengattung *Buellia* sensu lato in den Subtropen und Tropen. – Bibl. Lichenol. 74: 1-384.
- Marcelli M.P. et Seaward M.R.D. [eds.] (1998): Lichenology in Latin America: history, current knowledge and applications. – 179 p., CETESB, São Paulo.
- Martínez Moreno M.I. (1999): Taxonomía del género *Peltigera* Willd. (Ascomycetes Liquenizados) en la Península Ibérica y estudio de sus hongos liquenócolas. – Ruizia 15: 1-200.
- Meyer B. et Printzen C. (2000): Proposal for a standardized nomenclature and characterization of insoluble lichen pigments. – Lichenologist 32: 571-583.
- Molina M.C. et Crespo A. (2000): Comparison of development of axenic cultures of five species of lichen-forming fungi. – Mycol. Res. 104: 595-602.
- Motiejūnaitė J. (1999): Checklist of lichens and allied fungi of Lithuania. – Bot. Lithuan. 5: 251-269.
- Myllys L. et al. (1999): Applicability of ITS data in Roccellaceae (Arthoniales, Euascomycetes) phylogeny. – Lichenologist 31: 471-476.
- Nimis P.L. et Tretiach M. (1999): Itineria Adriatica - Lichens from the eastern part of the Italian peninsula. – Stud. Geobot. 18: 51-106.
- Nordin A. (2000): *Buellia chloroleuca* in the Nordic countries. – Graphis Scripta 11: 35-39.
- Nordin A. (2000): Taxonomy and phylogeny of *Buellia* species with pluriseptate spores (Lecanorales, Ascomycotina). – Symb. Bot. Ups. 33/1: 3-117.
- Olech M. (1998): Apophytes in the lichen Flora of Poland. - Phytocoenosis 10: 251-255.
- Ott S., Schröder T. et Jahns H.M. (2000): Colonization strategies and interactions of lichens on twigs. – Bibl. Lichenol. 175: 445-455.
- Parinkina O.M., Piin T.K. et Pereverznev V.N. (1998): Mineralization and humification of lichens under natural conditions of Kola Peninsula. – Eurasian Soil Sci. 31: 1107-1113.
- Paulsrød P. et Lindblad P. (1998): Sequence variation of the tRNA Leu^{UUR}tron as a marker for genetic diversity and specificity of symbiotic cyanobacteria in some lichens. – Appl. Environm. Microbiol. 64: 310-315.
- Pinna D., Salvadori O. et Tretiach M. (1998): An anatomical investigation of calcicolous endolithic lichens from the Trieste karst (NE Italy). – Plant Biosyst. 132: 183-195.
- Pirintos S.A. et al. (1998): Effects of grazing on epiphytic lichen vegetation in a Mediterranean mixed evergreen sclerophyllous and deciduous shrubland (northern Greece). – Israel J. Plant Sci. 46: 303-307.
- Platt J.L. et Spatafora J.W. (1999): A re-examination of generic concepts of baeomycetoid lichens based on phylogenetic analyses of nuclear SSU and LSU ribosomal DNA. – Lichenologist 31: 409-418.
- Platt J.L. et Spatafora J.W. (2000): Evolutionary relationships of nonsexual lichenized fungi: molecular phylogenetic hypotheses for the genera *Siphula* and *Thamnolia* from SSU and LSU rDNA. – Mycologia 92: 475-487.
- Prieto B., Edwards H.G.M. et Seaward M.R.D. (2000): A fourier transform-raman spectroscopic study of lichen strategies on granite monumentss. – Geomicrobiol. J. 17: 55-60.
- Printzen C. (1999): *Japewiella* gen. nov., a new lichen genus and a new species from Mexico. – Bryologist 102: 714-719.
- Printzen C. et Lumbsch H.T. (2000): Molecular evidence for the diversification of extant lichens in the Late Cretaceous and Tertiary. – Molec. Phylogen. Evol. 17: 379-387.
- Printzen C. et Tønsberg T. (1999): The lichen genus *Biatora* in northwestern North America. –

- Bryologist 102: 692-713.
- Prinzing A.J. (1999): Wind-acclimated thallus morphogenesis in a lichen (*Evernia prunastri*, *Parmeliaceae*) probably favored by grazing disturbances. – Amer. J. Bot. 86: 173-183.
- Ramstad S. et Hestmark G. (2000): Effective neighbourhoods for a saxicolous lichen. – Mycol. Res. 104: 198-204.
- Reyes A., Molina M.C. et Vicente C. (2000): Polymerase chain reaction (PCR) primers for amplifying ribulose-1,5-bisphosphate carboxylase from lichens. – J. Hattori Bot. Lab. 88: 61-67.
- Richardson D.H.S. (1999): War in the world of lichens: parasitism and symbiosis as exemplified by lichens and lichenicolous fungi. – Mycol. Res. 103: 641-650.
- Scheidegger C. et Axelrod E. (2000): Low-frequency acoustic radiation during desiccation of *Lobaria pulmonaria*. – Bibl. Lichenol. 175: 161-170.
- Schöller H. [ed.] (1997): Flechten. Geschichte, Biologie, Systematik, Ökologie, Naturschutz und kulturelle Bedeutung. – 247 p., W. Kramer, Frankfurt am Main.
- Scholz P. (2000): Katalog der Flechten und flechtenbewohnenden Pilze Deutschlands. – Schriftenr. Vegetationsk. 31: 1-298.
- Schulz F. et Schroeter B. (2000): The coupling of microclimate, CO₂-exchange and activity: a case study on two *Parmelia* species in Swedish Lapland. – Bibl. Lichenol. 175: 353-364.
- Sipman H. et Raus T. (1999): A lichenological comparison of the Paros and Santorini island groups (Aegean, Greece), with annotated checklist. – Willdenowia 29: 239-297.
- Smith E. et al. (1998): Intra-specific variation in the photosynthetic responses of cyanobiont lichens from contrasting habitats. – New Phytol. 138: 213-224.
- Sommerkorn M. (2000): The ability of lichens to benefit from natural CO₂ enrichment under a spring snow-cover: a study with two arctic-alpine species from contrasting habitats. – Bibl. Lichenol. 175: 365-380.
- St. Clair L.L. (1999): A color guidebook to common Rocky Mountain lichens. – 242 p., M.L. Bean Life Science Museum of Brigham Young University, Provo, Utah.
- Stolley G. (2000): Zum Vorkommen von Flechten auf Plastik. – Bibl. Lichenol. 175: 457-464.
- Sundberg B. et al. (1999): Lichen respiration in relation to active time, temperaure, nitrogen and ergosterol concentrations. – Function. Ecol. 13: 119-125.
- Takahagi T. et al. (2000): Influence of NaCl concentration on spore germination and hyphal growth of lichens. – J. Hattori Bot. Lab. 88: 69-77.
- Thor G., Lücking R. et Matsumoto T. (2000): The foliicolous lichens of Japan. – Symb. Bot. Upsal. 32/3: 1-72.
- Tretiach M. et Hafellner J. (2000): Lichens and lichenicolous fungi of Mt. Fleons (Carnic Alps, Italy) I. Epilithic and epigaeic species. – Herzogia 14: 99-111.
- Trinkaus U. et Mayrhofer H. (2000): Revision der *Buellia epigaea*-Gruppe (lichenisierte Ascomyceten, Physciaceae). I. Die Arten der Nordhemisphäre. – Nova Hedwigia 71: 271-314.
- Türk R. et Berger F. (1999): Neue und seltene Flechten sowie lichenicole Pilze aus den Ostalpen III. – Linzer Biol. Beitr. 31: 929-953.
- Vokou D., Pirintos S.A. et Loppi S. (1999): Lichens as bioindicators of temporal variations in air quality around Thessaloniki, northern Greece. – Ecol. Res. 14: 89-96.
- Wedin M. et al. (2000): Small subunit rDNA phylogeny shows the lichen families Caliciaceae and Physciaceae (Lecanorales, Ascomycotina) to form a monophyletic group. – Can. J. Bot. 78: 246-254.
- Wetmore C.M. et Kärnefelt E.I. (1998): The lobate and subfruticose species of *Caloplaca* in north and central America. – Bryologist 101: 230-255.
- Wirth V. (1999): Gefährdete Flechtenbiotope in Mitteleuropa. – Natur und Museum 129: 12-21.
- Wolseley P.A. et Pryor K.V. (1999): The potential of epiphytic twig communities on *Quercus petraea* in a welsh woodland site (Tycanol) for evaluating environmental changes. –

- Lichenologist 31: 41-61.
- Wynn-Williams D.D., Holder J.M. et Edwards H.G.M. (2000): Lichens at the limits of life: past perspectives and modern technology. – Bibl. Lichenol. 175: 275-288.
- Yamamoto Y. et al. (1998): Factors affecting discharge and germination of lichen ascospores. – J. Hattori Bot. Lab. 85: 267-278.
- Zimmer D. (2000): Dauerbeobachtung der epiphytischen Flechtenvegetation unter immissionsökologischen Aspekten an Boden-Dauerbeobachtungsflächen in Schleswig-Holstein. – Bibl. Lichenol. 175: 231-251.
- Zoller S., Lutzoni F. et Scheidegger C. (1999): Genetic variation within and among populations of the threatened lichen *Lobaria pulmonaria* in Switzerland and implications for its conservation. – Molekul. Ecol. 8: 2049-2059.