

NOVÁ LICHENOLOGICKÁ LITERATURA III.

cestavil J. Liška

- Adamo P., Marchettiello A. & Violante P. (1993): The weathering of mafic rocks by lichens. - *Lichenologist* 25: 285-297.
- Aguirre-Hudson B. (1991): A taxonomic study of the species referred to the ascomycete genus *Leptoraphis*. - *Bull. Brit. Mus. (Natur. Hist.) Botany* 21: 85-192.
- Ahti T. (1993): Names in current use in the Cladoniaceae (Lichen-forming Ascomycetes) in the ranks of genus to variety. - *Regnum Veget.* 128: 59-102.
- Alstrup V. (1992): *Weddelomyces aspiciliicola*, a new lichenicolous fungus found in Czechoslovakia. - *Folia Geobot. Phytotax.* 27: 185-187.
- Alstrup V. (1992): Ceased grazing: the consequences of overgrowing on the flora of soil and stone inhabiting lichens. - *Int. J. Mycol. Lichenol.* 5: 1-2.
- Alstrup V. & Olech M. (1990): Additions to the lichen flora of the Polish Tatra Mountains. II. - *Zesz. Nauk. Univ. Jagiellonsk.* - Pr. Bot. 968/21: 211-217.
- Alstrup V. & Olech M. (1992): Additions to the lichen flora of the Polish Tatra Mountains. III. - *Zesz. Nauk. Univ. Jagiellonsk.* - Pr. Bot. 1059/24: 179-183.
- Alstrup V. & Olech M. (1992): Checklist of the lichens of the Tatra National Park, Poland. - *Zesz. Nauk. Univ. Jagiellonsk.* - Pr. Bot. 1059/24: 185-206.
- Armstrong R.A. (1992): A comparison of the foliose lichen *Parmelia conspersa* determined by a cross-sectional study and by direct measurement. - *Environm. Exp. Bot.* 32: 221-227.
- Arup U. et al. (1993): High performance thin layer chromatography (HPTLC), an improved technique for screening lichen substances. - *Lichenologist* 25: 61-71.
- Berger F. & Türk R. (1991): Zur Kenntnis der Flechten und Flechtenbewohnenden Pilze von Oberösterreich und Salzburg III. - *Linzer Biol. Beitr.* 23: 425-453.
- Berger F. & Türk R. (1993): Neue und seltene Flechten und lichenicole Pilze aus Oberösterreich, Österreich. - *Linzer Biol. Beitr.* 25: 167-204.
- Berger F. & Türk R. (1993): Bemerkenswerte Flechtenfunde aus dem Donautal zwischen Passau und Aschach (Oberösterreich, Österreich). - *Herzogia* 9: 669-681.
- Bjazrov L.G. (1991): Changes of epiphytic lichen species composition in a deciduous forest in Moscow region during 21 years and some problems of lichen use in environmental monitoring. - In: Krivoluckij D.A., ed., Bioindication and monitoring, p. 101-108, Nauka Press, Moscow.
- Brackel W. v. (1993): Die Flechten- und Moos-Gesellschaften Süddeutschlands. - *Veröffentlichungen des Bundes des Ökologen Bayerns* 6: 1-63.
- Burgaz A.R. (1992): Bibliografía botánica Ibérica. 1990. Lichenes. - *Botanica Complutensis* 17: 218-220.
- Burgaz A.R., Escudero A. & Ahti T. (1993): Morphometric variation in primary squamules of *Cladonia foliacea* and *C. convoluta*. - *Nova Hedwigia* 57: 231-238.
- Clerc P., Scheidegger Ch. & Ammann K. (1992): Liste rouge des macrolichens de la Suisse. - *Bot. Helvet.* 102: 71-83.
- Coppins B.J., James P.W. & Hawksworth D.L. (1992): New species and combination in the lichen flora of the Great Britain and Ireland. - *Lichenologist* 24: 351-369.
- Daniels F.J.A. & Harkema M. (1992): Epilithic lichen vegetation on man-made, calcareous substrates in The Netherlands. - *Phytocenologia* 21: 209-235.
- DePriest P.T. & Been M.D. (1992): Numerous group I introns with variable distributions in the ribosomal DNA of a lichen fungus. - *J. Molecul. Biol.* 228: 315-321.

- Dobson F.S. (1992): Lichens. An illustrated guide to the British and Irish species. - 376 p., Richmond Publishing Co., Slough.
- Egea J.M. & Torrente P. (1993): The lichen genus *Bactrospora*. - *Lichenologist* 25: 211-255.
- Elix J.A. (1993): Progress in the generic delimitation of *Parmelia* sensu lato lichens (Ascomycotina: Parmeliaceae) and a synoptic key to the *Parmeliaceae*. - *Bryologist* 96: 359-383.
- Farkas E. & Sipman H.J.M. (1993): Bibliography and checklist of foliicolous lichenized fungi up to 1992. - *Tropical Bryology* 7: 93-148.
- Feige G.B. & Lumbsch H.T., eds. (1993): Phytochemistry and chemotaxonomy of lichenized Ascomycetes - A festschrift in honour of Siegfried Huneck. - *Biblioth. Lichenol.* 53: 1-288.
- Feige G.B. et al. (1993): Identification of lichen substances by a standardized high-performance liquid chromatographic method. - *J. Chromatogr.* 646: 417-427.
- Fil'son R. (1993): What is a lichen? - *Australasian Lichenol. Newslett.* 32: 1-3.
- Föllmann G. & Schulz M. (1993): Stress-induced changes in the secondary products of lichen thalli. - *Bibl. Lichenol.* 53: 75-86.
- Foucard T. (1992): Notes on the corticolous *Arthopyrenia*-species in Sweden. - *Graphis Scripta* 4: 49-60.
- Fröberg L., Baur A. & Baur B. (1993): Differential herbivore damage to calcicolous lichens by snails. - *Lichenologist* 25: 83-95.
- Galloway D.J. (1992): Checklist of New Zealand lichens. - DSIR Land Resources Report 26: 1-58.
- Galloway D. (1992): Biodiversity: a lichenological perspective. - *Biodiversity and Conservation* 1: 312-323.
- Gehrman C.K., Krumbein W.E. & Petersen K. (1992): Endolithic lichens and the corrosion of carbonate rocks - a study of biopitting. - *Int. J. Mycol. Lichenol.* 5: 37-48.
- Gnüchtel A. & Müller F. (1993): Zur Flechtenflora Sachsen (3. Beitrag). - *Sächs. Florist. Mitt.* 2/1992-93: 43-51.
- Golubkova N.S., ed. (1992): Actual problems in experimental lichenology in the USSR. - 102 p., Komarov Bot. Inst., Leningrad.
- Gowan S.P. & Ahti T. (1993): Status of the lichen genus *Porpidia* in eastern Fennoscandia. - *Ann. Bot. Fennici* 30: 53-75.
- Gustafsson L. et al. (1992): Factors of importance to some lichen species of deciduous broad-leaved woods in southern Sweden. - *Lichenologist* 24: 255-266.
- Hafellner J. (1993): Die Gattung *Pyrrhospora* in Europa. Eine erste Übersicht mit einem Bestimmungsschlüssel der Arten nebst Bemerkungen zu einigen ausereuropäischen Taxa (lichenisierte Ascomycotina, Lecanorales). - *Herzogia* 9: 725-747.
- Hansson L., ed. (1992): Ecological principles of nature conservation. Applications in temperate and boreal environments. - 436 p., Elsevier Applied Science, London & New York.
- Hocke B. & Daniels F. (1993): Über die epilithische Flechtenflora und -vegetation im Stadtgebiet von Münster. - *Natur & Heimat* 53: 41-54.
- Hofmann P. (1993): Die epiphytische Flechtenflora und -vegetation des östlichen Nordtirol unter Berücksichtigung immissionsökologischer Gesichtspunkte. - *Bibl. Lichenol.* 51: 1-299.
- Jacobsen P. et al. (1992): Bearbeitungsstand und methodische Probleme bei der Erstellung von Roten Listen gefährdeter Flechten. - *Schriftenr. Vegetationskde* 23: 113-121.
- Jørgensen P.M. (1991): On some Fennoscandian *Pannaria* species. - *Ann. Bot. Fenn.* 28: 87-91.

- Kappen L. (1993): Flechten. Algen als Partner oder Opfer? - Naturwiss. Rundschau 46: 260-267.
- Kärnfeldt I., Mattsson J.-E. & Thell A. (1992): Evolution and phylogeny of cetrarioid lichens. - Plant Syst. Evol. 183: 113-160.
- Kärnfeldt I., Mattsson J.-E. & Thell A. (1993): The lichen genera Arctocetraria, Cetraria and Cetrariella (Parmeliaceae) and their presumed evolutionary affinities. - Bryologist 96: 394-404.
- Kärnfeldt I. & Thell A. (1992): The evaluation of characters in lichenized families, exemplified with the alectoroïd and some parmelioid genera. - Plant Syst. Evol. 180: 181-204.
- Kärnfeldt I. & Thell A. (1993): Chemical evolution in cetrarioid lichens. - Bibl. Lichenol. 53: 115-127.
- Kinder M., Vagts I. & Müller J. (1992): Auswirkung der Flechtenvegetation auf die Sukzession eines Sandtrockenrasens. - Int. J. Mycol. Lichenol. 5: 67-76.
- Kondratjuk S.J. & Beznc N.G. (1990): Peculiarities of distribution of toxitolerant lichen *Lecanora conizaeoides* Nyl. ex Cromb in the Ukraine. - Ukrainsk. Bot. Žurn. 47: 33-36.
- Kondratjuk S.J. & Solonina E.F. (1990): *Scoliciosporum chlorococcum* (Graewe ex Stenhammar) Věžda, a toxitolerant lichen of the Ukraine. - Ukrainsk. Bot. Žurn. 47: 37-41.
- Laundon J. (1992): Lepraria in the British Isles. - Lichenologist 24: 315-350.
- Letrouit-Galinou M.-A., Seaward M.R.D. & Deruelle S. (1992): A propos du retour des lichens épiphytes dans le Jardin du Luxembourg (Paris). - Bull. Soc. Bot. Fr., Paris, 139, Lettres bot. 2: 115-126.
- Litterski B. (1993): Die Flechten der Insel Rügen. - Herzogia 9: 415-474.
- Masuch G. (1993): Biologie der Flechten. - 411 p., Quelle & Meyer Verlag, Heidelberg & Wiesbaden.
- Mattsson J.-E. (1993): A monograph of the genus *Vulpicida* (Parmeliaceae, Ascomycetes). - Opera Bot. 119: 1-61.
- Mattsson J.-E. & Ming-J. Lai (1993): *Vulpicida*, a new genus in Parmeliaceae (lichenized Ascomycetes). - Mycotaxon 46: 425-428.
- Mayrhofer H., Scheidegger Ch. & Sheard J.W. (1992): On the taxonomy of five saxicolous species of the genus *Rinodina* (lichenized Ascomycetes). - Nordic J. Bot. 12: 451-459.
- Moberg R. & Holmasen I. (1992): Flechten von Nord- und Mitteleuropa. - 237 p., G.Fischer Verlag, Stuttgart, Jena & New York.
- Moreno P.P. & Egea J.M. (1992): Estudios sobre el complejo Anema-Thyre-Peccania en el sureste de la Península Ibérica y norte de África. - Acta Bot. Barcinon. 41: 1-66.
- Nimis P.L. (1992): Chiavi analitiche del genere *Caloplaca* Th.Fr. in Italia. - Notiziario Soc. Lichenol. Ital. 5: 9-28.
- Nimis P.L. (1993): The lichens of Italy, an annotated catalogue. - 897 p., Museo Regionale di Scienze Naturali, Torino.
- Papastefanou C. & Manolopoulou M. (1992): Residence time and uptake rates of ¹³⁷Cs in lichens and mosses at temperate latitude (40°N). - Environm. Int. 18: 397-401.
- Paus S. (1992): Der *Cladonia chlorophaea*-Komplex im Westmünsterland, Deutschland. - Int. J. Mycol. Lichenol. 5: 99-112.
- Pirintos S.A. et al. (1993): An assessment of the sampling procedure for estimating air pollution using epiphytic lichens as indicators. - Lichenologist 25: 165-173.
- Poelt J. & Hinteregger E. (1993): Beiträge zur Kenntnis der Flechtenflora des Himalaya. VII. Die Gattungen *Caloplaca*, *Fulgensia* und *Ioplaca* (mit englischem Bestimmungsschlüssel). - Bibl. Lichenol. 50: 1-247.

- Poelt J. & Leuckert C. (1993): Substitution and supplementary addition of secondary products in the evolution of lichenized Ascomycotina. - *Bibl. Lichenol.* 53: 201-215.
- Purvis O.W., Coppins B.J. & James P.W. (1993): Checklist of lichens of Great Britain and Ireland. - *Brit. Lichen Soc. Bull.* 72 Suppl.: 1-75.
- Rambold G., Triebel D. & Hertel H. (1993): Icmadophilaceae, a new family in the Leotiales. - *Bibl. Lichenol.* 53: 217-240.
- Randlane T. & Saag A. (1993): World list of cetrarioid lichens. - *Mycotaxon* 47: 395-403.
- Ropin K. & Mayrhofer H. (1993): Zur Kenntnis corticoler Arten der Gattung *Rinodina* (lichenisierte Ascomyceten) in den Ostalpen und angrenzenden Gebieten. - *Herzogia* 9: 779-835.
- Ruoss E. (1990): Rentierflechten im Alpenraum. - *Mitt. Naturforsch. Ges. Luzern* 31: 59-80.
- Ruoss E. (1992): Flechten im Kanton Luzern, Untersuchungen zur Bioindikation und Floristik, sowie zur Immissionsökologie voralpiner Hochmoore. - *Veröff. Natur-Mus. Luzern* 3: 1-98.
- Salvadori O., Pinna D. & Grillini G.C. (1990): Deterioration induced by lichens on volcanite of the Vulsini complex (central Italy). - In: Charola A.E., ed., *Preprints of the contributions to the international meeting Lavas and volcanic tuffs*, p. 215-225, Dirección de bibliotecas, Archives y Museos.
- Santesson R. (1993): The lichens and lichenicolous fungi of Sweden and Norway. - 240 p., Lund.
- Scheidegger C. (1993): A revision of European saxicolous species of the genus *Buellia* de Not. and formerly included genera. - *Lichenologist* 25: 315-364.
- Scholz P. (1992): Synopse der Roten Listen Flechten. - *Schriftenr. Vegetationskde* 23: 73-111.
- Scholz P. (1992): Karten zur Pilzverbreitung in Ostdeutschland. 9. Serie: Die Baeomycetaceae und Umbilicariaceae. - *Boletus* 16: 76-94.
- Scholz P., ed. (1993): Flechtenmonitoring - ein kommunales Kontrollinstrument. Tagung des Unabhängigen Institutes für Umweltfragen. - 140 p., UfU, Halle.
- Schuhwerk F. (1992): Die Berücksichtigung der Ökologie in der Lichenometrie: Datierung mit Sukzessionsstadien von Flechtengesellschaften. - Stuttgart. *Geogr. Stud.* 117: 161-175.
- Sloof J.E. & Wolterbeek B.T. (1993): Substrate influence on epiphytic lichens. - *Environm. Monitor. Assessm.* 25: 225-234.
- Tibell L. (1992): Crustose lichens as indicators of forest continuity in boreal coniferous forests. - *Nordic J. Bot.* 12: 427-450.
- Trass H. (1992): Synopsis of the lichen genus *Heterodermia* (Ascomycotina, Physciaceae sive Pyxinaceae). - *Folia Cryptogam. Eston.* 29: 1-24.
- Tynnirinen S. et al. (1992): Comparison of several bioindicator methods in monitoring the effects on forest of a fertilizer plant and a strip mine. - *Ann. Bot. Fenn.* 29: 11-24.
- Valladares F. & Sancho L.G. (1993): Biología de las comunidades líquénicas de los posaderos rocosos de aves en el Sistema Central español. - *Rivasgodaya* 7: 5-68.
- Virtala M. (1992): Optimal harvesting of a plant-herbivore system: lichen and reindeer in northern Finland. - *Ecol. Modelling* 60: 233-255.
- Wirth V. (1990): Neufunde von Flechten in Baden-Württemberg und anderen Regionen Deutschlands. - *Herzogia* 8: 305-334.
- Wirth V. (1991): Zeigerwerte von Flechten. - In: Ellenberg H. et al., eds., *Zeigerwerte von Pflanzen in Mitteleuropa*, p. 215-237, E. Goltze KG, Göttingen.