

- Jelínková G. (1995): Lišejníky v českých botanických textech posledních 150 let. - 61 p., Ms. [Dipl. pr., Praha]
- Kožená R. (1992): Příspěvek k poznání rozšíření rodu *Cladonia* vybraných chráněných území ChKV Slavkovský les. - 60 p., Ms [Dipl. pr., Plzeň]
- Łackovičová A. (1981): Epifytické lišajníky a čistota ovzdušia v južnej časti Malých Karpát. - 187 p., Ms. [Kand. disert. pr., Bratislava]
- Lisická E. (1977): Čelaď Umbilicariaceae Fée v ČSSR. - 230 p. + Príl. (sine pag.), Ms. [Kand. disert. pr., Bratislava]
- Máderová J. (1993): Lišejníky. - 23 p., Ms. [Bakal.pr., Praha]
- Sluková J. (1969): Epifytická lišejníková společenstva vrchu "Peklo" u Kašperských Hor. - 64 p., Ms [Dipl. pr., Plzeň]

## NOVÁ BRYOLOGICKÁ LITERATURA IV.

sestavil Z. Soldán

- Akiyama H. (1994): Allozyme variability within and among populations of the epiphytic moss *Leucodon* (Leucodontaceae: Musci). - *Am. J. Bot.* 81: 1280-1287.
- Akiyama H. (1994): Suggestion for the delimitation of the Leucodontaceae and the infrageneric classification of the genus *Leucodon*. - *J. Hattori Bot. Lab.* 76: 1-12.
- Asakawa Y. (1994): Chemosystematics of Hepaticae. - *J. Hattori Bot. Lab.* 76: 293-312.
- Ashton N.W. et al. (1994): Molecular phylogenetic aspects of genetical and physiological studies of moss development. - *J. Hattori Bot. Lab.* 76: 41-58.
- Bajon C. et al. (1995): In situ detection of mRNAs in the nucleus of the motile spermatozoid in a bryophyte (*Marchantia polymorpha*). - *Cryptogamie, Bryol. Lichénol.* 16: 1-10.
- Bakken S. (1995): Regional variation in nitrogen, protein and chlorophyll concentration in *Dicranum majus* - a reciprocal transplantation experiment. - *J. Bryol.* 18: 425-437.
- Basile M.R. & Basile D.V. (1994): The role of growth suppression in leafy liverwort morphogenesis and phylogeny. - *J. Hattori Bot. Lab.* 76: 75-82.
- Becker H. (1994): Secondary metabolites from bryophytes in vitro cultures. - *J. Hattori Bot. Lab.* 76: 283-292.
- Blackstock T.H. (1995): *Nardia insecta* Lindb., an addition to the liverwort flora of Britain, with cytological observations and comparison with *N. geoscyphus* (De Not.) Lindb. - *J. Bryol.* 18: 485-492.
- Buzas G. (1995): Bestimmungsschlüssel für die mitteleuropäischen Arten des *Barbula-Didymodon* Komplexes (inklusive *Bryoerythrophyllum*, *Geheebia* und *Pseudocrossidium*). - *Bryol. Rundbriefe* 22: 4-8.
- Convey P. (1994): Photosynthesis and dark respiration in Antarctic mosses: an initial comparative study. - *Polar Biology* 14(1): 65-69.
- Crosby M.R. & Magill R.E. (1994): Index to mosses 1990-1992. - *Monographs Syst. Bot. Missouri Bot. Garden* 50: 1-87.
- Daroczy S. et al. (1994): Subsequent mapping of 137Cs fallout from Chernobyl in Hungary using the radioactivity found in mosses. - *Naturwissenschaften* 81: 175-177.
- Delgadillo M.C. (1994): Endemism in the neotropical moss flora. - *Biotropica* 26(1): 12-16.
- Delgadillo M.C., Bello B. & Cárdenas S.S. (1995): LATMOSS, a catalogue of neotropical mosses. - 191 p., Mexico.
- Demidova E.E. & Filin V.R. (1994): False columella and spore release in *Tetraplodon angustatus* (Hedw.) Bruch et Schimp. in B.S.G. and *T. mnioides* (Hedw.) Bruch et Schimp. in B.S.G. (Musci: Splachnaceae). - *Arctoa* 3: 1-6.
- Duckett J.G. (1994): Studies of protonemal morphogenesis in mosses V. *Diphyscium foliosum* (Hedw.) Mohr. (Buxbaumiales). - *J. Bryol.* 18: 223-238.

- Duckett J.G. (1994): Studies of protonemal morphogenesis in mosses VI. The foliar rhizoids of *Calliargon stramineum* (Brid.) Kindb. function as organs of attachment. - *J. Bryol.* 18: 239-252.
- Duckett J.G. & Matcham H.W. (1995): Studies of protonemal morphogenesis in mosses VII. The perennial rhizoids and gemmiferous protonema of *Dicranella heteromalla* (Hedw.) Schimp. - *J. Bryol.* 18: 407-424.
- Felix H. (1994): Calli, cell and plantlet suspension cultures of bryophytes. - *Candollea* 49: 141-158.
- Flatberg K.I. (1994): Norwegian Sphagna. A field colour guide. - University Trondheim Vitensk. Museum Report Botanisk, Bot. ser. 1994(3): 1-42.
- Frahm J.-P. (1993): Zur Untersuchung und Lektotypifikation von *Marsupella aquatica* (Hepaticae, Marsupellaceae). - *Fragm. Flor. Geobot.* 38(2): 489-501.
- Frahm J.-P. (1994): A new synopsis of the *Campylopus* species from Australia. - *J. Bryol.* 18: 311-327.
- Frahm J.-P. (1994): A type catalogue of Campylopodioideae and Paraleucobryoideae (Musci, Dicranaceae). Part I. - *Fragm. Flor. Geobot.* 39: 85-101.
- Frahm J.-P. (1995): Lexikon der deutschen Bryologen. - *Limprichtia* 6: 1-187.
- Frey W., Frahm J.-P., Fischer E. & Lobin W. (1995): Die Moos- und Farnpflanzen Europas. - *Kleine Kryptogamenflora* 4., 6 ed., 426 p., Fischer Verlag, Stuttgart.
- Furuki T. & Mizutani M. (1994): Checklist of Japanese Hepaticae and Anthocerotae, 1993. - *Proceedings of the Bryol. Soc. of Japan* 6(5): 75-83.
- Gao C. et al. (1994): *Flora Bryophytarum Sinicorum* vol. 1. Sphagnales, Andreaeales, Archidiales, Dicranales. - 368 p., Science Press, Beijing.
- Goncalves E.P.R. et al. (1994): Seasonal variations of heavy metals in sediments and aquatic mosses from the Cavado river basin, Portugal. - *Sc. Total Environm.* 142(3): 143-156.
- Gos L. (1993): Two species of *Seligeria* (Musci, Seligeriaceae) new to Finland. - *Fragm. Flor. Geobot.* 38(2): 509-512.
- Greven H.C. (1995): *Grimmia* Hedw. (Grimmiaceae, Musci) in Europe. - 160 p., Backhuys Publishers, Leiden.
- Hartung W., Hellwege E.M. & Volk O.H. (1994): The function of abscisic acid in bryophytes. - *J. Hattori Bot. Lab.* 76: 59-66.
- Hasegawa J. (1994): New classification of Anthocerotae. - *J. Hattori Bot. Lab.* 76: 21-34.
- Hébrard J.-P. & Pierrot R.B. (1994): *Didymodon bistratosus* (Pottiaceae, Musci), espèce nouvelle du sud de l'Espagne (environs de Ronda, province Málaga). - *Nova Hedwigia* 59: 353-364.
- Hill M.O., Preston C.D. & Smith A.J.E. (1994): Atlas of bryophytes of Britain and Ireland 3. Mosses (Diplolepidaceae). - 419 p., Harley Books, Colchester.
- Hogg E.H., Malmer N. & Wallen B. (1994): Microsite and regional variation in the potential decay rate of Sphagnum magellanicum in south Swedish raised bogs. - *Ecography* 17(1): 50-59.
- Homm T. (1995): Zur Bestimmung von *Scleropodium caespitosum* (C.Müll.) L.Koch. - *Bryol. Rundbriefe* 22: 1-3.
- Hyvonen J. & Enroth J. (1994): Cladistic analysis of the genus *Pinatella* (Neckeraceae, Musci). - *Bryologist* 97(3): 305-312.
- Ingerpuu N. et al. (1994): Eesti sammalde nimestik - List of the Estonian bryophytes. - *The Naturalist's Notebook (Abiks Loodusevaatlajale)*, Tartu, 94: 1-175.
- Izumi Y. & Ono K. (1994): Pattern of the plastid division in spore mother cells of the hornwort *Anthoceros punctatus*. - *J. Plant Research* 107: 147-152.
- Knight C.D. (1994): Studying plant development in mosses: the transgenic route. - *Plant Cell and Environment* 17: 669-674.
- Kooijman A.M. & Baker C. (1994): The acidification capacity of wetland bryophytes as influenced by simulated clean and polluted rain. - *Aquatic Botany* 48: 133-144.
- Long D.G. (1994): Mosses of Bhutan II. A checklist of the mosses of Bhutan. - *J. Bryol.* 18: 339-364.

- Longton R.E. (1994): Genetic differentiation within the moss species *Polytrichum alpestre* Hoppe. - J. Hattori Bot. Lab. 75: 1-14.
- Longton R.E. (1994): Reproductive biology in bryophytes. The challenge and opportunities. - J. Hattori Bot. Lab. 76: 159-172.
- Marstaller R. (1994): Zur Verbreitung bemerkenswerter Moose in der Umgebung von Budapest (Ungarn). - Feddes Repert. 105: 531-547.
- Mischler B.D. et al. (1994) Phylogenetic relationships of the 'Green Algae' and 'Bryophytes'. - Annals Missouri Bot. Garden 81: 451-483.
- Moya J.J. et al. (1995): *Weissia papillosissima* Laz. (Pottiaceae, Musci), a species new to the European bryophyte flora. - J. Bryol. 18: 493-498.
- Müller F. (1995): *Riccia rhenana* Lorbeer in Europa nicht einheimisch? - Bryol. Rundbriefe 23: 6.
- Neinhuis C. & Jetter R. (1995): Ultrastructure and chemistry of epicuticular wax crystals in *Polytrichales* sporophytes. - J. Bryol. 18: 399-406.
- Newton A.E. & Mischler B.D. (1994): The evolutionary significance of asexual reproduction in mosses. - J. Hattori Bot. Lab. 76: 127-146.
- Ochrya R. (1994): Nomenclatural notes on mosses: 3. *Syntrichia virescens* (Pottiaceae). - Fragm. Flor. Geobot. 39: 311-314.
- Ochrya R., eds. (1995): *Munera Bryologica Georgio Szweykowski professori amicissimo et clarissimo septuagesimum vitae annum claudenti oblata*. - Fragm. Flor. Geobot. 40: 1-544.
- Ros R.M., Cano M.J. & Guerra J. (1994): *Barbula nana* C.Müll., a new synonym of *Trichostomum brachydontium* Bruch. - J. Bryol. 18: 375-376.
- Rydin H. (1995): Effects of density and water level on recruitment, mortality and shoot size in *Sphagnum* populations. - J. Bryol. 18: 439-453.
- Sastad S.M. & Flatberg K.I. (1994): Leaf size and shape in the *Sphagnum recurvum* complex: taxonomic significance and habitat variation. - J. Bryol. 18: 261-275.
- So M.L. (1995): Mosses and liverworts of Hong Kong. - 170 p., Hong Kong.
- Söderström L. (1994): Scope and significance of studies on reproductive biology of bryophytes. - J. Hattori Bot. Lab. 76: 97-104.
- Söderström L., ed. (1995): Preliminary distribution maps of bryophytes in Norden. Vol. 1. Hepaticae and Anthocerotae. - Mossornas Vänner, Trondheim.
- Söderström L. & Hedenäs L. (1994): Checklista över Sveriges Mossor Nyfynd och Rättelser. - Myrnia 4(2): 30-33.
- Takio S. (1994): Growth characteristics of bryophyte cell culture. - J. Hattori Bot. Lab. 76: 67-74.
- Tallis J.H. (1994): Pool-and-hummock patterning in a mire II. The formation and erosion of the pool system. - J. Ecol. 82: 789-803.
- Tallis J.H. & Livett E.A. (1994): Pool-and-hummock patterning in a southern Pennine blanket mire I. Stratigraphic profiles for the last 2800 years. - J. Ecol. 82: 775-788.
- Wojtuń B. (1994): Element contents of *Sphagnum* mosses of peat bogs of Lower Silesia (Poland). - Bryologist 97: 284-295.
- Woolgrove C.E. & Woodin S.J. (1994): Relationships between the duration of snowlie and the distribution of bryophyte communities within snowbeds in Scotland. - J. Bryol. 18: 253-260.
- Wyatt R. (1994): Population genetics of bryophytes in relation to their reproductive biology. - J. Hattori Bot. Lab. 76: 147-158.
- Wyatt R., Stoneburner A. & Odrzykoski I.J. (1994): Phylogenetic relationships within *Plagiomnium* section *Rosulata*. - J. Hattori Bot. Lab. 76: 87-96.
- Zechmeister H.G. (1995): Growth rates of five pleurocarpous moss species under various climatic conditions. - J. Bryol. 18: 455-468.